

4. Reduced ideals

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **36 (1990)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Then we have $GCD(a, b, c) = 1$, where $\frac{D - b^2}{4a} = c$, and so $GCD(a, bf, cf^2) = 1$,

showing that $I = \left[a, f \left(\frac{b + \sqrt{D}}{2} \right) \right]$ is a primitive ideal of $O_{D'}$. Hence C is the image of the class of I under θ .

COROLLARY 4. *If the class C of $O_{D'}$ contains the primitive ideal $I = \left[a, \frac{b + \sqrt{D'}}{2} \right]$, where $f^2 | a$, then $f | b$ and the class $\theta(C)$ contains the primitive ideal $J = \left[\frac{a}{f^2}, \frac{\frac{b}{f} + \sqrt{D}}{2} \right]$ of O_D .*

Proof. As $D' = Df^2 = b^2 + 4ac$, and $f^2 | a$, we see that $f | b$, and so $GCD(f, c) = 1$. By Corollary 2 we have $I = \left(\frac{\sqrt{D'} - b}{2a} \right) \left[c, \frac{-b + \sqrt{D'}}{2} \right]$ and so, by Theorem 1, we see that $\left[c, \frac{-\frac{b}{f} + \sqrt{D}}{2} \right] \in \theta(C)$. Finally, by Corollary 2, $J = \left[\frac{a}{f^2}, \frac{b/f + \sqrt{D}}{2} \right] = \frac{\left(\sqrt{D} + \frac{b}{f} \right)}{2c} \left[c, \frac{-\frac{b}{f} + \sqrt{D}}{2} \right]$, showing that $J \in \theta(C)$.

4. REDUCED IDEALS

From now on in this paper we suppose that $D_0 > 0$ so that we are only considering ideals in orders of a real quadratic field. An ideal I of O_D can be written in the form $I = ad[1, \phi]$, where $\phi = \frac{b + \sqrt{D}}{2a}$. By Proposition 1 (ii), if $I = a'd'[1, \phi']$ is another representation of I , then $a' = \pm a$ and $\phi' \equiv \frac{a}{a'} \phi \pmod{1}$. A real number of the form $\frac{b + \sqrt{D}}{2a}$, where $c = \frac{D - b^2}{4a}$ is an integer and $GCD(a, b, c) = 1$ is called a quadratic irrationality of discriminant D .

Definition 9. (Reduced number). The quadratic irrationality $\phi = \frac{b + \sqrt{D}}{2a}$

of discriminant D is said to be *reduced* if

$$(4.1) \quad \phi > 1, \quad -1 < \bar{\phi} < 0.$$

It is easy to check that (4.1) is equivalent to each of the inequalities in (4.2)

$$(4.2) \quad \begin{aligned} \text{(i)} \quad & 0 < \sqrt{D} - b < 2a < \sqrt{D} + b, \\ \text{(ii)} \quad & 0 < \sqrt{D} - b < 2c < \sqrt{D} + b. \end{aligned}$$

Moreover (4.2) implies

$$(4.3) \quad 0 < a < \sqrt{D}, \quad 0 < b < \sqrt{D}, \quad 0 < c < \sqrt{D}.$$

Definition 10. (Reduced ideal). The ideal $I = ad[1, \phi]$ of O_D , where $\phi = \frac{b + \sqrt{D}}{2a}$, is said to be *reduced* if, and only if, ϕ can be chosen to be reduced.

From (4.3) we see that the number of reduced, primitive ideals of O_D is finite.

PROPOSITION 4. ([12]: Definition and Theorem 3.5). *The ideal*

$$I = d \left[a, \frac{b + \sqrt{D}}{2} \right]$$

of O_D , where $a > 0$ and $d > 0$, is reduced if, and only if, I does not contain a nonzero element α satisfying $|\alpha| < da$, $|\bar{\alpha}| < da$.

Proof. It suffices to prove that I is reduced if, and only if, the Z -module $[1, \phi]$ does not contain a nonzero element $\lambda = x + y\phi$ such that

$$(4.4) \quad |\lambda| < 1, \quad |\bar{\lambda}| < 1.$$

If I is reduced we can suppose that $\phi > 1$, $-1 < \bar{\phi} < 0$. Let x and y be integers such that $0 < \lambda = x + y\phi < 1$.

Clearly we have $y \neq 0$. If $y \geq 1$, then we have $y\phi > 1$, so $x \leq -1$, showing that $\bar{\lambda} = x + y\bar{\phi} < -1$. If $y \leq -1$, then we have $y\phi < -1$, so $x \geq 2$, showing that $\bar{\lambda} = x + y\bar{\phi} > 2$. This proves that $[1, \phi]$ does not contain an element $\lambda \neq 0$ such that $|\lambda| < 1$, $|\bar{\lambda}| < 1$.

Now suppose the Z -module $[1, \phi]$ does not contain an element $\lambda \neq 0$ satisfying (4.4). We can choose ϕ so that $-1 < \bar{\phi} < 0$, in which case

$\phi = \bar{\phi} + \frac{\sqrt{D}}{a} > -1$. Hence, as ϕ cannot satisfy (4.4), we must have $\phi > 1$, so I is reduced.

LEMMA 4. If $I = d \left[a, \frac{b + \sqrt{D}}{2} \right]$ is an ideal of O_D with $0 < a < \frac{\sqrt{D}}{2}$ then I is reduced.

Proof. We can write $I = da[1, \phi]$ with $-1 < \bar{\phi} < 0$. Then we have $\phi = \bar{\phi} + \frac{\sqrt{D}}{a} > 1$ so that I is reduced.

5. LAGRANGE'S REDUCTION PROCEDURE

In this section we describe Lagrange's reduction procedure which was first introduced in [2]. This procedure uses Lagrange neighbours and so is based on the continued fraction algorithm. The procedure, when applied to a given primitive ideal I of O_D , gives all the reduced ideals of O_D which are equivalent to I .

Let $\{a, b\}$ be a representation of the primitive ideal I of O_D . The Lagrange neighbour of $\{a, b\}$ is the representation $\{a', b'\}$ of the primitive ideal I' of O_D given as follows:

$$(5.1) \quad \begin{cases} q = [\phi] = \left[\frac{b + \sqrt{D}}{2a} \right], & \phi = q + \frac{1}{\phi'}, \\ b' = -b + 2aq, & a' = \frac{D - b'^2}{4a} = \frac{D - b^2}{4a} + bq - aq^2, \end{cases}$$

(see (2.10) and (2.11)). We write $\{a, b\} \xrightarrow{L} \{a', b'\}$. The primitive ideal $I' = a'[1, \phi']$ is also called the Lagrange neighbour of I .

We note that

$$\phi' = \frac{1}{\phi - q} > 1, [\phi'] \geq 1,$$

as $q = [\phi]$. We also remark that if a is kept fixed and ϕ is changed modulo 1 then ϕ' , b' and a' do not change. Hence the Lagrange neighbour of $\{a, b\}$ depends only upon the sign of a . If $\{a, b\} \xrightarrow{L} \{a', b'\}$ then by Corollary 1 the