

Objekttyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **36 (1990)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

the Fourier transform $\hat{1}_E$ does not vanish on any nonempty open subset of the dual group \widehat{G} . Thus bounded Borel subsets of \mathbf{R}^n of positive Lebesgue measure are determining sets by the analyticity of $\hat{1}_E$. Classical quasi-analyticity results apply to give conditions on the growth of an unbounded subset $E \subseteq \mathbf{R}^n$ to be a determining set. Settling a problem that was open for some time, Kargaev ([17]) proved the existence of sets $E \subseteq \mathbf{R}^n$ of finite Lebesgue measure which are not determining sets for finite complex measures.

The problem of determining sets has also been studied with the class of probability measures replaced by other classes of measures, e.g., a class of infinite measures with growth/decay conditions (see [22], [11] and [28]). Also different groups of homeomorphisms acting on X have been considered in these studies.

Finally, we refer to the following form of the *support problem* analogous to the well known problem in the case of Radon transform. Let X be a symmetric space (Euclidean, compact or non-compact). Let x_0 be a fixed point of X . If E is a relatively compact subset of positive measure and if $\int_{gE} f = 0$ for all $g \in G$ with $d(x_0, gx_0) > R$ what can one say about the support of f with respect to the reference point x_0 ? (Here, d stands for the geodesic distance.) Some partial answers to this question are known (see [26] and [28]).

We have not addressed ourselves in this paper to the situation when X is an infinite-dimensional Hilbert space or X is an arbitrary Riemannian manifold. Another important problem we have not considered is the *local version* of the Pompeiu problem. (For this, we refer the reader to [5] and [6]). We have restricted ourselves to the situation of symmetric spaces and locally compact groups and the relationship of the Pompeiu problem with harmonic analysis and representation theory.

Acknowledgement. The authors thank Lawrence Zalcman for pointing out some errors in the original draft of this paper. They would also like to express their gratitude to K. R. Parthasarathy for initially getting them interested in the Pompeiu problem.

REFERENCES

- [1] BAGCHI, S. C. and A. SITARAM. Spherical mean periodic functions on semisimple Lie groups. *Pacific J. Math.* 84 (1979), 241-250.
- [2] BAGCHI, S. C. and A. SITARAM. Determining sets for measures on \mathbf{R}^n . *Illinois J. Math.* 26 (1982), 419-422.

- [3] BERENSTEIN, C. A. An inverse spectral theorem and its relation to the Pompeiu problem. *J. Analyse Math.* 37 (1980), 128-144.
- [4] —— Spectral synthesis on symmetric spaces. *Contemporary Math.* 63 (1987), 1-25.
- [5] BERENSTEIN, C. A. and R. GAY. Le problème de Pompeiu local. *Journal d'Analyse Mathématique*, LII (1989), 133-166.
- [6] BERENSTEIN, C. A. and R. GAY. A local version of the two circles theorem. *Israel J. Math.* 55 (1986), 267-288.
- [7] BERENSTEIN, C. A. and M. SHAHSHAHANI. Harmonic analysis and the Pompeiu problem. *Amer. J. Math.* 105 (1983), 1217-1229.
- [8] BERENSTEIN, C. A. and P. YANG. An inverse Neumann problem. To appear in *J. Reine Angew. Math.*
- [9] BERENSTEIN, C. A. and L. ZALCMAN. Pompeiu's problem on symmetric spaces. *Comment. Math. Helvetici* 55 (1980), 593-621.
- [10] BROWN, L. and J. P. KAHANE. A note on the Pompeiu problem for convex domains. *Math. Annalen* 259 (1982), 107-110.
- [11] BROWN, L., F. SCHNITZER and A. L. SHIELDS. A note on a problem of D. Pompeiu. *Math. Zeitschr.* 105 (1968), 59-61.
- [12] BROWN, L., B. M. SCHREIBER and B. A. TAYLOR. Spectral synthesis and the Pompeiu problem. *Ann. Inst. Fourier (Grenoble)* 23 (1973), 125-154.
- [13] EHRENPREIS, L. and F. MAUTNER. Some properties of the Fourier transform on semi-simple Lie groups I. *Ann. Math.* 61 (1955), 406-439.
- [14] EHRENPREIS, L. and F. MAUTNER. Some properties of the Fourier transform on semisimple Lie groups II. *Trans. Amer. Math. Soc.* 84 (1957), 1-55.
- [15] HELGASON, S. *Differential Geometry, Lie Groups and Symmetric Spaces*. Academic Press. 1978.
- [16] —— *Groups and Geometric Analysis*. Academic Press. 1984.
- [17] KARGAEV, P. P. The Fourier transform of the characteristic function of a set vanishing on an interval. *Math. USSR Sbornik* 45 (1983), 397-410.
- [18] POMPEIU, D. Sur une propriété des fonctions continues dépendant de plusieurs variables. *Bull. Sci. Math.* (2) 53 (1929), 328-332.
- [19] —— Sur une propriété intégrale des fonctions de deux variables réelles. *Bull. Sci. Acad. Royale Belgique* (5) 15 (1929), 265-269.
- [20] RANA, I. K. Determination of probability measures through group actions. *Zeitsch. Warsch.* 53 (1980), 197-206.
- [21] —— *Determination of probability measures through group actions*. Ph. D. Thesis, Indian Statistical Institute, 1979.
- [22] SAPAGOV, N. A. A uniqueness problem for finite measures in Euclidean spaces — Problems in the theory of probability distributions. *Zap. Naucen Sem. (LOMI) Leningrad* 41 (1974), 3-13.
- [23] SCHWARTZ, L. Théorie générale des fonctions moyennes-périodiques. *Ann. of Math.* (2) 48 (1947), 857-929.
- [24] SCOTT, D. and A. SITARAM. Some remarks on the Pompeiu problem for groups. *Proc. Amer. Math. Soc.* 104 (1988), 1261-1266.
- [25] SERRIN, J. A symmetry problem in potential theory. *Arch. Rational Mech. Anal.* 43 (1971), 304-318.
- [26] SHAHSHAHANI, M. and A. SITARAM. The Pompeiu problem in exterior domains in symmetric spaces. *Contemporary Math.* 63 (1987), 267-277.
- [27] SITARAM, A. A theorem of Cramér and Wold revisited. *Proc. Amer. Math. Soc.* 87 (1983), 714-716.
- [28] —— Fourier analysis and determining sets for Radon measures on \mathbf{R}^n . *Illinois J. Math.* 28 (1984), 339-347.

- [29] — Some remarks on measures on noncompact semisimple Lie groups. *Pacific J. Math.* 110 (1984), 429-434.
- [30] — On an analogue of the Wiener-Tauberian theorem for symmetric spaces of the non-compact type. *Pacific J. Math.* 133 (1988), 197-208.
- [31] WAWRZYNCZYK, A. Spectral analysis and mean-periodic functions on rank one symmetric spaces. *Bol. Soc. Mat. Mexicana* (2), 30 (1985), 15-29.
- [32] WEIT, Y. On Schwartz's theorem for the motion group. *Ann. Inst. Fourier (Grenoble)* 30 (1980), 91-107.
- [33] WILLIAMS, S. A. A partial solution of the Pompeiu problem. *Math. Annalen* 223 (1976), 183-190.
- [34] — Analyticity of the boundary for Lipschitz domains without Pompeiu property. *Indiana Univ. Math. J.* 30 (1981), 357-369.
- [35] ZALCMAN, L. Analyticity and the Pompeiu problem. *Arch. Rational Mech. Anal.* 47 (1972), 237-254.

(Reçu le 6 novembre 1989)

S. C. Bagchi

Division of Theoretical Statistics and Mathematics
 Indian Statistical Institute
 203, B.T. Road
 Calcutta — 700035 (India)

A. Sitaram

Division of Theoretical Statistics and Mathematics
 Indian Statistical Institute
 Bangalore Centre
 8th Mile, Mysore Road
 Bangalore — 560059 (India)

vide-leer-empty