

## §2. The Maass space

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.05.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

only on the discriminant  $D := r^2 - 4mn$  and the residue class  $r \pmod{2m}$ .

The Petersson scalar product on  $J_{k,m}^{\text{cusp}}$  is normalized by

$$\langle \phi, \psi \rangle = \int_{\Gamma_1^J \backslash \mathcal{H} \times \mathbb{C}} \phi(\tau, z) \overline{\psi(\tau, z)} \exp(-4\pi m y^2 / v) v^{k-3} du dv dx dy$$

$$(\tau = u + iv, z = x + iy).$$

For basic facts about Jacobi forms we refer to [9].

## §2. THE MAASS SPACE

### 2.1. RESULTS

Let  $F$  be a Siegel modular form of integral weight  $k$  on  $\Gamma_2$  and write the Fourier expansion of  $F$  in the form

$$(1) \quad F(Z) = \sum_{m \geq 0} \phi_m(\tau, z) e^{2\pi i m \tau'} \quad \left( Z = \begin{pmatrix} \tau & z \\ z & \tau' \end{pmatrix} \in \mathcal{H}_2 \right).$$

Using the injection

$$(2) \quad \Gamma_1^J \rightarrow \Gamma_2, \quad \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix}, ((\lambda, \mu), \kappa) \right) \mapsto \begin{pmatrix} a & 0 & b & \mu \\ \lambda' & 1 & \mu' & \kappa \\ c & 0 & d & -\lambda \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

where  $(\lambda', \mu') = (\lambda, \mu) \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , and the transformation formula of  $F$  it is easy to see that the functions  $\phi_m$  are in  $J_{k,m}$ . The expansion (1) is referred to as the Fourier-Jacobi expansion of  $F$ .

Thus for any  $m \in \mathbf{N}_0$  we obtain a linear map

$$(3) \quad \rho_m: M_k(\Gamma_2) \rightarrow J_{k,m}, \quad F \mapsto \phi_m.$$

Note that  $\rho_0$  is equal to the Siegel  $\Phi$ -operator.

We shall be interested in the case  $m = 1$ . For  $k$  odd,  $\rho_1$  is the zero map; in fact, any Jacobi form of odd weight and index one must vanish identically as is easily seen.

For  $k$  even,  $\rho_1$  was studied in detail by Maass [28, 29] who showed the existence of a natural map  $V: J_{k,1} \rightarrow M_k(\Gamma_2)$  such that the composite  $\rho_1 \circ V$  is the identity. More precisely, let  $\phi \in J_{k,1}$  with Fourier coefficients  $c(n, r)$  ( $n, r \in \mathbf{Z}; r^2 \leq 4n$ ) and for  $m \in \mathbf{N}_0$  define

$$(4) \quad (V_m \phi)(\tau, z) := \sum_{n, r \in \mathbf{Z}, r^2 \leq 4mn} \left( \sum_{d \mid (n, r, m)} d^{k-1} c\left(\frac{mn}{d^2}, \frac{r}{d}\right) \right) q^n \zeta^r$$

(if  $m = 0$ , the term  $\sum_{d \mid 0} d^{k-1} c(0, 0)$  on the right of (4) has to be interpreted

as  $\frac{1}{2} \zeta(1-k)$ ; note that  $V_1 \phi = \phi$ ). Using a more invariant definition

of  $V_m$  in terms of the action of a set of representatives for  $\Gamma_1 \setminus \{M \in \mathbf{Z}^{(2,2)} \mid \det M = m\}$  one checks that  $V_m \phi \in J_{k,m}$  [9, §4]. Put

$$(V\phi)(Z) := \sum_{m \geq 0} (V_m \phi)(\tau, z) e^{2\pi i m \tau'} \quad \left( Z = \begin{pmatrix} \tau & z \\ z & \tau' \end{pmatrix} \in \mathcal{H}_2 \right).$$

We denote by  $T_n (n \in \mathbf{N})$  the usual Hecke operators on  $M_k(\Gamma_2)$  resp.  $S_k(\Gamma_2)$  [12, IV; 1, II]; thus, if  $p$  is a prime,  $T_p$  resp.  $T_{p^2}$  correspond to the two generators

$$\Gamma_2 \begin{pmatrix} 1_2 & 0 \\ 0 & p1_2 \end{pmatrix} \Gamma_2 \text{ resp. } \Gamma_2 \text{diag}(1, p, p^2, p) \Gamma_2$$

of the local Hecke algebra of  $\Gamma_2$  at  $p$ . We denote by  $T_{J,n} (n \in \mathbf{N})$  the Hecke operators on  $J_{k,m}$  resp.  $J_{k,m}^{\text{cusp}}$  [9, §4].

**THEOREM 1.** (Maass [28, 29], Andrianov [2]). *Suppose that  $k$  is even. The map  $\phi \mapsto V\phi$  gives an injection  $J_{k,1} \rightarrow M_k(\Gamma_2)$  which sends cusp forms to cusp forms and is compatible with the action of Hecke operators. If  $p$  is a prime, one has  $T_p \circ V = V \circ (T_{J,p} + p^{k-2}(p+1))$  and  $T_{p^2} \circ V = V \circ (T_{J,p}^2 + p^{k-2}(p+1)T_{J,p} + p^{2k-2})$ .*

The image of  $J_{k,1}$  under  $V$  is called the Maass space and will be denoted by  $M_k^*(\Gamma_2)$ . One knows that  $M_k^*(\Gamma_2) = \mathbf{C}E_k^{(2)} \oplus S_k^*(\Gamma_2)$  where  $E_k^{(2)}$  is the Siegel-Eisenstein series of weight  $k$  on  $\Gamma_2$  and  $S_k^*(\Gamma_2) := M_k^*(\Gamma_2) \cap S_k(\Gamma_2)$ . Observe that  $\dim M_k^*(\Gamma_2) = \dim J_{k,1}$  grows linearly in  $k$  while  $\dim M_k(\Gamma_2)$  grows like  $k^3$ .

Note that Theorem 1 implies that  $M_k^*(\Gamma_2)$  is stable under all Hecke operators and that it is annihilated by the operator

$$(5) \quad \mathcal{C}_p := T_p^2 - p^{k-2}(p+1)T_p - T_{p^2} + p^{2k-2},$$

for every prime  $p$ .

Let  $F \in M_k(\Gamma_2)$  be a non-zero Hecke eigenform and denote by  $\lambda_n (n \in \mathbf{N})$  its eigenvalues under  $T_n$ . If  $p$  is a prime, we put

$$Z_{F,p}(X) := 1 - \lambda_p X + (\lambda_p^2 - \lambda_{p^2} - p^{2k-4})X^2 - \lambda_p p^{2k-3}X^3 + p^{4k-6}X^4$$

so that  $Z_{F,p}(p^{-s})$  ( $s \in \mathbb{C}$ ) is the local spinor zeta function of  $F$  at  $p$ . We put

$$Z_F(s) := \prod_p Z_{F,p}(p^{-s}) \quad (\operatorname{Re}(s) \geq 0).$$

One has

$$Z_F(s) = \zeta(2s - 2k + 4)^{-1} \sum_{n \geq 1} \lambda_n n^{-s} \quad (\operatorname{Re}(s) \geq 0).$$

If  $F$  is an Eisenstein series, then it is well-known that  $Z_F(s)$  can be expressed in terms of products of Hecke  $L$ -functions of elliptic modular forms.

Suppose that  $F$  is cuspidal. Then it was proved in [1, Chap. 3] that  $Z_F(s)$  has a meromorphic continuation to  $\mathbb{C}$  which is holomorphic everywhere if  $k$  is odd and is holomorphic except for a possible simple pole at  $s = k$  if  $k$  is even. Moreover, the global function  $Z_F^*(s) := (2\pi)^{-s} \Gamma(s) \Gamma(s - k + 2) Z_F(s)$  is  $(-1)^k$ -invariant under  $s \mapsto 2k - 2 - s$ .

Let  $M_{2k-2}(\Gamma_1)$  be the space of modular forms of weight  $2k - 2$  on  $\Gamma_1$ . Recall that a Hecke eigenform in  $M_{2k-2}(\Gamma_1)$  is called normalized if its first Fourier coefficient is equal to 1.

**THEOREM 2** (Saito-Kurokawa conjecture; Andrianov [2], Maass [28, 29], Zagier [39]). *Let  $k$  be even and let  $F$  be a non-zero Hecke eigenform in  $M_k^*(\Gamma_2)$ . Then there is a unique normalized Hecke eigenform  $f$  in  $M_{2k-2}(\Gamma_1)$  such that*

$$Z_F(s) = \zeta(s - k + 1) \zeta(s - k + 2) L_f(s)$$

where  $L_f(s)$  is the Hecke  $L$ -function attached to  $f$ .

Theorem 2 in particular shows that  $Z_F(s)$  has a pole at  $s = k$  if  $F$  is a Hecke eigenform in  $S_k^*(\Gamma_2)$ . The converse is also true as shown by Evdokimov [10] and Oda [31], i.e. the function  $Z_F(s)$  is holomorphic everywhere if and only if  $F$  lies in the orthogonal complement of  $S_k^*(\Gamma_2)$ .

Using Theorem 2 one can show that  $M_k^*(\Gamma_2) = \bigcap_p \ker \mathcal{O}_p$  where  $\mathcal{O}_p$  is defined by (5). Finally let us mention that Theorem 2 implies that a Hecke eigenform  $F$  in  $S_k^*(\Gamma_2)$  does not satisfy the generalized Ramanujan-Petersson conjecture which would require that  $\lambda_n \ll_{\varepsilon, F} n^{k-3/2+\varepsilon}$  ( $\varepsilon > 0$ ).

The proof of Theorem 1 is based on the fact that the function  $V\phi$ , by definition, is symmetric w.r.t.  $\tau$  and  $\tau'$  and that  $\Gamma_2$  is generated by the matrix  $\operatorname{diag} \left( \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right)$  (which acts on  $\mathcal{H}_2$  by interchanging  $\tau$  and  $\tau'$ ) and the image of  $\Gamma_1^J$  under the map (2). For the compatibility statement of  $V$  with Hecke operators one has to check the action of the latter on Fourier coeffi-

cients. The proof of Theorem 2 is based on a trace formula. We do not give here any more details. Good expositions can be found in [9] and [39].

## 2.2. PROBLEMS

i) Since for fixed  $k$  the dimension of  $J_{k,m}$  grows linearly in  $m$ , the map  $\rho_m$  defined by (3) for  $m \gg_k 0$  cannot be surjective. Is there any simple or nice description of the image of  $\rho_m$  or  $(\text{im } \rho_m | S_k(\Gamma_2))^\perp$ ? Let us mention here that one can express the Fourier-Jacobi coefficients of Poincaré series of exponential type on  $\Gamma_2$  which generate  $S_k(\Gamma_2)$ , as certain infinite linear combinations of Poincaré series on  $\Gamma_1^J$  [22]. Taking scalar products one obtains a characterization of  $(\text{im } \rho_m | S_k(\Gamma_2))^\perp$  as the kernel of certain infinite systems of linear equations. This description, however, does not seem to be very illuminating (for example, it does not imply in any obvious way that  $\rho_1$  is surjective).

ii) A skew-holomorphic Jacobi form of weight  $k \in \mathbb{Z}$  and index  $m \in \mathbb{N}_0$  on  $\Gamma_1^J$  as introduced by Skoruppa is a complex-valued  $C^\infty$ -function  $\phi(\tau, z)$  ( $\tau \in \mathcal{H}$ ,  $z \in \mathbb{C}$ ) satisfying the following properties: 1)  $\phi$  is holomorphic in  $z$  and is annihilated by the heat operator  $8\pi i m \partial/\partial \tau - \partial^2/\partial z^2$ ; 2)  $\phi$  satisfies the same transformation formula under  $\Gamma_1^J$  as a holomorphic Jacobi form of weight  $k$  and index  $m$  (cf. § 1.2) except that the factor  $(c\tau + d)^k$  has to be replaced by  $(c\bar{\tau} + d)^{k-1} |c\tau + d|$ ; 3)  $\phi$  has a Fourier expansion of type

$$\phi(\tau, z) = \sum_{n, r \in \mathbb{Z}, r^2 \geq 4mn} c(n, r) \exp \left( -\pi \frac{r^2 - 4mn}{m} v \right) q^n \zeta^r \quad (v = \text{Im}(\tau)).$$

Note that a skew-holomorphic Jacobi form of even weight and index 1 is identically zero as is easily seen.

Despite of the importance of skew-holomorphic Jacobi forms as demonstrated in [34, 36] it is not quite clear so far how they are related to Siegel modular forms. One difficulty, for example, is that if one starts with a real-analytic Siegel modular form of genus 2, the coefficients of the partial Fourier expansion of  $F(Z)$  w.r.t.  $e^{2\pi i \tau'}$  (where as usual  $Z = \begin{pmatrix} \tau & z \\ z & \tau' \end{pmatrix}$ ) not only depend on  $\tau$  and  $z$  but also on  $\text{Im}(\tau')$ , and it is a priori not obvious how to get rid of the latter variable and to produce “true” Jacobi forms.

Let  $k$  be an odd integer and denote by  $M_{1/2, k-1/2}(\Gamma_2)$  the space of Siegel-Maass wave forms “of type  $(1/2, k-1/2)$ ” as defined in [26], i.e. the space of real-analytic functions  $F: \mathcal{H}_2 \rightarrow \mathbb{C}$  which satisfy

$$F(M \langle Z \rangle) = \det(C\bar{Z} + D)^{k-1} | \det(CZ + D) | F(Z)$$

for all  $M = \begin{pmatrix} \cdot & \cdot \\ C & D \end{pmatrix} \in \Gamma_2$  and which are annihilated by the matrix differential operator

$$\Omega_{1/2, k-1/2} := (Z - \bar{Z}) \left( (Z - \bar{Z}) \frac{\partial}{\partial Z} \right)' \frac{\partial}{\partial \bar{Z}} + \frac{1}{2} (Z - \bar{Z}) \frac{\partial}{\partial \bar{Z}} - \left( k - \frac{1}{2} \right) (Z - \bar{Z}) \frac{\partial}{\partial Z}$$

$$\text{where } \frac{\partial}{\partial Z} = \begin{pmatrix} \frac{\partial}{\partial \tau} & \frac{1}{2} \frac{\partial}{\partial z} \\ \frac{1}{2} \frac{\partial}{\partial \bar{z}} & \frac{\partial}{\partial \tau'} \end{pmatrix}$$

and  $\frac{\partial}{\partial \bar{Z}}$  is defined analogously (the notation “of type  $(1/2, k-1/2)$ ” comes from the fact that the factor of automorphy of  $F$  can be written as  $\det(C\bar{Z} + D)^{k-1/2} \det(CZ + D)^{1/2}$  with appropriate choice of the square root).

Using certain invariance properties of  $\Omega_{1/2, k-1/2}$  under the action of  $\text{Sp}_2(\mathbf{R})$  one can define Hecke operators  $T_n (n \in \mathbf{N})$  on  $M_{1/2, k-1/2}(\Gamma_2)$  in the usual way. Let

$$E_{1/2, k-1/2}^{(2)}(Z) := \sum_{(C, D)} \det(CZ + D)^{-k+1} |\det((CZ + D))|^{-1} \quad (k > 3)$$

be the Maass-Siegel-Eisenstein series in  $M_{1/2, k-1/2}(\Gamma_2)$  ([26; 27, §18]; summation over all pairs  $(C, D)$  of relatively prime symmetric  $(2, 2)$ -matrices inequivalent under left-multiplication by  $GL_2(\mathbf{Z})$ ). Then the following can be shown:

- 1) The function  $E_{1/2, k-1/2}^{(2)}$  is a Hecke eigenform whose spinor zeta function (defined in the same way as above) is equal to  $\zeta(s-k+1) \zeta(s-k+2) L_{E_{2k-2}}(s)$  where  $E_{2k-2}$  is the normalized Eisenstein series of weight  $2k-2$  on  $\Gamma_1$  (this implies that  $E_{1/2, k-1/2}^{(2)}$  for all primes  $p$  is annihilated by the Hecke operator  $\mathcal{E}_p$  defined analogously as in (5));
- 2) if  $e_{1/2, k-1/2; m}(\tau, z, \text{Im}(\tau'))$  is the  $m$ -th Fourier-Jacobi coefficient of  $E_{1/2, k-1/2}^{(2)}$  and if for  $m > 0$  one carries out a similar limit process as in [19, §2, Remark ii) after the proof of Thm. 1], i.e. essentially replaces  $\text{Im}(\tau')$  by  $(\text{Im}(z))^2 / \text{Im}(\tau) + \delta$  and lets  $\delta \rightarrow \infty$ , then one obtains a skew-holomorphic Eisenstein series of weight  $k$  and index  $m$  (in fact, finite linear combinations of such Eisenstein series if  $m$  is not squarefree).

The following questions therefore are suggestive:

- 1) if one starts with an arbitrary  $F \in M_{1/2, k-1/2}(\Gamma_2)$ , does the above limit process produce skew-holomorphic Jacobi forms of weight  $k$ ?
- 2) define  $M_{1/2, k-1/2}^*(\Gamma_2)$  as the subspace of  $M_{1/2, k-1/2}(\Gamma_2)$  consisting of the intersection of the kernels of the operators  $\mathcal{E}_p$  for all primes  $p$ . Does there exist a natural map  $V$  from skew-holomorphic Jacobi forms of weight  $k$  and index 1 to  $M_{1/2, k-1/2}^*(\Gamma_2)$  similar as in the case of holomorphic Jacobi forms?

Recently, N.-P. Skoruppa [36] has developed a theory of theta lifts from skew-holomorphic Jacobi forms to automorphic forms on  $\mathrm{Sp}_2$ . It would be interesting to investigate if his lifts would provide (at least partial) answers to the above questions.

iii) So far a generalization of the Maass space to higher genus  $n > 2$  has not been given; in fact, in the general case it does not seem to be quite clear what one has to look for, except that (the cuspidal part) of a “Maass space” eventually should be generated by Hecke eigenforms which do not satisfy a generalized Ramanujan-Petersson conjecture. Note that there is a partial negative result by Ziegler [40, 4.2. Thm.] who showed by means of specific examples that for  $n \geq 33$  the map which sends a Siegel modular form of weight 16 on  $\Gamma_n := \mathrm{Sp}_n(\mathbb{Z})$  to its first Fourier-Jacobi coefficient is not surjective.

On the other hand, there are very interesting numerical calculations for  $n = 3$  due to Miyawaki [30] which suggest that a Siegel-Hecke eigenform  $F$  of even integral weight  $k$  on  $\Gamma_3$  could be constructed from a pair  $(f, g)$  of elliptic Hecke eigenforms of weights  $(k_1, k_2)$  equal to  $(k, 2k - 4)$  or  $(k - 2, 2k - 2)$  such that the (formal) spinor zeta function of  $F$  should be equal to  $L_f(s - k_2/2) L_f(s - k_2/2 + 1) L_{f \otimes g}(s)$  where  $L_{f \otimes g}(s)$  essentially is the Rankin convolution of  $f$  and  $g$  ([*loc. cit.*, §4]; note that for  $n > 2$  the analytic continuation of the spinor zeta function of a holomorphic Hecke eigenform on  $\Gamma_n$  is not known).

### §3. SPINOR ZETA FUNCTIONS

#### 3.1. RESULTS

Although the Maass space  $S_k^*(\Gamma_2)$  as discussed in the previous section is an important subspace of  $S_k(\Gamma_2)$  in its own right, one quickly realizes that the “true” Siegel cusp forms on  $\Gamma_2$  should lie in the orthogonal complement of  $S_k^*(\Gamma_2)$  (cf. Theorem 2 in §2 and its discussion). It is therefore even more