

## 2. Realization of cubic forms

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

where the 'degree'  $d$  corresponds to the cubic form. Such a 4-tuple is admissible iff  $d(2x + W)^3 \equiv (p + 24T) \cdot (2x + W) \pmod{48}$  holds for every integer  $x$ . This is equivalent to  $p \equiv 4d \pmod{24}$  if  $\bar{W} = 0$ , and to  $p \equiv d + 24T \pmod{48}$  with  $d \equiv 0 \pmod{2}$  if  $\bar{W} \neq 0$ .

Two admissible 4-tuples  $(\bar{W}, \bar{T}, d, p)$  and  $(\bar{W}', \bar{T}', d', p')$  are equivalent iff  $\bar{W}' = \bar{W}$ ,  $\bar{T}' = \bar{T}$  and  $(d', p') = \pm(d, p)$ . Taking the degree  $d$  non-negative, we find:

**PROPOSITION 1.** *There is a 1-1 correspondence between oriented homeomorphism types of cores  $X_0$  with  $b_2(X_0) = 1$ , and 4-tuples  $(\bar{W}, \bar{T}, d, p)$ , normalized so that  $d \geq 0$ , and  $p \geq 0$  if  $d = 0$ , which satisfy  $p \equiv 4d \pmod{24}$  if  $\bar{W} = 0$ , and  $d \equiv 0 \pmod{2}$ ,  $p \equiv d + 24T \pmod{48}$  if  $\bar{W} \neq 0$ .*

In order to classify the associated homotopy types we first have to determine the subgroup  $U_F$  associated to a given cubic form  $F$ . By definition we find  $U_F = 0$  if  $d \equiv 0 \pmod{2}$ ,  $U_F = \mathbf{Z}_{/2}$  if  $d \equiv 1 \pmod{2}$ . Two normalized 4-tuples  $(\bar{W}, \bar{T}, d, p)$  and  $(\bar{W}', \bar{T}', d', p')$  are weakly equivalent iff  $d' = d$ ,  $\bar{W}' = \bar{W}$ , and  $p + 24T \equiv p' + 24T' \pmod{48}$  if  $d \equiv 0 \pmod{2}$ ,  $p \equiv p' \pmod{24}$  if  $d \equiv 1 \pmod{2}$ .

Putting everything together, we find a single oriented homotopy type for every odd degree  $d \geq 0$ , which is necessarily spin, and 3 oriented homotopy types for every even degree  $d \geq 0$ ; one of these 3 types has  $\bar{W} \neq 0$ , the other two are spin, and they are distinguished by  $p + 24T \pmod{48}$  i.e.  $p \equiv 4d \pmod{48}$ , or  $p \equiv 4d + 24 \pmod{48}$ .

## 2. REALIZATION OF CUBIC FORMS

In the previous section the (homotopy) topological classification of 1-connected, closed, oriented, 6-dimensional manifolds with torsion-free homology has been transformed into an arithmetical moduli problem: to describe the sets of (weak) equivalence classes of admissible systems of invariants. In this section we begin to investigate the latter problem; we give a simple criterion for the realizability of cubic forms by smooth manifolds, and we describe, at least in principle, the classification of homotopy types of manifolds with a given cohomology ring.

## 2.1 COHOMOLOGY RINGS OF 6-MANIFOLDS

Let  $(r, H, w, \tau, F, p)$  be a system of invariants as in section 1; recall that it is admissible iff for every  $W \in H$ ,  $T \in H^\vee$  with  $\bar{W} = w(\bmod 2)$ ,  $\bar{T} \equiv \tau(\bmod 2)$  the following congruence holds:

$$(*) \quad W^3 \equiv (p + 24T)(W) \pmod{48}.$$

LEMMA 1.  $(r, H, w, \tau, F, p)$  is admissible if and only if there exist  $W_o \in H$ ,  $T_o \in H^\vee$  with  $\bar{W}_o \equiv w(\bmod 2)$ ,  $\bar{T}_o \equiv \tau(\bmod 2)$ , such that

- i)  $W_o^3 \equiv (p + 24T_o)(W_o) \pmod{48}$
- ii)  $p(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H$ .

*Proof.* Obvious since the set of integral lifts of  $w$  is a coset  $W_o + 2H$ .

DEFINITION 3. Let  $F \in S^3 H^\vee$  be a symmetric trilinear form on a finitely generated free abelian group  $H$ . An element  $W \in H$  is characteristic for  $F$  iff

$$(**) \quad x \cdot y \cdot (x + y + W) \equiv 0 \pmod{2} \quad \forall x, y \in H.$$

LEMMA 2.  $W \in H$  is a characteristic element for  $F \in S^3 H^\vee$  if and only if the function  $l_W: H \rightarrow \mathbf{Z}$ ,  $l_W(x) := 4x^3 + 6x^2 W + 3x W^2$  is linear in  $x$  modulo 24.

*Proof.*  $l_W(x + y) = l_W(x) + l_W(y) + 12(x^2 y + x y^2 + x y W)$ , whence the assertion.

The existence of characteristic elements is a necessary and sufficient condition for a cubic form  $F \in S^3 H^\vee$  to be realizable by a manifold. In fact, we have:

PROPOSITION 2. A given cubic form  $F \in S^3 H^\vee$  on a finitely generated free abelian group  $H$  is realizable as cup-form of a 1-connected, closed, oriented, 6-dimensional manifold with torsion-free homology if and only if it possesses a characteristic element.

*Proof.* If  $(r, H, w, \tau, F, p)$  is an admissible system of invariants, and  $W_o \in H$  any integral lift of  $w$ , then we have  $p(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H$ , i.e. the function  $l_{W_o}: H \rightarrow \mathbf{Z}$  is linear modulo 24, and  $W_o$  is therefore characteristic for  $F$ . Conversely, suppose  $W_o \in H$  is a characteristic element for a cubic form  $F \in S^3 H^\vee$ ; let  $w := \bar{W}_o(\bmod 2)$ ,  $r := 0$ .

By the main lemma we have to construct linear forms  $p, T \in H^\vee$ , such that

- i)  $W_\circ^3 \equiv (p + 24T)(W_\circ) \pmod{48}$
- ii)  $p(x) \equiv 4x^3 + 6x^2 W_\circ + 3x W_\circ^2 \pmod{24} \quad \forall x \in H$ .

The function  $l_{W_\circ}: H \rightarrow \mathbf{Z}, l_{W_\circ}(x) = 4x^3 + 6x^2 W_\circ + 3x W_\circ^2$  is linear modulo 24 since  $W_\circ$  is a characteristic element for  $F$ : we therefore choose a linear form  $p_\circ \in H^\vee$  with  $p_\circ(x) \equiv l_{W_\circ}(x) \pmod{24} \quad \forall x \in H$ . Substituting  $x = W_\circ$  we find  $p_\circ(W_\circ) \equiv 13 W_\circ^3 \pmod{24}$ ; but since  $W_\circ$  is characteristic we have  $W_\circ^3 \equiv 0 \pmod{2}$ , thus  $p_\circ(W_\circ) \equiv W_\circ^3 \pmod{24}$ . Write  $p_\circ(W_\circ) = W_\circ^3 + 24k$  for some  $k \in \mathbf{Z}$ .

case 1)  $k \equiv 0 \pmod{2}$ : define  $p := p_\circ, T := 0$ .

case 2)  $k \equiv 1 \pmod{2}$ : we must find a linear form  $T_\circ \in H^\vee$  with  $T_\circ(W_\circ) \equiv 1 \pmod{2}$ ; clearly this can be done if and only if  $W_\circ$  is not divisible by 2. If  $W_\circ$  were divisible by 2,  $W_\circ = 2V_\circ$  for some  $V_\circ \in H$ , then  $2p_\circ(V_\circ) = p_\circ(W_\circ) = W_\circ^3 + 24k = 8V_\circ^3 + 24k$  would give  $p_\circ(V_\circ) = 4V_\circ^3 + 12k$ ; then, using  $p_\circ(V_\circ) \equiv 4V_\circ^3 + 6V_\circ^2 W_\circ + 3V_\circ W_\circ^2 \pmod{24}$  we would find  $k \equiv 0 \pmod{2}$ , which is not the case by assumption.

This shows that  $F \in S^3 H^\vee$  is realizable by a topological manifold with Pontrjagin class  $p_\circ$  and non-vanishing triangulation obstruction  $\tau_\circ := \bar{T}_\circ \pmod{2}$ . In order to realize  $F$  by a smooth manifold, one can take  $p := p_\circ + 24T_\circ$ , and  $\tau := 0$ .

REMARK 3. The topological counterpart of the existence of a characteristic element for a given cubic form  $F \in S^3 H^\vee$  is the existence of a mod-2 Steenrod-algebra structure, which is a necessary condition for a ring to be a cohomology ring.

The existence and the classification of characteristic elements for a given cubic form is essentially a linear algebra problem over  $\mathbf{Z}_{/2}$ . To see this, let  $F \in S^3 H^\vee$  be a fixed cubic form on a finitely generated free abelian group  $H$ . Associated with  $F$  we have a linear map  $F^t: H \rightarrow S^2 H^\vee$  sending an element  $h \in H$  to the bilinear form  $F^t(h): H \otimes H \rightarrow \mathbf{Z}, (x, y) \rightarrow x \cdot y \cdot h$ . Let  $\bar{H} := H/_{2H}$ .  $\bar{F} \in S^3 \bar{H}^\vee$  be the reductions of  $H$  and  $F$  modulo 2, and let  $-: H \rightarrow \bar{H}$  be the natural epimorphism. The symmetric trilinear form  $\bar{F}$  on the  $\mathbf{Z}_{/2}$ -module  $\bar{H}$  defines a natural symmetric bilinear form  $q_{\bar{F}} \in S^2 \bar{H}^\vee$  given by  $q_{\bar{F}}(\bar{x}, \bar{y}) := \bar{x} \cdot \bar{y} \cdot (\bar{x} + \bar{y})$ .

LEMMA 3.  $F \in S^3 H^\vee$  admits characteristic elements if and only if  $q_{\bar{F}}$  lies in the image of  $\bar{F}^t \in \text{Hom}_{\mathbf{Z}}(H, S^2 \bar{H}^\vee)$ . The set of all characteristic elements for  $F$  is a coset of the form  $W_\circ + \text{Ker}(\bar{F}^t)$ .

*Proof.*  $W_o$  is characteristic for  $F$  if and only if  $q_{\bar{F}} = \bar{F}^t(W_o)$ .

In terms of a  $\mathbf{Z}$ -basis  $\{e_1, \dots, e_b\}$  for  $H$  the condition  $q_{\bar{F}} \in \text{Im}(\bar{F}^t)$  translates into a simple rank condition over  $\mathbf{Z}_{/2}$ : the  $\mathbf{Z}_{/2}$ -rank of the  $b \times \binom{b+1}{2}$ -matrix  $A$  representing  $\bar{F}^t$  must be equal to the  $\mathbf{Z}_{/2}$ -rank of the matrix  $A$  extended by the column  $(\bar{e}_i \cdot \bar{e}_j \cdot (\bar{e}_i + \bar{e}_j))_{1 \leq i \leq j \leq b}$

EXAMPLE 3. Let  $H = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$  be free of rank 2,  $F \in S^3 H^\vee$  given by  $e_1^3 = a$ ,  $e_1^2 e_2 = b$ ,  $e_1 e_2^2 = c$ ,  $e_2^3 = d$  with  $a, b, c, d \in \mathbf{Z}$ . The rank condition becomes

$$rk_2 \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \\ \bar{b} & \bar{c} \end{bmatrix} = rk_2 \begin{bmatrix} \bar{a} & \bar{b} & \bar{0} \\ \bar{c} & \bar{d} & \bar{0} \\ \bar{b} & \bar{c} & \overline{b+c} \end{bmatrix}$$

## 2.2 HOMOTOPY TYPES WITH A GIVEN COHOMOLOGY RING

Our next task is to describe the set of oriented homotopy types of 1-connected, closed, oriented, 6-dimensional manifolds with a fixed torsion-free cohomology ring.

From Žubr's classification theorem we know that in algebraic terms this means the following: fix a non-negative integer  $r_o$ , a finitely generated free abelian group  $H_o$ , and a symmetric trilinear form  $F_o \in S^3 H_o^\vee$  which admits characteristic elements.

Let  $\mathcal{M}(r_o, H_o, F_o)$  be the set of 1-connected, closed, oriented, 6-dimensional manifolds  $X$  with  $b_3(X) = 2r_o$ , such that there exists an isomorphism  $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$  with  $\alpha^* F_X = F_o$ . Denote by  $\text{Aut}(F_o)$  the subgroup of  $\mathbf{Z}$ -automorphisms of  $H_o$  which leave  $F_o \in S^3 H_o^\vee$  invariant;  $\text{Aut}(F_o)$  acts on pairs  $(w, [l]) \in \bar{H}_o \times H_o^\vee / {}_{48}H_o^\vee / {}_{U_{F_o}}$  in a natural way:

$$\gamma \cdot (w, [l]) := (\gamma(w), (\gamma^{-1})^*[l]) .$$

Let  ${}_{\text{Aut}(F_o)} \backslash \bar{H}_o \times H_o^\vee / {}_{48}H_o^\vee / {}_{U_{F_o}}$  be the set of  $\text{Aut}(F_o)$ -orbits.

A manifold  $X$  in  $\mathcal{M}(r_o, H_o, F_o)$  and an isomorphism  $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$  with  $\alpha^* F_X = F_o$  yields a well-defined  $\text{Aut}(F_o)$ -orbit:

$$(\alpha^{-1}(w_2(X)), \alpha^*[p_1(X) + 24T]) \text{ (modulo } \text{Aut}(F_o) \text{)},$$

where  $T \in H^4(X, \mathbf{Z})$  is an arbitrary integral lifting of  $\tau(X) \in H^4(X, \mathbf{Z}_{/2})$ .

The set of oriented homotopy types  $\mathcal{M}(r_o, H_o, F_o) / \simeq$  of manifolds in  $\mathcal{M}(r_o, H_o, F_o)$  can now be described in the following way:

PROPOSITION 3. *The assignment  $X \mapsto (\alpha^{-1}(w_2(X)), \alpha^*[p_1(X) + 24T])$  (modulo  $\text{Aut}(F_o)$ ) defines an injection.*

$$I: \mathcal{M}(r_o, H_o, F_o)/\simeq \rightarrow_{\text{Aut}(F_o)} \bar{H}_o \times H_o^\vee /_{48H_o^\vee} /_{U_{F_o}}.$$

*Proof.* Suppose  $X$  and  $X'$  are manifolds in  $\mathcal{M}(r_o, H_o, F_o)$ ,  $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$  and  $\alpha': H_o \rightarrow H^2(X', \mathbf{Z})$  isomorphisms with  $\alpha^*F_X = F_o$  and  $(\alpha')^*F_{X'} = F_o$ .  $X$  and  $X'$  have the same image under  $I$  iff there exists an automorphism  $\gamma \in \text{Aut}(F_o)$  with  $\gamma\alpha^{-1}(w_2(X)) = (\alpha')^{-1}w_2(X')$  and  $(\gamma^{-1})^*\alpha^*[p_1(X) + 24T] = (\alpha')^*[p_1(X') + 24T']$ . Consider  $\beta := \alpha \circ \gamma \circ \alpha^{-1}: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ ;  $\beta$  is obviously an isomorphism with  $\beta^*F_{X'} = F_X$ ,  $\beta w_2(X) = w_2(X')$ , and  $\beta^*[p_1(X') + 24T'] = [p_1(X) + 24T]$ ; but this means that the systems of invariants associated with  $X$  and  $X'$  are weakly equivalent, and therefore  $X$  and  $X'$  oriented homotopy equivalent.

A complete description of the set  $\mathcal{M}(r_o, H_o, F_o)/\simeq$  i.e. of the image of  $I$  is only possible if the automorphism group  $\text{Aut}(F_o)$  is known; this can be a serious problem, but we will see that the ‘general’ automorphism group is finite (and usually small), so that the next proposition gives a reasonable estimate for the number of elements in  $\mathcal{M}(r_o, H_o, F_o)/\simeq$ .

PROPOSITION 4. *Fix  $r_o \in \mathbf{N}$ , a finitely generated free abelian group  $H_o$ , and a symmetric trilinear form  $F_o \in S^3 H_o^\vee$  which admits characteristic elements. Set  $b := rk_{\mathbf{Z}} H_o$ ,  $s := rk_{\mathbf{Z}/2}(\bar{F}_o^t)$ , and let  $t := rk_{\mathbf{Z}/2}(\cdot_{\bar{F}_o})$  be the  $\mathbf{Z}/2$ -rank of the  $\mathbf{Z}/2$ -linear square map  $\cdot_{\bar{F}_o}: \bar{H}_o \rightarrow \bar{H}_o^\vee$  sending  $\bar{u} \in \bar{H}_o$  to  $\bar{u}^2 \in \bar{H}_o^\vee$ . Then  $\mathcal{M}(r_o, H_o, F_o)/\simeq$  contains at most  $2^{2b-s-t}$  elements.*

*Proof.* Fix any admissible system of invariants  $(r_o, H_o, w_o, \tau_o, F_o, p_o)$  for a manifold in  $\mathcal{M}(r_o, H_o, F_o)$ . Given  $(r_o, H_o, F_o)$ , we know from the last lemma that the possible elements  $w_o$  form a coset of  $\text{Ker}(\bar{F}_o^t)$  in  $\bar{H}_o$ , so that there exist precisely  $2^{b-s}$  such elements. It remains to count the classes  $[l] \in H_o^\vee /_{48H_o^\vee} /_{U_{F_o}}$ , such that the  $\text{Aut}(F_o)$ -orbit of  $(w_o, [p_o + 24T_o + l])$  lies in the image of  $I$ .

To understand the latter condition we fix integral liftings  $W_o, \in H_o, T_o \in H_o^\vee$  of  $w_o$  and  $\tau_o$  satisfying the admissibility conditions

- i)  $W_o^3 \equiv (p_o + 24T_o)(W_o) \pmod{48}$
- ii)  $p_o(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H_o.$

Clearly the  $\text{Aut}(F_o)$ -orbit of  $(w_o, [p_o + 24T_o + l])$  lies in the image of  $I$  if and only if

$$\text{i}') \quad W_o^3 \equiv (p_o + 24T_o + l)(W_o) \pmod{48},$$

$$\text{ii}') \quad (p_o + l)(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H_o,$$

which is equivalent to  $l(W_o) \equiv 0 \pmod{48}$ , and  $l \equiv 0 \pmod{24H_o^\vee}$  because of i) and ii).

Now, by definition of the subgroup  $U_{F_o} \subset H_o^\vee / {}_{48}H_o^\vee$  we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & \text{Ker}(\cdot \bar{F}_o) & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 \rightarrow \text{Ker}(24 \cdot \bar{F}_o) & \hookrightarrow & H_o / {}_2H_o & \xrightarrow{24 \cdot \bar{F}_o} & U_{F_o} & \rightarrow & 0 \\
 & & \downarrow \cdot \bar{F}_o & & \downarrow & & \\
 & 0 \rightarrow & H_o^\vee / {}_2H_o^\vee & \xrightarrow{24} & H_o^\vee / {}_{48}H_o^\vee & \rightarrow & H_o^\vee / {}_{24}H_o^\vee \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 \rightarrow \text{Coker}(\cdot \bar{F}_o) & \rightarrow & H_o^\vee / {}_{48}H_o^\vee / U_{F_o} & \rightarrow & H_o^\vee / {}_{24}H_o^\vee & \rightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 
 \end{array}$$

The number of elements  $[l] \in H_o^\vee / {}_{48}H_o^\vee / U_{F_o}$  to be counted coincides therefore with the cardinality of the kernel of the map  $ev(w_o): \text{Coker}(\cdot \bar{F}_o) \rightarrow \mathbf{Z}_{/2}$  induced by evaluation in  $w_o$ . This number is at most  $2^{b-t}(2^{b-t-1}$  if  $w_o \neq 0$  and  $t \neq b$ ).

**COROLLARY 2.** *If the  $\mathbf{Z}_{/2}$ -rank  $s = rk_{\mathbf{Z}_{/2}}(\cdot \bar{F}_o)$  is maximal, then  $\mathcal{M}(r_o, H_o, F_o)/\simeq$  contains at most one class.*

*Proof.* Suppose  $\cdot \bar{F}_o: \bar{H}_o \rightarrow \bar{H}_o^\vee$  is surjective; then  $\bar{F}_o^t: \bar{H}_o \rightarrow S^2 \bar{H}_o^\vee$  must have a trivial kernel, since  $\bar{h}\bar{x}^2 = 0$  for all  $\bar{x} \in \bar{H}_o$  implies  $\bar{h} = 0$  if every linear form is a square. But this means  $s = t = b$ , so that  $\mathcal{M}(r_o, H_o, F_o)/\simeq$  has at most one element.

**EXAMPLE 4.** Let  $H_o = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$ ,  $e_1^3 = a$ ,  $e_1^2 e_2 = b$ ,  $e_1 e_2^2 = c$ ,  $e_2^3 = d$ . If  $\bar{b} \equiv \bar{c} \pmod{2}$ , and  $\bar{a}\bar{d} - \bar{b}\bar{c} \equiv 1 \pmod{2}$ , then  $\mathcal{M}(r_o, H_o, F_o)/\simeq$  contains precisely one class for every  $r_o \geq 0$ .