

2.1 COHOMOLOGY RINGS OF 6-MANIFOLDS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2.1 COHOMOLOGY RINGS OF 6-MANIFOLDS

Let (r, H, w, τ, F, p) be a system of invariants as in section 1; recall that it is admissible iff for every $W \in H$, $T \in H^\vee$ with $\bar{W} = w \pmod{2}$, $\bar{T} \equiv \tau \pmod{2}$ the following congruence holds:

$$(*) \quad W^3 \equiv (p + 24T)(W) \pmod{48}.$$

LEMMA 1. (r, H, w, τ, F, p) is admissible if and only if there exist $W_0 \in H$, $T_0 \in H^\vee$ with $\bar{W}_0 \equiv w \pmod{2}$, $\bar{T}_0 \equiv \tau \pmod{2}$, such that

- i) $W_0^3 \equiv (p + 24T_0)(W_0) \pmod{48}$
- ii) $p(x) \equiv 4x^3 + 6x^2W_0 + 3xW_0^2 \pmod{24} \quad \forall x \in H.$

Proof. Obvious since the set of integral lifts of w is a coset $W_0 + 2H$.

DEFINITION 3. Let $F \in S^3 H^\vee$ be a symmetric trilinear form on a finitely generated free abelian group H . An element $W \in H$ is characteristic for F iff

$$(**) \quad x \cdot y \cdot (x + y + W) \equiv 0 \pmod{2} \quad \forall x, y \in H.$$

LEMMA 2. $W \in H$ is a characteristic element for $F \in S^3 H^\vee$ if and only if the function $l_W: H \rightarrow \mathbf{Z}$, $l_W(x) := 4x^3 + 6x^2W + 3xW^2$ is linear in x modulo 24.

Proof. $l_W(x + y) = l_W(x) + l_W(y) + 12(x^2y + xy^2 + xyW)$, whence the assertion.

The existence of characteristic elements is a necessary and sufficient condition for a cubic form $F \in S^3 H^\vee$ to be realizable by a manifold. In fact, we have:

PROPOSITION 2. A given cubic form $F \in S^3 H^\vee$ on a finitely generated free abelian group H is realizable as cup-form of a 1-connected, closed, oriented, 6-dimensional manifold with torsion-free homology if and only if it possesses a characteristic element.

Proof. If (r, H, w, τ, F, p) is an admissible system of invariants, and $W_0 \in H$ any integral lift of w , then we have $p(x) \equiv 4x^3 + 6x^2W_0 + 3xW_0^2 \pmod{24} \quad \forall x \in H$, i.e. the function $l_{W_0}: H \rightarrow \mathbf{Z}$ is linear modulo 24, and W_0 is therefore characteristic for F . Conversely, suppose $W_0 \in H$ is a characteristic element for a cubic form $F \in S^3 H^\vee$; let $w := \bar{W}_0 \pmod{2}$, $r := 0$.

By the main lemma we have to construct linear forms $p, T \in H^\vee$, such that

- i) $W_\circ^3 \equiv (p + 24T)(W_\circ) \pmod{48}$
- ii) $p(x) \equiv 4x^3 + 6x^2 W_\circ + 3x W_\circ^2 \pmod{24} \quad \forall x \in H.$

The function $l_{W_\circ}: H \rightarrow \mathbf{Z}$, $l_{W_\circ}(x) = 4x^3 + 6x^2 W_\circ + 3x W_\circ^2$ is linear modulo 24 since W_\circ is a characteristic element for F : we therefore choose a linear form $p_\circ \in H^\vee$ with $p_\circ(x) \equiv l_{W_\circ}(x) \pmod{24} \quad \forall x \in H$. Substituting $x = W_\circ$ we find $p_\circ(W_\circ) \equiv 13W_\circ^3 \pmod{24}$; but since W_\circ is characteristic we have $W_\circ^3 \equiv 0 \pmod{2}$, thus $p_\circ(W_\circ) \equiv W_\circ^3 \pmod{24}$. Write $p_\circ(W_\circ) = W_\circ^3 + 24k$ for some $k \in \mathbf{Z}$.

case 1) $k \equiv 0 \pmod{2}$: define $p := p_\circ$, $T := 0$.

case 2) $k \equiv 1 \pmod{2}$: we must find a linear form $T_\circ \in H^\vee$ with $T_\circ(W_\circ) \equiv 1 \pmod{2}$; clearly this can be done if and only if W_\circ is not divisible by 2. If W_\circ were divisible by 2, $W_\circ = 2V_\circ$ for some $V_\circ \in H$, then $2p_\circ(V_\circ) = p_\circ(W_\circ) = W_\circ^3 + 24k = 8V_\circ^3 + 24k$ would give $p_\circ(V_\circ) = 4V_\circ^3 + 12k$; then, using $p_\circ(V_\circ) \equiv 4V_\circ^3 + 6V_\circ^2 W_\circ + 3V_\circ W_\circ^2 \equiv 4V_\circ^3 \pmod{24}$ we would find $k \equiv 0 \pmod{2}$, which is not the case by assumption.

This shows that $F \in S^3 H^\vee$ is realizable by a topological manifold with Pontrjagin class p_\circ and non-vanishing triangulation obstruction $\tau_\circ := \bar{T}_\circ \pmod{2}$. In order to realize F by a smooth manifold, one can take $p := p_\circ + 24T_\circ$, and $\tau := 0$.

REMARK 3. The topological counterpart of the existence of a characteristic element for a given cubic form $F \in S^3 H^\vee$ is the existence of a mod-2 Steenrod-algebra structure, which is a necessary condition for a ring to be a cohomology ring.

The existence and the classification of characteristic elements for a given cubic form is essentially a linear algebra problem over $\mathbf{Z}_{/2}$. To see this, let $F \in S^3 H^\vee$ be a fixed cubic form on a finitely generated free abelian group H . Associated with F we have a linear map $F^t: H \rightarrow S^2 H^\vee$ sending an element $h \in H$ to the bilinear form $F^t(h): H \otimes H \rightarrow \mathbf{Z}$, $(x, y) \mapsto x \cdot y \cdot h$. Let $\bar{H} := H_{/2H}$. $\bar{F} \in S^3 \bar{H}^\vee$ be the reductions of H and F modulo 2, and let $-: H \rightarrow \bar{H}$ be the natural epimorphism. The symmetric trilinear form \bar{F} on the $\mathbf{Z}_{/2}$ -module \bar{H} defines a natural symmetric bilinear form $q_{\bar{F}} \in S^2 \bar{H}^\vee$ given by $q_{\bar{F}}(\bar{x}, \bar{y}) := \bar{x} \cdot \bar{y} \cdot (\bar{x} + \bar{y})$.

LEMMA 3. $F \in S^3 H^\vee$ admits characteristic elements if and only if $q_{\bar{F}}$ lies in the image of $\bar{F}^t \in \text{Hom}_{\mathbf{Z}}(H, S^2 \bar{H}^\vee)$. The set of all characteristic elements for F is a coset of the form $W_\circ + \text{Ker}(\bar{F}^t)$.

Proof. W_\circ is characteristic for F if and only if $q_{\bar{F}} = \bar{F}^t(W_\circ)$.

In terms of a \mathbf{Z} -basis $\{e_1, \dots, e_b\}$ for H the condition $q_{\bar{F}} \in \text{Im}(\bar{F}^t)$ translates into a simple rank condition over $\mathbf{Z}_{/2}$: the $\mathbf{Z}_{/2}$ -rank of the $b \times \binom{b+1}{2}$ -matrix A representing \bar{F}^t must be equal to the $\mathbf{Z}_{/2}$ -rank of the matrix A extended by the column $(\bar{e}_i \cdot \bar{e}_j \cdot (\bar{e}_i + \bar{e}_j))_{1 \leq i \leq j \leq b}$

EXAMPLE 3. Let $H = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$ be free of rank 2, $F \in S^3 H^\vee$ given by $e_1^3 = a$, $e_1^2 e_2 = b$, $e_1 e_2^2 = c$, $e_2^3 = d$ with $a, b, c, d \in \mathbf{Z}$. The rank condition becomes

$$rk_2 \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \\ \bar{b} & \bar{c} \end{bmatrix} = rk_2 \begin{bmatrix} \bar{a} & \bar{b} & \bar{0} \\ \bar{c} & \bar{d} & \bar{0} \\ \bar{b} & \bar{c} & \bar{b+c} \end{bmatrix}$$

2.2 HOMOTOPY TYPES WITH A GIVEN COHOMOLOGY RING

Our next task is to describe the set of oriented homotopy types of 1-connected, closed, oriented, 6-dimensional manifolds with a fixed torsion-free cohomology ring.

From Žubr's classification theorem we know that in algebraic terms this means the following: fix a non-negative integer r_\circ , a finitely generated free abelian group H_\circ , and a symmetric trilinear form $F_\circ \in S^3 H_\circ^\vee$ which admits characteristic elements.

Let $\mathcal{M}(r_\circ, H_\circ, F_\circ)$ be the set of 1-connected, closed, oriented, 6-dimensional manifolds X with $b_3(X) = 2r_\circ$, such that there exists an isomorphism $\alpha: H_\circ \rightarrow H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_\circ$. Denote by $\text{Aut}(F_\circ)$ the subgroup of \mathbf{Z} -automorphisms of H_\circ which leave $F_\circ \in S^3 H_\circ^\vee$ invariant; $\text{Aut}(F_\circ)$ acts on pairs $(w, [l]) \in \bar{H}_\circ \times H_\circ^\vee /_{48 H_\circ^\vee} /_{U_{F_\circ}}$ in a natural way:

$$\gamma \cdot (w, [l]) := (\gamma(w), (\gamma^{-1})^* [l]).$$

Let ${}_{\text{Aut}(F_\circ)} \backslash \bar{H}_\circ \times H_\circ^\vee /_{48 H_\circ^\vee} /_{U_{F_\circ}}$ be the set of $\text{Aut}(F_\circ)$ -orbits.

A manifold X in $\mathcal{M}(r_\circ, H_\circ, F_\circ)$ and an isomorphism $\alpha: H_\circ \rightarrow H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_\circ$ yields a well-defined $\text{Aut}(F_\circ)$ -orbit:

$$(\alpha^{-1}(w_2(X)), \alpha^* [p_1(X) + 24T]) \pmod{\text{Aut}(F_\circ)},$$

where $T \in H^4(X, \mathbf{Z})$ is an arbitrary integral lifting of $\tau(X) \in H^4(X, \mathbf{Z}_{/2})$.

The set of oriented homotopy types $\mathcal{M}(r_\circ, H_\circ, F_\circ)/_{\sim}$ of manifolds in $\mathcal{M}(r_\circ, H_\circ, F_\circ)$ can now be described in the following way: