

## **5.2 3-folds with $b_2 = 2$**

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.05.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

There will certainly be some gaps for algebraic 3-folds. In order to show this, we prove the following finiteness theorem for families of Kähler structures:

**THEOREM 7.** *Fix a positive constant  $c$ . There exist only finitely many families of 1-connected, smooth projective 3-folds  $X$  with  $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$ ,  $w_2(X) \neq 0$ , and with  $b_3(X) \leq c$ .*

*Proof.* Let  $X$  be a smooth projective 3-fold with  $H_1(X, \mathbf{Z}) = \{0\}$ ,  $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$ , and with  $w_2(X) \neq 0$ . Clearly  $\text{Pic}(X) \cong H^2(X, \mathbf{Z})$ , and we can choose a basis  $e \in H^2(X, \mathbf{Z})$  corresponding to the ample generator of  $\text{Pic}(X)$ .

Let  $c_1(X) = c_1 e$ ,  $c_2(X) = c_2 \varepsilon$  where  $e^2 = d\varepsilon$ ,  $\varepsilon(e) = 1$ . If  $c_1$  is positive, then  $X$  is Fano, and there are only finitely many possibilities [Mu]. The case  $c_1 = 0$  is excluded, so that we are left with  $c_1 < 0$ , i.e. the canonical bundle of  $X$  is ample.

The Riemann-Roch formula  $\chi(X, \mathcal{O}_X) = 1 - h^3(X, \mathcal{O}_X) = \frac{1}{24} c_1 c_2$  shows that the set of possible Chern numbers  $c_1 c_2$  is bounded from below:  $24(1 - c) \leq c_1 c_2$ . Using Yau's inequality  $8c_1(X)c_2(X) \leq 3c_1(X)^3$  we find that  $d | c_1 |^3 \leq 64(c - 1)$ , i.e. the degree  $d$  and the order of divisibility  $| c_1 |$  of  $c_1(X)$  is bounded. Now Kollar's finiteness theorem [Ko2] yields the assertion.

**EXAMPLE 15.** Let  $X$  be a 1-connected, smooth projective 3-fold with  $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$  and  $w_2(X) \neq 0$ . If  $b_3(X) \leq 2$ , then  $h^3(X, \mathcal{O}_X) \leq 1$  and  $X$  must be Fano of index 1 or 3. For  $b_3(X) = 4$  we have that  $X$  is either Fano, or  $h^3(X, \mathcal{O}_X) = 2$  and  $X$  is of general type with  $d | c_1 |^3 \leq 64$ .

Note that the assumption  $w_2 \neq 0$  was only used to exclude Calabi-Yau 3-folds.

## 5.2 3-FOLDS WITH $b_2 = 2$

Let  $X$  be a 1-connected, closed, oriented, 6-dimensional differentiable manifold with  $H_2(X, \mathbf{Z}) \cong \mathbf{Z}^2$ .

We choose a basis  $e_1, e_2$  for  $H^2(X, \mathbf{Z})$  and set  $a_0 = e_1^3, a_1 = e_1^2 e_2, a_2 = e_1 e_2^2, a_3 = e_2^3$ ; the cubic polynomial  $f$  associated to the cup-form of  $X$  is then given by  $f = a_0 X^3 + 3a_1 X^2 Y + 3a_2 X Y^2 + a_3 Y^3$ . The discriminant of  $f$  is by definition  $\Delta(f) = a_0^2 a_3^2 - 3a_1^2 a_2^2 - 6a_0 a_1 a_2 a_3 + 4a_0 a_2^3 + 4a_1^3 a_3$ ; up to a factor it is simply the discriminant of the Hessian  $H_f = 6^2[(a_0 a_2 - a_1^2) X^2 + (a_0 a_3 - a_1 a_2) X Y + (a_1 a_3 - a_2^2) Y^2]$  of  $f$ :  $\Delta(f) = (a_0 a_3 - a_1 a_2)^2 - 4(a_0 a_2 - a_1^2)(a_1 a_3 - a_2^2)$ .

The last identity shows that  $\Delta(f)$  is always a square modulo 4, i.e.  $\Delta(f) \equiv 0, 1 \pmod{4}$ .

**PROPOSITION 17.** *Every integer  $\Delta \equiv 0, 1 \pmod{4}$  is realizable as discriminant of a compact complex 3-fold.*

*Proof.* Consider the projectivization  $X = \mathbf{P}_{\mathbf{P}^2}(E)$  of a holomorphic rank-2 vector bundle  $E$  over the plane. In terms of the standard basis of  $H^2(X, \mathbf{Z})$  ( $e_1 = \pi^* h, e_2 = c_1(\mathcal{O}_{\mathbf{P}(E)}(1))$ ) the cubic polynomial associated to  $X$  is given by  $f = (c_1^2 - c_2)X^3 + 3(-c_1)X^2Y + 3XY^2$ , where  $c_i = c_i(E)$  are the Chern classes of  $E$  considered as integers. Inserting this into the discriminant formula yields  $\Delta(f) = c_1^2 - 4c_2$ . Since every pair  $c_1, c_2$  occurs as pair of Chern classes of a holomorphic rank-2 bundle on  $\mathbf{P}^2$ , every integer  $\Delta \equiv 0, 1 \pmod{4}$  can be realized as discriminant of a holomorphic projective bundle  $\mathbf{P}_{\mathbf{P}^2}(E)$ .

Recall from section 3.2 that there are 4 different types of  $SL(2)$ -orbits of complex binary cubics: non-singular forms  $f$  (with  $\Delta(f) \neq 0$ ), and three orbits of singular cubics, represented by the normal forms  $X^2Y, X^3$ , and 0.

**PROPOSITION 18.** *All four types of complex binary cubics are realizable by complex 3-folds.*

*Proof.* We have seen this already for non-singular cubics. Clearly the product  $\mathbf{P}^1 \times \mathbf{P}^2$  realizes the normal form  $X^2Y$ . The cubics of normal forms  $X^3$  or 0 are degenerate, i.e. their Hessians vanish identically. Therefore they can only be realized by non-Kählerian 3-folds. To realize  $X^3$  one can blow up a point in an elliptic fiber bundle over a surface  $Y$  with  $b_2(Y) = 3$ ; the trivial form occurs for elliptic fiber bundles over a surface with  $b_2 = 4$ .

More detailed investigations of the possible homotopy types of real or complex manifolds with  $b_2 = 2$  will appear elsewhere [Sch].

Here we only want to illustrate an interesting phenomenon which relates the ample cone of a projective 3-fold with  $b_2 = 2$  to the Hessian of its cup-form.

**PROPOSITION 19.** *Let  $X$  be a smooth projective 3-fold with  $b_2(X) = 2$ . The ample cone  $\mathcal{C}_X$  is contained in the Hesse cone  $\mathcal{H}_F := \{h \in H^2(X, \mathbf{R}) \mid \det(F^t(h)) < 0\}$ .*

*Proof.* This is only a special case of our general result in section 4.3.

REMARK 14. The Hessian of a binary form  $F \in S^3 H^\vee$  is identically zero iff  $F$  is degenerate; it is negative semi-definite if  $F$  is non-degenerate and  $\Delta(F) \leq 0$ ; it is indefinite iff  $\Delta(F) > 0$  [Ca]. Only in the indefinite case  $\Delta(F) > 0$  can the closure  $\bar{\mathcal{H}}_F := \{h \in H_{\mathbf{R}} \mid \det F'(h) \leq 0\}$  of the Hesse cone be a proper subset of  $H_{\mathbf{R}}$ .

EXAMPLE 16. Let  $P = \mathbf{P}_{\mathbf{P}^2}(E)$  be the projectivization of a rank-2 vector bundle  $E$  with Chern classes  $c_i = c_i(E)$ . The cup-form of  $P$  yields the cubic polynomial  $f = (c_1^2 - c_2)X^2 + 3(-c_1)X^2Y + 3XY^2$  whose Hessian is  $H_f = (-c_2)X^2 + c_1XY - Y^2$ . Rewriting  $H_f$  as  $H_f = -\frac{1}{4}[(2Y - c_1X)^2 + X^2(4c_2 - c_1^2)] = \frac{-1}{4}[(2Y - c_1X)^2 - \Delta(f)X^2]$  we find 3 possibilities for the Hesse cone:

- i)  $\Delta(f) < 0$ :  $\mathcal{H}_f = H^2(P, \mathbf{R}) \setminus \{0\}$
- ii)  $\Delta(f) = 0$ :  $\mathcal{H}_f = H^2(P, \mathbf{R}) \setminus L_{c_1}$  for a real line  $L_{c_1}$  depending on  $c_1$  ( $L_{c_1} = \mathbf{R}(2, c_1)$  in the coordinates  $X, Y$ )
- iii)  $\Delta(f) > 0$ :  $\mathcal{H}_f$  is an open cone whose angle is determined by  $\Delta(f)$   $((Z + \sqrt{\Delta(f)}X)(Z - \sqrt{\Delta(f)}X) > 0$  in coordinates  $X, Z := 2Y - c_1X$ ).

### 5.3 3-FOLDS WITH $b_2 \geq 3$

Let  $X$  be a 1-connected, compact complex 3-fold with  $H_2(X, \mathbf{Z}) \cong \mathbf{Z}^{\oplus 3}$ . The cup-form of  $X$  gives rise to a curve  $C_X$  of degree 3 in the projective plane  $\mathbf{P}(H^2(X, \mathbf{C}))$ :

$$C_X := \{ \langle h \rangle \in \mathbf{P}(H^2(X, \mathbf{C})) \mid h^3 = 0 \} .$$

A first natural question is which types of plane cubic curves occur in this way?

Recall that there are 10 types of plane cubics, namely: 1) non-singular cubics, 2) irreducible cubics with a node, 3) irreducible cubics with a cusp, 4) reducible cubics consisting of a smooth conic and a transversal line, 5) smooth conics with a tangent line, 6) three lines forming a triangle, 7) three distinct lines through a common point, 8) a double line with a third skew line, 9) a triple line, 10) the trivial ‘cubic’ with equation 0.

LEMMA 4. *If the 3-fold  $X$  has a non-trivial Hodge number  $h^{2,0}(X) \neq 0$ , then  $C_X$  is of type 4), 6) 9) or 10).*