

6. Topologically Tame Manifolds

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

semidirect product of K and a connected simply-connected nilpotent Lie group and Γ is a discrete cocompact subgroup of G [12, Theorem 6.4]. We may as well assume that $X = \Gamma \backslash G/K$. By passing to a finite cover, we may assume that K is trivial. That is, X is a nilmanifold. From [27, Corollary 7.28], $H^p(X; \mathbf{C}) \cong H^p(g, \mathbf{C})$, the Lie algebra cohomology of g . From [7], $H^p(g, \mathbf{C}) \neq 0$ for all $p \in [0, \dim(X)]$. Thus for all $p \in [0, \dim(X)]$, $H^p(X; \mathbf{C}) \neq 0$.

Now let ω be a nonzero harmonic p -form on X . Let $\pi^*\omega$ be its pullback to \tilde{X} . The idea is to construct low-energy square-integrable p -forms on X by multiplying $\pi^*\omega$ by appropriate functions on X . We define the functions as in [2, §2]. Take a smooth triangulation of X and choose a fundamental domain F for the lifted triangulation of \tilde{X} . If E is a finite subset of $\pi_1(X)$, let χ_H be the characteristic function of $H = \bigcup_{g \in E} g \cdot F$. Given numbers $0 < \epsilon_1 < \epsilon_2 < 1$, choose a nonincreasing function $\psi \in C_0^\infty([0, \infty))$ which is identically one on $[0, \epsilon_1]$ and identically zero on $[\epsilon_2, \infty)$. Define a compactly-supported function f_E on \tilde{X} by $f_E(m) = \psi(d(m, H))$. Then there is a constant $C_1 > 0$, independent of E , such that

$$(5.12) \quad \int_{\tilde{X}} |df_E|^2 \leq C_1 \operatorname{area}(\partial H).$$

Define $\rho_E \in \Lambda^p(\tilde{X})$ by $\rho_E = f_E \cdot \pi^*\omega$. We have $d\rho_E = df_E \wedge \pi^*\omega$ and $d^*\rho_E = -i(df_E)\pi^*\omega$. As f_E is identically one on H , it follows that there is a constant $C > 0$, independent of E , such that

$$(5.13) \quad \frac{\int_{\tilde{X}} [|d\rho_E|^2 + |d^*\rho_E|^2]}{\int_{\tilde{X}} |\rho_E|^2} \leq C \frac{\operatorname{area}(\partial H)}{\operatorname{vol}(H)}.$$

As $\pi_1(X)$ is amenable, by an appropriate choice of E this ratio can be made arbitrarily small. Thus $0 \in \sigma(\Delta_p)$. \square

QUESTION. Does the conclusion of Proposition 20 hold if we only assume that $\pi_1(X)$ is amenable?

6. TOPOLOGICALLY TAME MANIFOLDS

Another class of manifolds for which one can hope to get some nontrivial results about the zero-in-the-spectrum question is given by *topologically tame* manifolds, meaning manifolds M which are diffeomorphic to the interior of a compact manifold N with boundary. If M has finite volume then $\operatorname{Ker}(\Delta_0) \neq 0$,

so we restrict our attention to the infinite volume case. A limited result is given by Corollary 2.

An interesting class of topologically tame manifolds consists of those which are radially symmetric. This means that M is diffeomorphic to \mathbf{R}^n , with a metric which is given on $\mathbf{R}^n - \{0\} \cong (0, \infty) \times S^{n-1}$ by

$$(6.1) \quad g = dr^2 + \phi^2(r)d\Omega^2.$$

Here $d\Omega^2$ is the standard metric on S^{n-1} , $r \in (0, \infty)$, $\phi \in C^\infty([0, \infty))$, $\phi(0) = 0$, $\phi'(0) = 1$ and $\phi(r) > 0$ for $r > 0$.

PROPOSITION 21. *Suppose that there is a constant $c \geq 0$ such that $\text{Ricci}_M \geq -c^2$. Then $0 \in \sigma(\Delta_p)$ for some p .*

Proof. We may assume that $\text{vol}(M) = \infty$. Suppose first that

$$\liminf_{r \rightarrow \infty} \phi(r) < \infty.$$

Then there is a constant $C > 0$ and a sequence $\{r_j\}_{j=1}^\infty$ such that $\lim_{j \rightarrow \infty} r_j = \infty$ and $\phi(r_j) \leq C$. Let D_j be the domain in M defined by $r \leq r_j$. Then $\text{area}(D_j) \leq C^{n-1} \text{vol}(S^{n-1})$ and $\lim_{j \rightarrow \infty} \text{vol}(D_j) = \infty$. Thus M is not open at infinity. By Proposition 6, $0 \in \sigma(\Delta_0)$.

Now suppose that $\liminf_{r \rightarrow \infty} \phi(r) = \infty$. We want to show that M is hyperEuclidean and apply Proposition 7. Consider a map $F : M \rightarrow \mathbf{R}^n$ given in polar coordinates by

$$(6.2) \quad F(r, \theta) = (s(r), \theta),$$

for some $s : [0, \infty) \rightarrow [0, \infty)$. The condition for F to be distance-nonincreasing is

$$(6.3) \quad |s'(r)| \leq 1, \quad s(r) \leq \phi(r).$$

If $\lim_{r \rightarrow \infty} s(r) = \infty$ then F is a proper map of degree one. It remains to construct s satisfying (6.3).

Put

$$(6.4) \quad \tilde{\phi}(r) = \inf_{v \in [r, \infty)} \phi(v).$$

Replacing ϕ by $\tilde{\phi}$, we may assume that ϕ is monotonically nondecreasing. Thinking of $\phi(r)$ as representing the trajectory of a car in front of us which is blocking the road, with our car's velocity bounded above by one, it is intuitively clear that we can find a trajectory $s(r)$ for our car such that

$\lim_{r \rightarrow \infty} s(r) = \infty$. More precisely, let $\rho \in C^\infty([0, 2])$ be a nondecreasing function which is identically zero near 0, identically one near 2 and satisfies $\rho'(x) \leq 1$ for all $x \in [0, 2]$. Put $r_0 = 0$ and define $\{r'_j\}_{j=0}^\infty$ and $\{r_j\}_{j=1}^\infty$ inductively by

$$(6.5) \quad \begin{aligned} r'_j &= \inf\{r : r \geq r_j + 2 \text{ and } \phi(r) \geq j + 1\}, \\ r_{j+1} &= r'_j + 2. \end{aligned}$$

Define s by

$$(6.6) \quad s(r) = \begin{cases} j & \text{if } r \in [r_j, r'_j] \\ j + \rho(r - r'_j) & \text{if } r \in [r'_j, r_{j+1}]. \end{cases}$$

Then s satisfies (6.3) and $\lim_{r \rightarrow \infty} s(r) = \infty$. \square

QUESTION. What can one say in the radially symmetric case without the assumption of a lower bound on the Ricci curvature?

Another interesting class of topologically tame manifolds consists of those which are hyperbolic, that is, of constant sectional curvature -1 . Complete hyperbolic manifolds are divided into those which are *geometrically finite* and those which are *geometrically infinite*. Roughly speaking, M is geometrically finite if its set of ends consists of a finite number of standard cusps and flares.

PROPOSITION 22 (Mazzeo-Phillips [23, Theorem 1.11]). *Let M be an infinite-volume geometrically finite hyperbolic manifold. If $\dim(M) = 2k$ then $\dim(\text{Ker}(\Delta_k)) = \infty$. If $\dim(M) = 2k + 1$ then $\sigma(\Delta_k) = \sigma(\Delta_{k+1}) = [0, \infty)$.*

The paper [23] also computes $\dim(\text{Ker}(\Delta_p))$ for such manifolds.

In general, geometrically infinite hyperbolic manifolds can have wild end behavior. However, in three dimensions one can show that the ends have a fairly nice structure. This is used to prove the next result.

PROPOSITION 23 (Canary [4, Theorem A]). *If M is a geometrically infinite topologically tame hyperbolic 3-manifold then $0 \in \sigma(\Delta_0)$.*

Proof. The method of proof is to show that M is not open at infinity and then apply Theorem 6. See [4] for details. \square

Thus zero lies in the spectrum of all topologically tame hyperbolic 3-manifolds. From Proposition 2, the same statement is true for compactly-supported modifications of such manifolds.

REFERENCES

- [1] AHLFORS L. and L. SARIO. *Riemann Surfaces*. Princeton University Press, Princeton (1960).
- [2] BROOKS, R. The Fundamental Group and the Spectrum of the Laplacian. *Comment. Math. Helv.* 56 (1981), 581–598.
- [3] BUSER, P. A Note on the Isoperimetric Constant. *Ann. Sci. Éc. Norm. Sup.* 15 (1982), 213–230.
- [4] CANARY, R. On the Laplacian and the Geometry of Hyperbolic 3-Manifolds. *J. Diff. Geom.* 36 (1992), 349–367.
- [5] CHEEGER, J. and M. GROMOV. L^2 -Cohomology and Group Cohomology. *Topology* 25 (1986), 189–215.
- [6] DE LA HARPE, P. and A. VALETTE. La Propriété (T) de Kazhdan pour les Groupes Localement Compacts. *Astérisque* 175 (1989), Paris.
- [7] DIXMIER, J. Cohomologie des Algèbres de Lie Nilpotentes. *Acta Sci. Math. Szeged* 16 (1955), 246–250.
- [8] DODZIUK, J. and W. KENDALL. Combinatorial Laplacians and Isoperimetric Inequality. In *From Local Times to Global Geometry, Control and Physics*, ed. K. Elworthy, Longman Scientific and Technical, Essex (1986), 68–74.
- [9] DONNELLY, H. The Differential Form Spectrum of Hyperbolic Space. *Manuscripta Math.* 33 (1981), 365–385.
- [10] —— Essential Spectrum and Heat Kernel *J. Funct. Anal.* 75 (1987), 362–381.
- [11] DRANISHNIKOV, A., S. FERRY and S. WEINBERGER. Large Riemannian Manifolds Which are Flexible. Preprint (1994).
- [12] FARRELL, F. and W.-C. HSIANG. Topological Characterization of Flat and Almost Flat Riemannian Manifolds M^n ($n \neq 3, 4$). *Amer. J. Math.* 105 (1983), 641–672.
- [13] GAFFNEY, M. A Special Stokes' Theorem for Complete Riemannian Manifolds. *Ann. of Math.* 60 (1954), 140–145.
- [14] GALLOT, S. and D. MEYER. Opérateur de Courbure et Laplacien des Formes Différentielles d'une Variété Riemannienne. *J. Math. Pures et Appl.* 54 (1975), 259–284.
- [15] GROMOV, M. Asymptotic Invariants of Infinite Groups. *Geometric Group Theory*, Vol. 2, ed. by Graham Niblo and Martin Roller, *London Math. Soc. Lecture Notes* 182, Cambridge University Press, Cambridge (1993).
- [16] GROMOV, M. and H. B. LAWSON. Positive Scalar Curvature and the Dirac Operator on Complete Riemannian Manifolds. *Publ. Math. IHES* 58 (1983), 83–196.
- [17] HAUSMANN, J.-C. and S. WEINBERGER. Caractéristiques d'Euler et Groupes Fondamentaux des Variétés de Dimension 4. *Comm. Math. Helv.* 60 (1985), 139–144.