

# 6. Embedding of tori

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

In particular, it follows that  $J(\omega)$ , the elliptic modular function [5], is real at  $\omega$ . In each case one has to determine the possible reflections  $\rho$ , determine their fixed-point sets, and add a suitable  $\tau_1$ .

We consider the rectangular case (1) of the lemma, for application in the next section. Let

$$(5.19) \quad \omega_1 = 1, \omega_2 = \omega = i\omega'', \omega'' > 1$$

be a normalized basis. For  $a = 1$ ,  $l_a$  is the real axis,  $\omega_0 = 0$ , or  $\omega_0 = 1$ ,  $b = ib_2$ , or  $b = \frac{1}{2} + ib_2$ ,  $0 \leq b_2 < \omega''$ . In the first case  $\omega'_0 = 0$ , or  $\omega'_0 = \omega$ , while there is no  $\omega'_0$  in the second case. Thus, we have

$$(5.20) \quad \rho(t) = \bar{t} + ib_2, \text{FP}(\rho) = \{\text{Im } t = b_2/2\} \cup \{\text{Im } t = (b_2 + \omega'')/2\}.$$

For  $a = -1$ ,  $l_a$  is the imaginary axis,  $\omega_0 = 0$  or  $\omega_0 = \omega$ ,  $b = b_1$ , or  $b = b_1 + i\omega''/2$ ,  $0 \leq b_1 < 1$ .  $\omega'_0 = 0, 1$  in the first case, and there is no  $\omega'_0$  in the second case. We have

$$(5.21) \quad \rho(t) = -\bar{t} + b_1, \text{FP}(\rho) = \{\text{Re } t = b_1/2\} \cup \{\text{Re } t = (b_1 + 1)/2\}.$$

If  $\varepsilon_1 = -1$ , then

$$(5.22) \quad \text{FP}(\tau_1) = \{c_1/2, (c_1 + \omega_1)/2, (c_1 + \omega_2)/2, (c_1 + \omega_1 + \omega_2)/2\}.$$

If we have  $\varepsilon_1 = +1$ ,  $2c_1 \in \Lambda$ ,  $c_1 \notin \Lambda$ , then  $\tau_1$  has no fixed points.  $\tau_1$  is then the deck transformation of an unbranched covering of another torus.

## 6. EMBEDDING OF TORI

We turn to the problem of concretely realizing the data of the previous section in the main case. Given a complex torus  $\Gamma = \mathbf{C}/\Lambda$ , with a pair of holomorphic involutions induced by

$$(6.1) \quad \tau_i(t) = -t + c_i, i = 1, 2,$$

we look for a pair of two-fold branched coverings

$$(6.2) \quad \pi_i: \Gamma \rightarrow \mathbf{P}_1, \pi_i \circ \tau_i = \pi_i, i = 1, 2.$$

The problem is immediately solved by taking

$$(6.3) \quad z_i = \pi_i(t) \equiv \mathcal{P}(t - c_i/2), i = 1, 2,$$

where

$$(6.4) \quad \mathcal{P}(t) = \frac{1}{t^2} + \sum_{\omega \in \Lambda - \{0\}} \left( \frac{1}{(t - \omega)^2} - \frac{1}{\omega^2} \right)$$

is the Weierstrass  $\mathcal{P}$ -function [5], [6]. We set

$$(6.5) \quad \pi(t) = (\pi_1(t), \pi_2(t)) .$$

If  $\pi(s_0) = \pi(t_0)$ ,  $s_0 \neq t_0$ , then  $s_0 \equiv -t_0 + c_i$ , mod  $\Lambda$ . Thus  $\pi$  will be one-to-one, as a map into  $\mathbf{P}_1 \times \mathbf{P}_1$ , if we assume

$$(6.6) \quad c_2 - c_1 \notin \Lambda .$$

To represent  $\pi$  as a map into  $\mathbf{P}_2$  with homogeneous coordinates  $\zeta$ ,  $z_1 = \zeta_1/\zeta_0$ ,  $z_2 = \zeta_2/\zeta_0$ , we again use the sigma function (4.13). We have [6]

$$(6.7) \quad \mathcal{P}(t) = -\partial_t^2 \log S(t) = -\frac{\Delta}{S(t)^2}, \quad \Delta = S(t)S''(t) - S'(t)^2 .$$

Since  $\Delta(0) = -S'(0)^2 \neq 0$ , we may write  $\pi$  as

$$(6.8) \quad \begin{aligned} \zeta_0 &= S(t - c_1/2)^2 S(t - c_2/2)^2 \\ \zeta_1 &= \Delta(t - c_1/2) S(t - c_2/2)^2 \\ \zeta_2 &= \Delta(t - c_2/2) S(t - c_1/2)^2 \end{aligned}$$

The branch points of the map  $\pi_1$  are given by (5.22), with  $\omega_1 = 1$ , and  $\omega_2 = \omega$ . By (6.6) the curve  $\pi$  has no finite singular points. Since  $\pi_i(t)$  has a pole of order two at  $t = c_i/2$ ,  $i = 1, 2$ ; the plane curve has two cusps on the line at  $\infty$  corresponding to these two parameter values. Such curves are considered in [3], for example.

To find the equation  $G(z_1, z_2) = 0$  of this plane curve, we change the variable,  $t \rightarrow t - c_1/2$ , so that  $G(\mathcal{P}(t), \mathcal{P}(t + c)) = 0$ , where

$$(6.9) \quad c = (c_1 - c_2)/2 .$$

We set

$$(6.10) \quad \begin{aligned} x &= \mathcal{P}(t + c), p = \mathcal{P}(t), p' = \mathcal{P}'(t), \\ \beta &= \mathcal{P}(c), \beta' = \mathcal{P}'(c) . \end{aligned}$$

The addition theorem and differential equation satisfied by  $\mathcal{P}$  [6] give

$$x + p + \beta = \frac{1}{4} \left( \frac{p' - \beta'}{p - \beta} \right)^2, \quad p'^2 = 4p^3 - g_2p - g_3 .$$

We rewrite these as

$$(p' - \beta')^2 = A(x, p), p'^2 = B(p) ,$$

and eliminate  $p'$ . This gives

$$(6.11) \quad F(x, p) \equiv F(x, p, \beta, \beta') \equiv (A - B - \beta'^2)^2 - 4\beta'^2 B = 0 .$$

Note that  $A - B$  is quadratic in  $p$ , and  $\beta'^2 = B(\beta)$ . Since  $F$  is an even function of  $\beta'$ , and  $\mathcal{P}$  is an even function, changing  $c$  to  $-c$  shows that we also have  $F(p, x) = 0$ . Since the coefficient of  $x^2$  in  $F$  is  $16(p - \beta)^2$ , we must have

$$F(x, p) = G(x, p) (p - \beta)^2 .$$

Expanding in powers of  $p - \beta$  gives

$$(6.12) \quad \begin{aligned} F(x, \beta) &= 0, \partial_p F(x, \beta) = 0 , \\ G(x, p) &= (1/2) \partial_p^2 F(x, \beta) + (1/6) \partial_p^3 F(x, \beta) (p - \beta) \\ &\quad + (1/24) \partial_p^4 F(x, \beta) (p - \beta)^2 . \end{aligned}$$

After some computation we get

$$(6.13) \quad \begin{aligned} G(z_1, z_2) &= (z_1 - \beta)^2 (z_2 - \beta)^2 + \beta_1 (z_1 - \beta) (z_2 - \beta) \\ &\quad + \beta_2 (z_1 + z_2 - 2\beta) + \beta_3 , \end{aligned}$$

where

$$(6.14) \quad \begin{aligned} \beta_1 &= -(12\beta^2 - g_2)/2, \beta_2 = -B(\beta), \\ \beta_3 &= (12\beta^2 - g_2)^2 - 3\beta B(\beta) . \end{aligned}$$

Next we consider the reality condition (3.11). From (5.9) and (6.4) we get

$$(6.15) \quad \overline{\mathcal{P}(t)} = a^2 \mathcal{P}(a\bar{t}) .$$

By definition  $g_2 = 60G_2$ ,  $g_3 = 140G_3$ , where [6]

$$G_k = \sum_{\omega \in \Lambda - \{0\}} \frac{1}{\omega^{2k}} .$$

It follows from (5.9) that  $\bar{G}_k = a^{2k} G_k$ , so that

$$(6.16) \quad \bar{g}_2 = a^4 g_2, \bar{g}_3 = a^6 g_3 .$$

By (5.7) we have  $c = (c_1 - a\bar{c}_1 - b + a\bar{b})/2$ , so that  $a\bar{c} = -c$ . Hence,

$$(6.17) \quad \bar{\beta} = a^2 \beta, \bar{\beta}_1 = a^4 \beta_1, \bar{\beta}_2 = a^6 \beta_2, \bar{\beta}_3 = a^8 \beta_3 .$$

To satisfy (3.11) we *redefine*

$$(6.18) \quad \pi_i(t) = a \mathcal{P}(t - c_i/2),$$

and set

$$(6.19) \quad G_0(z_1, z_2) = a^4 G(z_1/a, z_2/a),$$

so that

$$(6.20) \quad \overline{G_0(z_1, z_2)} = G_0(\bar{z}_2, \bar{z}_1).$$

In summary we have

**PROPOSITION 6.1.** *Let  $\Lambda = \mathbf{C}/\Lambda$  have the holomorphic involutions (6.1) intertwined by the anti-holomorphic involution (5.6). Then  $(\Gamma, \rho, \tau_i)$  is realized by the map (6.5), (6.18) onto the quartic curve  $G_0(z_1, z_2) = 0$  given by (6.13), (6.14), (6.19). If the fixed-point set of  $\rho$  is non-empty, then this is the complexification of the real curve  $G_0(z, \bar{z}) = 0$ .*

## 7. A RECTANGULAR LATTICE

We consider the special case of  $\Lambda, \rho, \tau_i$  as given in (5.19), (5.6), (6.1), with

$$(7.1) \quad a = +1, b = 0, \bar{c}_2 = c_1 = c'_1 + i c''_1, c = i c''_1.$$

From (6.16), (6.15) it follows that  $g_2, g_3, \beta$  are real, and  $\beta'$  is purely imaginary. Thus, the coefficients  $\beta_1, \beta_2, \beta_3$  of  $G(z_1, z_2)$  are real. With  $t = t' + i t''$ , we have

$$(7.2) \quad FP(\rho) = \{t'' = 0\} \cup \{t'' = \omega''/2\},$$

$$(7.3) \quad \tau_1\{t'' = 0\} = \{t'' = c''_1\}, \quad \tau_1\{t'' = \omega''/2\} = \{t'' = c''_1 + \omega''/2\}.$$

Let us assume that  $0 < c''_1 < \omega''/2$ . Then the torus  $\Lambda$  is divided into four annuli

$$A_1 = \{0 < t'' < c''_1\}, \quad A_2 = \{c''_1 < t'' < \omega''/2\},$$

$$A_3 = \{\omega''/2 < t'' < c''_1 + \omega''/2\}, \quad A_4 = \{c''_1 + \omega''/2 < t'' < \omega''\}.$$

The fixed points of  $\tau_1$  are, by (5.22),

$$(7.4) \quad c_1/2, (c_1 + 1)/2 \in A_1,$$

$$(7.5) \quad (c_1 + i\omega'')/2, (c_1 + 1 + i\omega'')/2 \in A_3.$$