

# §5. Additive functions

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

### § 5. ADDITIVE FUNCTIONS

As we noted in the first section, the study of uniform distribution on divisors of additive functions is made much easier by the fact that the Weyl sums are multiplicative. Indeed this is the only case when we are able to achieve estimates for the discrepancy which go beyond the statistical bound  $\sqrt{\tau(n)}$ .

We shall prove the following theorem, which provides a simple, but nevertheless effective criterion. We write for integer  $v$

$$(102) \quad L_v(x; f) := \sum_{p \leq x} \frac{\|vf(p)\|^2}{p}, \quad C_v(x; f) = \sum_{p \leq x} \frac{1 - |\cos v\pi f(p)|}{p},$$

and note that we have for all  $x$

$$(103) \quad 4L_v(x; f) \leq C_v(x; f) \leq \frac{1}{2} \pi^2 L_v(x; f).$$

**THEOREM 13.** *Let  $f$  be an additive function. Then  $f$  is erd if, and only if,*

$$(104) \quad \sum_p \frac{\|vf(p)\|^2}{p} = \infty \quad (v \neq 0).$$

*Moreover, if this is the case then we have*

$$(105) \quad \sum_{n \leq x} \frac{\Delta(n; f)}{n} \left(\frac{y}{2}\right)^{\Omega(n)} \ll (\log x)^y \left\{ \frac{1}{T} + \sum_{1 \leq v \leq T} \frac{e^{-yC_v(x; f)}}{v} \right\}$$

*uniformly for  $x \geq 2$ ,  $T \geq 1$ ,  $0 \leq y \leq y_0 < 4$ .*

The qualitative result corresponding to this statement (i.e. the criterion (104)) was first established by Kátai [18].

By (104) and the lower bound of (103), the expression between curly brackets tends to zero for suitable  $T = T(x, y)$ . Hence (105) provides, as described in Theorem 6, a family of effective ppl estimates for  $\Delta(n; f)$  which may of course be further optimised with respect to  $y$ . This process must lead to a non trivial result since  $y = 1$  is admissible.

We give below two applications of Theorem 13, respectively devoted to the functions  $f(d) = \theta\Omega(d)$  for irrational  $\theta$ , and  $f(d) = \log d$ . Recovering a result first obtained independently by Hall and Kátai, the latter case furnishes the best known ppl upper bound for the discrepancy for any function we are aware of, although this falls short of the current conjecture stated in section 2. The former case provides an example of a function which

is erd but has the same rate of growth than, and indeed is asymptotically equal to, a function which is not, namely  $\theta \log_2 n$  — see Corollary 3. Actually, if we define  $\Omega(n; E)$  as the number of those prime factors of  $n$  which belong to some set of primes  $E$ , then Theorem 13 implies that  $\theta\Omega(n; E)$  is erd for all  $\theta \in \mathbf{R} \setminus \mathbf{Q}$  if, and only if,  $\sum_{p \in E} 1/p = \infty$ . We thus exhibit erd additive functions with an arbitrarily slow growth.

The effective bounds for  $\Delta(n; \theta\Omega)$  naturally depend on the rational approximations to  $\theta$ . We set  $Q_1(x) := \sqrt{\log_2 x / \log_3 x}$ , and define

$$(106) \quad q_1(x; \theta) := \inf_{t \geq x} \max \{q \leq Q_1(t) : \| \theta q \| \leq 1/Q_1(t)\}.$$

Then  $q_1(x; \theta)$  increases to  $\infty$  for all irrational  $\theta$  and furthermore  $q_1(x; \theta) = (\log_2 x)^{1/2 + o(1)}$  for almost all  $\theta$ .

COROLLARY 10. *Let  $\theta \in \mathbf{R} \setminus \mathbf{Q}$ . Then the function  $\theta\Omega(n)$  is erd and we have*

$$(107) \quad \Delta(n; \theta\Omega) < \tau(n) q_1(n; \theta)^{-1 + o(1)} \text{ ppl}.$$

COROLLARY 11 (Hall [8]; Kátai [17]). *Let  $\alpha > (\log(4/\pi))/\log 2 \approx 0.34850$ . We have*

$$(108) \quad \Delta(n; \log) < \tau(n)^\alpha \text{ ppl}.$$

We now embark on the proof of Theorem 13. That the condition is necessary is a straightforward consequence of the definition of uniform distribution on divisors in the form (1). Indeed, suppose that (104) fails to hold for  $v \neq 0$ . Writing  $F(z; n) := \sum_{d|n, (f(d)) \leq z} 1$ , we have

$$\begin{aligned} g_v(n) &:= \sum_{d|n} e(vf(n)) = \int_0^1 e(vz) dF(z; n) = \int_0^1 e(vz) d(F(z; n) - z\tau(n)) \\ &= 2\pi i v \int_0^1 e(vz) (F(z; n) - z\tau(n)) dz, \end{aligned}$$

hence

$$(109) \quad 2\pi |v| |\Delta(n; f)| \geq |g_v(n)|.$$

Now for all  $\varepsilon \in ]0, 1[$ , we have

$$\begin{aligned} (110) \quad \sum_{n \leq x} \frac{\mu(n)^2 |g_v(n)|}{n\tau(n)} &\geq \sum_{P^+(n) \leq x^\varepsilon} \frac{\mu(n)^2 |g_v(n)|}{n\tau(n)} - \sum_{\substack{n > x \\ P^+(n) \leq x^\varepsilon}} \frac{1}{n} \\ &= \prod_{p \leq x^\varepsilon} \left( 1 + \frac{|\cos \pi vf(p)|}{p} \right) - O(e^{-1/2\varepsilon} \log x), \end{aligned}$$

where the  $O$ -estimate follows from (94) by partial summation. The product is

$$\gg \exp \left\{ \sum_{p \leq x^\varepsilon} \frac{|\cos \pi v f(p)|}{p} \right\} \geq \exp \left\{ \sum_{p \leq x^\varepsilon} \frac{1 - \frac{1}{2} \pi^2 \|vf(p)\|^2}{p} \right\} \gg \varepsilon \log x ,$$

by our assumption that the series (104) converges. Inserting this into (110) and choosing  $\varepsilon$  small enough but fixed, we obtain that the left-hand side is  $\gg \log x$ , whence by (109)

$$\sum_{n \leq x} \frac{\Delta(n; f)}{n \tau(n)} \gg \log x .$$

This cannot hold if  $\Delta(n; f) = o(\tau(n)) \text{pp}$  (or even  $\text{ppl}$ ), and we obtain the required necessity assertion.

The sufficiency part of the theorem readily follows from the upper bound (21) which we recall for convenience: we have uniformly for  $x \geq 2$ ,  $T \geq 1$ ,  $0 \leq y \leq y_0 < 4$ ,

$$(111) \quad \sum_{n \leq x} \frac{\Delta(n; f)}{n} \left( \frac{y}{2} \right)^{\Omega(n)} \ll \frac{(\log x)^y}{T} + \sum_{1 \leq v \leq T} \frac{1}{v} \sum_{n \leq x} \frac{|g_v(n)|}{n} \left( \frac{y}{2} \right)^{\Omega(n)} .$$

The inner  $n$ -sum on the right does not exceed

$$\sum_{P^+(n) \leq x} \frac{|g_v(n)|}{n} \left( \frac{y}{2} \right)^{\Omega(n)} \ll \exp \left\{ \sum_{p \leq x} \frac{y |\cos \pi v f(p)|}{p} \right\} \ll (\log x)^y e^{-y C_v(x; f)} .$$

Inserting this into (111) yields (105). Taking  $y = 1$  implies  $\Delta(n; f) = o(2^{\Omega(n)}) \text{ppl}$ , from which we deduce in turn that  $\Delta(n; f) = o(\tau(n)) \text{ppl}$  by a familiar argument. Corollary 1 hence implies that  $f$  is erd, and this finishes the proof of Theorem 13.

*Proof of Corollary 10.* We apply the upper bound (105) of Theorem 13 with  $y = 1$ . We have

$$C_v(x; \theta \Omega) = (1 - |\cos \pi v \theta|) \log_2 x + O(1)$$

and need a lower bound for this. For given  $x$ , there exist integers  $a, q$  with  $(a, q) = 1$ ,  $q \leq Q_1(x) = \sqrt{\log_2 x / \log_3 x}$ , and  $|\theta - a/q| \leq 1/q Q_1(x)$ . Furthermore, we have

$$(112) \quad q \geq q_1(x; \theta) ,$$

where the right-hand side is defined by (106). We select  $T := q / \log q$ , and note that for  $v \leq T$  we have  $|\theta v - a_v/q_v| \leq T/q Q_1(x) \leq 1/q \log q$ , with  $a_v = av/(q, v)$ ,  $q_v = q/(q, v)$ . This implies  $\|\theta v\| \geq (1/q) - 1/q \log q$ , hence, for large  $x$ ,

$$1 - |\cos \pi v \theta| \geq \left( \frac{1}{2} \pi^2 + o(1) \right) / q^2 > (\log_3 x) / \log_2 x .$$

Inserting this into (105), we arrive at

$$(\log x)^{-1} \sum_{n \leq x} \frac{\Delta(n; \theta\Omega)}{n 2^{\Omega(n)}} \ll \frac{1}{T} + \frac{\log q}{\log_2 x} \ll q_1(x; \theta)^{-1+o(1)}.$$

Since  $q_1(x; \theta)$  is a non-decreasing function of  $x$ , this implies (107) and the proof is thereby completed.

*Proof of Corollary 11.* Put  $\tau(n, \theta) := \sum_{d|n} d^{i\theta}$ . When  $f = \log$ , we have that  $g_v(n) = \tau(n, 2\pi v)$ . By lemma 30.2 of [14] we infer that, uniformly for  $1 \leq |\theta| \leq \exp/\sqrt{\log x}$ ,

$$\sum_{p \leq x} \frac{|\tau(p, \theta)|}{p} = \sum_{p \leq x} \frac{|\cos(\frac{1}{2}\theta \log p)|}{p} = \frac{2}{\pi} \log x + O(1).$$

This is proved by partial summation from a strong form of the prime number theorem. Thus we obtain that we have uniformly for  $1 \leq v \leq \log x$

$$C_v(x; \log) = (1 - 2/\pi) \log_2 x + O(1).$$

Inserting this into (105) with  $T = \log x$  and choosing optimally  $y = \pi/2$ , we obtain

$$\Delta(n; f) < \xi(n) (\log_2 n) \left(\frac{4}{\pi}\right)^{\Omega(n)} \text{ppl}$$

for all  $\xi(n) \rightarrow \infty$ . This implies the required result by a now standard device.

## 6. METRIC RESULTS

In this last section, we investigate the problem of uniform distribution on divisors from a further statistical point of view, regarding as random not only the integer  $n$  but also the function  $f$ . Thus, we define a measure  $\mu$  on the set  $\mathbf{A}$  of all real valued arithmetical function as the inverse image of the Haar measure on the compact group  $(\mathbf{R}/\mathbf{Z})^N$  by the canonical mapping  $f \mapsto \langle f \rangle$ . In other words,  $\mu$  is characterised by the property that for all finite families  $\{E_j : 1 \leq j \leq k\}$  of measurable subsets of the torus  $\mathbf{R}/\mathbf{Z}$  and for all integers  $n_1, n_2, \dots, n_k$ , we have

$$\mu \{f \in \mathbf{A} : \langle f(n_j) \rangle \in E_j \ (1 \leq j \leq k)\} = \prod_{j=1}^k \lambda(E_j),$$