

### **3. ESTIMATES FOR THE BOUNDARY SUBPOLYHEDRA**

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **42 (1996)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.05.2024**

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

#### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Each polyhedron  $V(s)$  is homotopically equivalent to  $V$ . More precisely we have

**PROPOSITION 2.3.** *For every sufficiently large  $s$  the locally symmetric space  $V$  is homeomorphic to the interior of the polyhedron  $V(s)$  in  $V$ , and  $V(s)$  is a strong deformation retract of  $V$ .*

For the proof see [L3], Theorems 5.2 and 5.5.

### 3. ESTIMATES FOR THE BOUNDARY SUBPOLYHEDRA

We wish to apply Proposition 1.1 to the polyhedra  $V(s)$  in the above exhaustion and then take the limit for  $s \rightarrow \infty$ . To that end we need estimates for the second fundamental forms and the volumes of the (lower dimensional) boundary polyhedra.

For each Siegel set  $\mathcal{S}_i := q_i \mathcal{S}$  which is part of the fundamental set  $\Omega$  we have its truncated part

$$\mathcal{S}_i(s) := \mathcal{S}_i - \bigcup_{\alpha \in \Delta} (\mathcal{B}_{i\alpha}(s) \cap \mathcal{S}_i) .$$

The top dimensional boundary faces of  $\mathcal{S}_i(s)$  in  $\mathcal{S}_i$  (resp. of  $\Omega(s)$  in  $\Omega$ ) are subsets of horospheres :

$$\mathcal{H}_{i\alpha}(s) := \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\} \cap \mathcal{S}_i(s) , \quad \alpha \in \Delta .$$

The “horospherical” pieces  $\mathcal{H}_{i\alpha}(s)$  together with their  $\Gamma$ -translates form the boundary of the manifold with corners  $X(s)$  in  $X$ . For any nonempty subset  $\Theta$  of  $\Delta$  we set

$$\mathcal{H}_{i\Theta}(s) := \bigcap_{\alpha \in \Theta} \mathcal{H}_{i\alpha}(s) \subset \mathcal{S}_i(s) .$$

The various boundary subpolyhedra of  $V(s)$  are then unions of projections of the pieces  $\mathcal{H}_{i\Theta}(s)$  under the canonical projection  $\pi : X \rightarrow V$ . More precisely, as explained in Section 2, for any subset  $\Theta \subset \Delta$ , we have the equivalence relation on the set  $I = \{1, \dots, m\}$

$$j \sim_\Theta l \quad \text{if and only if} \quad \Gamma q_j P_\Theta = \Gamma q_l P_\Theta$$

(the  $q_i$  are as in Proposition 2.1). This relation  $\sim_\Theta$  induces a partition,  $I(\Theta)$ , of the set  $I$  whose components will be denoted by  $E$ . Let  $n = \dim X = \dim V$ , let  $k$  be the cardinality of  $\Theta$  and let  $E \in I(\Theta)$ . Then  $V_E^{n-k}(s) := \pi(\bigcup_{i \in E} \mathcal{H}_{i\Theta}(s))$  is a  $(n-k)$ -dimensional boundary polyhedron of  $V(s)$ ; and moreover, any boundary polyhedron arises in this way (see [L3] §4).

REMARK. The minimal possible dimension which occurs is  $n - q$  where  $q$  is the  $\mathbb{Q}$ -rank of  $\mathbf{G}$ . It is also interesting to note (though not needed below) that the outer angles are isomorphic to  $\mathbb{Q}$ -Weyl chambers and their walls at infinity.

We shall use the following well-known fact about Jacobi fields in symmetric spaces (see [K] Theorem 2.2.9). A Jacobi field along a geodesic ray is called *stable* if its length is bounded.

LEMMA 3.1. *Let  $r : [0, \infty) \rightarrow X$  be a unit-speed geodesic ray in the symmetric space  $X$  (of noncompact type). Set  $p = r(0)$ . Then the unique stable Jacobi field  $J_u(s)$  along  $r(s)$  with  $J_u(0) = u \in T_p X$  can be written as*

$$J_u(s) = \sum_j e^{-\lambda_j s} a_j E_j(s)$$

where  $\{E_j(s)\}$  is an orthonormal frame of parallel fields along  $r$ , the  $\lambda_j$  are non-negative (uniform) constants and  $u = \sum_j a_j E_j(0)$ .

LEMMA 3.2. *Let  $s \geq 0$ . The second fundamental forms of every boundary polyhedron  $V_E^{n-k}(s)$  with respect to outer angles in  $V(s)$  are uniformly bounded by a constant independent of  $E, k$  and  $s$ .*

*Proof.* Since the claim is local we can work in the universal covering space  $X$ . As we noted above the preimage of  $V_E^{n-k}(s)$  in  $X$  under the projection  $\pi$  is the union of a finite number of horospherical sets

$$\mathcal{H}_{i\Theta}(s) = \bigcap_{\alpha \in \Theta} \mathcal{H}_{i\alpha}(s) \subset \bigcap_{\alpha \in \Theta} \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\},$$

where  $\Theta$  is a subset of  $\Delta$  with  $k$  elements. The (inner) unit normal field of the horosphere  $\{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\}$  is given by  $Z_{i\alpha} := -\text{grad } \tilde{h}_{i\alpha}$  (see e.g. [HI] Proposition 3.1). Using  $d\pi$  any element in the outer angle  $O(\pi(p))$  of  $V_E^{n-k}(s)$  at a point  $\pi(p) \in V_E^{n-k}(s)$  can then be identified with a *positive* linear combination (of norm 1) of the radial fields  $Z_{i\alpha}(p)$ ,  $\alpha \in \Theta$ . It therefore suffices to show that for any pair  $(i, \alpha)$  the second fundamental form of  $V_E^{n-k}(s)$  relative to  $d\pi Z_{i\alpha}$  is uniformly bounded. We fix  $i$  and  $\alpha$  and write  $Z$  for  $Z_{i\alpha}$ . For  $p \in X$  let  $\langle \cdot, \cdot \rangle_p$  denote the Riemannian metric of  $X$  at  $p$ . Let  $u, v \in T_p X$  be such that  $d\pi(u), d\pi(v) \in T_{\pi(p)} V_E^{n-k}(s)$ . Using the above identifications the second fundamental form of  $V_E^{n-k}(s) \subset V(s)$  with respect to  $Z$  can be written as

$$\Pi_Z(u, v)(p) = \langle D_u Z, v \rangle_p .$$

According to [HI], Proposition 3.1, we have  $D_u Z(p) = J'_u(0)$  where  $J_u$  is the stable Jacobi field along the (unique) geodesic ray, say  $r$ , in  $X$  which joins  $p$  to  $c_{i\alpha}(\infty) \in \partial_\infty X$  and with initial value  $J_u(0) = u$ . By Lemma 3.1 there are orthonormal parallel fields  $E_j(s)$  along  $r$  and constants  $\lambda_j \geq 0$  such that  $J_u(s) = \sum_j e^{-\lambda_j s} a_j E_j(s)$  with  $u = \sum_j a_j E_j(0)$ . Consequently we get  $J'_u(0) = -\sum_j \lambda_j a_j E_j(0)$  and finally, for  $v = \sum_j b_j E_j(0)$ ,  $|\Pi_Z(u, v)(p)| = \left| -\sum_j \lambda_j a_j b_j \right| \prec \|u\| \|v\|$ .  $\square$

We next estimate the volumes of the boundary polyhedra. Recall from Section 2.1 the parametrization of  $X$  by horocyclic coordinates

$$\mu: Y = U \times Z \times A \longmapsto X; (u, \tau(m), a) \longmapsto um a \cdot x_0.$$

Let  $dx^2$  be the  $G$ -invariant Riemannian metric on  $X$  induced by the Cartan-Killing form of the Lie algebra  $\mathfrak{g}$  of  $G$  and let  $dz^2$  be the invariant metric on  $Z$ . Further let  $da^2$  (resp.  $du^2$ ) be the left-invariant metric on  $A$  (resp.  $U$ ). Finally set  $dy^2 := \mu^* dx^2$ .

**LEMMA 3.3.** *Let  $dv_Y$ ,  $dv_U$ ,  $dv_Z$  and  $dv_A$  denote the volume elements of the metrics  $dy^2$ ,  $du^2$ ,  $dz^2$  and  $da^2$ . Then at the point  $(u, z, a) \in Y$  we have*

$$2^e dv_Y = \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_A$$

where  $e = \frac{1}{2} \dim U$  and  $\rho$  is the sum of all positive roots (counted with multiplicity); it can be written in the form  $\rho = \sum_{\alpha \in \Delta} c_\alpha \alpha$ ,  $c_\alpha > 0$ .

For the proof see [B2] Corollary 4.4.

**LEMMA 3.4.** *For the  $(n-k)$ -dimensional volume of each boundary polyhedron  $V_E^{n-k}(s)$  of  $V(s)$  one has the estimate*

$$\text{Vol}(V_E^{n-k}(s)) \prec s^{q-k} e^{-cs},$$

where  $q = \dim A$  is the  $\mathbb{Q}$ -rank of  $\mathbf{G}$  and  $c > 0$  is a constant (independent of  $E, k$  and  $s$ ).

*Proof.* We again consider the preimage of  $V_E^{n-k}(s)$  in  $X$  under the map  $\pi$ . We need to estimate the volume of each horospherical piece

$$\mathcal{H}_{i\Theta}(s) = \bigcap_{\alpha \in \Theta} \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\} \cap \mathcal{S}_i(s), \quad i \in E.$$

It clearly suffices to carry out the estimates for  $i = 1$ ; note that  $q_1 = e$ . For the horocyclic coordinate map  $\mu: Y \rightarrow X$  and the canonical projection

$\pi_A : Y \rightarrow A$  we set  $A_\Theta(s) := \pi_A \circ \mu^{-1}(\mathcal{H}_{1\Theta}(s)) \subset A$ . The set  $A_\Theta(s)$  is contained in an “affine” subspace of  $A$  of the form  $a_1 a_*(s) A^{q-k}$  where  $a_1 a_*(s) \in A$  and  $A^{q-k}$  is a  $q-k$ -dimensional subgroup of  $A$  (see Sections 3 and 4 of [L2]). We denote the restriction of  $dv_A$  to  $A^{q-k}$  by  $dv_{A^{q-k}}$ ; for  $k = q$  we have  $A^0 = e$  and we set  $dv_{A^0} \equiv 1$ . By Lemma 3.3 we have (for  $k$  equal to the number of elements of  $\Theta$ )

$$\text{Vol}(V_E^{n-k}(s)) \prec \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}}.$$

Since the horospherical piece  $\mathcal{H}_{1\Theta}(s)$  is part of a Siegel set  $\mathcal{S}_{\omega,\tau}$  with  $\omega$  relatively compact (and hence of finite volume) in  $UM$  we get

$$\begin{aligned} \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}} &\prec \\ &\prec \int_{\omega} dv_U \wedge dv_Z \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}} \prec \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}}. \end{aligned}$$

Also by definition of a Siegel set we have  $\alpha(a) \geq \tau \succ 1$  for all  $\alpha \in \Delta$ . Moreover, the computations in the proof of Lemma 4.1 (and Lemma 3.5) in [L2] show that for all  $\alpha \in \Theta$  one has  $\alpha(a_1 a_*(s)) \succ e^{\mu_\alpha s}$  with  $\mu_\alpha > 0$ . Hence, as  $\Theta \subset \Delta$  is not empty and since  $\rho = \sum_{\alpha \in \Delta} c_\alpha \alpha$  ( $c_\alpha > 0$ ), there is a uniform constant  $c > 0$  such that  $\rho(a)^{-1} \prec e^{-cs}$  for all  $a \in A_\Theta(s)$ . As noted above the set  $A_\Theta(s)$  is contained in a  $(q-k)$ -dimensional affine cone in  $A$ . It is similar (in the sense of Euclidean geometry) to  $A_\Theta(0)$  with similarity factor  $s$  (see the proof of Lemma 4.1 in [L2]). Hence we eventually get  $\int_{A_\Theta(s)} dv_{A^{q-k}} \prec s^{q-k}$  and the Lemma follows.  $\square$

#### 4. A NEW PROOF OF THE GAUSS-BONNET FORMULA

In this section we present a new simplified proof of the Gauss-Bonnet theorem for higher rank locally symmetric spaces.

**THEOREM 4.1.** *Let  $X$  be a Riemannian symmetric space of noncompact type and  $\mathbb{R}$ -rank  $\geq 2$  and let  $\Gamma$  be an irreducible, torsion-free (non-uniform) lattice in the group of isometries of  $X$ . Then for the locally symmetric space  $V = \Gamma \backslash X$  the Gauss-Bonnet formula holds :*

$$\chi(V) = \int_V \Psi dv.$$