

# 4. POLYGONS WITH GIVEN SIDES – KÄHLER STRUCTURES

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

section collapsed. The trace function on  $\mathcal{M}_{m \times 2}(\mathbf{C})$  descends to  $\tilde{\mathbf{G}}_2(\mathbf{C}^m)$  and to the Casimir function “perimeter” on  ${}^m\mathcal{PP}_+^3$ .

#### 4. POLYGONS WITH GIVEN SIDES – KÄHLER STRUCTURES

We now use the map  $\ell : {}^m\tilde{\mathcal{P}}^k, {}^m\mathcal{P}_+^k, {}^m\mathcal{P}^k \rightarrow \mathbf{R}^m$  defined in (2.4). Recall that  $\ell(\rho)$ , for  $\rho \in {}^m\tilde{\mathcal{P}}^k$ , is the length of the successive sides of a representative of  $r$  with total perimeter 2.

For  $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbf{R}_{\geq 0}^m$  with  $\sum_{i=1}^m \alpha_i = 2$ , we define

$${}^m\tilde{\mathcal{P}}^k(\alpha) :=: \tilde{\mathcal{P}}^k(\alpha) := \{\rho \in {}^m\tilde{\mathcal{P}}^k \mid \ell(\rho) = \alpha\} \subset {}^m\tilde{\mathcal{P}}^k.$$

The space  $\tilde{\mathcal{P}}^k(\alpha)$  is invariant under the action of  $O_k$ . We define the moduli spaces

$$\mathcal{P}_+^k(\alpha) := SO_k \backslash \tilde{\mathcal{P}}^k(\alpha) = \ell^{-1}(\alpha) \subset {}^m\mathcal{P}_+^k$$

and

$$\mathcal{P}^k(\alpha) := O_k \backslash \tilde{\mathcal{P}}^k(\alpha) = \ell^{-1}(\alpha) \subset {}^m\mathcal{P}^k.$$

The space  $\tilde{\mathcal{P}}^1(\alpha)$  consists of a finite number of points and is generically empty. We call  $\alpha$  *generic* if  $\tilde{\mathcal{P}}^1(\alpha) = \emptyset$ .

**THEOREM 4.1.** *The map  $\mu := \ell \circ \hat{\Phi} : \mathbf{G}_2(\mathbf{C}^m) \longrightarrow \mathbf{R}^m$  is a moment map for the action of  $U_1^m$  on  $\mathbf{G}_2(\mathbf{C}^m)$ .*

*Proof.* As seen in (3.13), the moment map  $\Psi : \mathbf{G}_2(\mathbf{C}^m) \longrightarrow \mathcal{H}(m)$  for the  $U_m$ -action on  $\mathbf{G}_2(\mathbf{C}^m)$  is induced from  $\tilde{\Psi} : \mathcal{M}_{m \times 2}(\mathbf{C}) \longrightarrow \mathcal{H}(m)$  given by  $\tilde{\Psi}(a, b) := (a, b) \cdot (a, b)^*$ . A moment map  $\mu$  for the action of  $U_1^m$  is obtained by composing  $\Psi$  with the projection  $\mathcal{H}(m) \longrightarrow \mathbf{R}^m$  associating to a matrix its diagonal entries. So, if  $\Pi \in \mathbf{G}_2(\mathbf{C}^m)$  is generated by  $a$  and  $b$  with  $(a, b) \in \mathbf{V}_2(\mathbf{C}^m)$ , one has

$$\mu(\Pi) = (|a_1|^2 + |b_1|^2, \dots, |a_m|^2 + |b_m|^2) = \ell \circ \hat{\Phi}(a, b). \quad \square$$

A now classic theorem of Atiyah and Guillemin-Sternberg [Au, §III.4.2] asserts that the image of a moment map for a torus action is a convex polytope (the *moment polytope*). The restriction of the moment map to the fixed point set of an anti-symplectic involution has the same image [Du]. In our case, one gets these facts directly:

COROLLARY 4.2. *The moment map  $\mu: \mathbf{G}_2(\mathbf{C}^m) \longrightarrow \mathbf{R}^m$  satisfies  $\mu(\mathbf{G}_2(\mathbf{C}^m)) = \mu(\mathbf{G}_2(\mathbf{R}^m)) = \Xi_m$ , where  $\Xi_m$  is the hypersimplex*

$$\Xi_m := \{(x_1, \dots, x_m) \in \mathbf{R}^m \mid 0 \leq x_i \leq 1 \text{ and } \sum_{i=1}^m x_i = 2\}.$$

*Proof.* One has  $\text{Image}(\mu) = \text{Image}(\ell)$ . Further it is manifest that  $\text{Image}(\ell) \subset \Xi_m$ . A proof that  $\text{Image}(\ell) = \Xi_m$  is actually provided in [KM1], Lemma 1, or [Ha]. We give here however another argument, for the pleasure of constructing a continuous section  $\sigma: \Xi_m \longrightarrow {}^m\mathcal{P}^2$  of  $\ell$ . If  $m = 3$ , we have already mentioned in (2.7) that  ${}^3\mathcal{P}^2$  is homeomorphic to  $\Xi_3$  via the map  $\ell$ . Let  $\alpha \in \Xi_m$ . Define  $\beta_i := \sum_{j=1}^i \alpha_j$  and

$$r(\alpha) := \min\{i \mid \beta_i \leq 1 \text{ and } \beta_{i+1} \geq 1\}.$$

The numbers  $\beta_r, \alpha_r, 2 - \beta_{r+1}$  form a triple of  $\Xi_3$  and are then the lengths of a unique triangle  $\tau(\alpha) \in {}^3\mathcal{P}^2$ , which can be subdivided in the obvious way to define the element  $\sigma(\alpha) \in {}^m\mathcal{P}^2(\alpha)$  (see Figure 1).




FIGURE 1:  $\tau(\alpha)$

The continuity of  $\sigma$  comes from the fact that if the map  $r$  is discontinuous at some  $\alpha$ , the triangle  $\tau(\alpha)$  is then lined.  $\square$

REMARKS. 1) Corollary 4.2 is also a consequence of our stronger result (5.4).

2) The word “hypersimplex” is introduced in [GM]. Observe that  $H$  is obtained by taking the convex hull of the middle point of each edge of a standard  $(m - 1)$ -simplex.

We also obtain the critical values of  $\mu$  (compare [Ha]):

PROPOSITION 4.3. *The set of critical values of  $\mu$  on  $\mathbf{G}_2(\mathbf{C}^m) \rightarrow \Xi_m$  or  $\mathbf{G}_2(\mathbf{R}^m) \rightarrow \Xi_m$  consists of those points  $(x_1, \dots, x_m) \in \Xi_m$  satisfying one of the following conditions:*

- a) *one  $x_i$  vanishes;*
- b) *one  $x_i$  is equal to 1;*
- c) *there exist  $\varepsilon_i = \pm 1$  such that  $\sum_{i=1}^m \varepsilon_i x_i = 0$ , with at least two  $\varepsilon_i$ 's of each sign.*

REMARK. Points satisfying a) and b) constitute the boundary of  $\Xi_m$ . Points satisfying c) are “inner walls”. Points satisfying a) correspond to non-proper polygons. Those satisfying b) or c) are non-generic  $\alpha$ 's (Condition b) implies that there exist  $\varepsilon_i = \pm 1$  such that  $\sum_{i=1}^m \varepsilon_i x_i = 0$  with all but one  $\varepsilon_i$  of the same sign.)

*Proof.* The critical points of the moment map  $\mu$  are the points of  $\mathbf{G}_2(\mathbf{C}^m)$  for which the  $U_1^m$ -action has a stabilizer of dimension bigger than 1. They are the images of those  $(2 \times m)$ -matrices in  $\mathbf{V}_2(\mathbf{C}^m)$  for which the  $(U_1^m \times_{U_1} U_2)$ -action has a non-discrete stabilizer. There are such points whose stabilizer is contained in  $U_1^m \times \{1\}$ ; they are the matrix with one row vanishing and their values under  $\mu$  are the points of  $\Xi_m$  satisfying a). The other points give rise to points in  ${}^m\tilde{\mathcal{P}}^3 = U_1^m / \mathbf{V}_2(\mathbf{C}^m)$  so that the action of  $U_2 / \{\text{center of } U_2\} \simeq SO_3$  has non discrete stabilizer. Those points are the lined configurations  ${}^m\tilde{\mathcal{P}}^1$ . Their values in  $\Xi_m$  are the non generic  $\alpha$ 's, which are the points in  $\Xi_m$  satisfying b) or c).  $\square$

We have proven most of the main result of this section: for generic and proper  $\alpha$ , the space  $\mathcal{P}^3(\alpha)$  is a Kähler sub-quotient of  $\mathbf{G}_2(\mathbf{C}^m)$ .

THEOREM 4.4. *For  $\alpha \in \text{int } \Xi_m$  generic,  $\mathcal{P}_+^3(\alpha)$  is a Kähler manifold isomorphic to the Kähler reduction  $U_1^m \backslash \mu^{-1}(\alpha)$ . The involution  $\smile$  is antiholomorphic and  $\mathcal{P}^2(\alpha)$  can be seen as the real part of  $\mathcal{P}_+^3(\alpha)$ .*

*Proof.* By 4.1, one has  $\mathcal{P}^3(\alpha) = \ell^{-1}(\alpha) = U_1^m \backslash \mu^{-1}(\alpha)$  and we have seen in 3.9 that  $\widehat{\Phi}(\bar{a}, \bar{b}) = \Phi(a, b)^\smile$ .  $\square$

We shall now compare the Kähler structure obtained on  $\mathcal{P}_+^3(\alpha)$  from the Grassmannian to that introduced by Klyachko [Kl] or Kapovich-Millson ([KM2], §3). Using the standard cross product  $\times$  and scalar product  $\langle \cdot, \cdot \rangle$  on  $\mathbf{R}^3$ , these authors put on the sphere  $S_r^2$  of radius  $r$  the complex structure  $\tilde{J}$

defined by

$$\tilde{J}v := \frac{1}{r} x \times v \quad (v \in T_x S_r^2)$$

and the Kähler metric

$$\tilde{h}(u, v) := \frac{1}{r} \langle u, v \rangle - \frac{i}{r^2} \langle x, u \times v \rangle \quad (u, v \in T_x S_r^2)$$

with associated symplectic form  $\tilde{\omega}(u, v) := \langle \frac{x}{r^2} u \times v \rangle$ . Let  $W(\alpha) := \prod_{i=1}^m S_{\alpha_i}^2$ . The map  $\beta : W_\alpha \rightarrow \mathbf{R}^3$  defined by  $\beta(z_1, \dots, z_m) := \sum_{i=1}^m z_i$  is the moment map for the diagonal action of  $SO_3$  on  $W_\alpha$ . The space  $\mathcal{P}_+^3(\alpha)$  thus occurs as the symplectic reduction  $SO_3 \backslash \beta^{-1}(0)$ .

**PROPOSITION 4.5.** *The complex structure  $J$  and Kähler metric  $h$  of 4.4 compare with those  $\tilde{J}$  and  $\tilde{h}$  of Kapovich-Millson in the following way:*

$$\tilde{J} = J \quad \text{and} \quad \tilde{h}(u, v) = 4h(u, v).$$

*Proof.* Starting from the Hermitian vector space  $\mathcal{M} = \mathcal{M}_{m \times 2}(\mathbf{C})$  one sees that  $\mathcal{P}^3(\alpha)$  is obtained by two successive symplectic reductions

$$\mathbf{G}_2(\mathbf{C}^m) = \tilde{\Phi}^{-1}(0)/U_2 \quad \text{and} \quad \mathcal{P}^3(\alpha) = U_1^m \backslash \mu^{-1}(\alpha)$$

(we use the notation of §3). One can perform the reductions in the reverse order. We first get

$$U_1^m \backslash \tilde{\Psi}^{-1}(\alpha) = \prod_{i=1}^m \mathbf{C}P_{\alpha_i}^1$$

where  $\mathbf{C}P_r^1$  is the quotient of the 3-dimensional sphere

$$\{(u, v) \in \mathbf{C}^2 \mid |u|^2 + |v|^2 = r\}$$

by the diagonal action of  $U_1$ . The moment map  $\tilde{\Phi} : \mathcal{M} \rightarrow \mathcal{H}(2)$  gives a moment map (still called  $\tilde{\Phi}$ ) from the product of projective spaces into  $\mathcal{H}_0(2)$ . One has a commutative diagram

$$\begin{array}{ccc} \prod_{i=1}^m \mathbf{C}P_{\alpha_i}^1 & \xrightarrow[\simeq]{\Pi^\phi} & \prod_{i=1}^m S_{\alpha_i}^2 \\ \tilde{\Phi} \downarrow & & \downarrow \beta \\ \mathcal{H}_0(2) & \xrightarrow[\simeq]{\psi} & \mathbf{R}^3 \end{array}$$

where  $\psi : \mathcal{H}_0(2) \rightarrow \mathbf{R}^3 \simeq \mathbf{R} \times \mathbf{C}$  sends the matrix  $\begin{pmatrix} u & z \\ \bar{z} & -u \end{pmatrix}$  to  $(u, z)$ .

To prove Proposition 4.5, it is enough to establish that for all  $a \in \mathbf{CP}_r^1$ , the tangent map  $T_a\phi : T_a\mathbf{CP}_r^1 \longrightarrow T_{\phi(a)}S_r^2$  satisfies

$$T_a\phi(Jv) = \tilde{J}T_a\phi(v) \quad \text{and} \quad \tilde{\omega}(T_a\phi(v), T_a\phi(Jv)) = 4\omega(v, Jv).$$

By  $U_2$ -equivariance, we can restrict ourselves to  $a = [\sqrt{r}, 0]$ . The tangent space  $T_a\mathbf{CP}_r^1$  is identified with  $\{0\} \times \mathbf{C}$  and one can take  $v = (0, 1)$  and  $Jv = (0, i)$ . One has  $\phi(a) = (r, 0, 0)$ ,

$$T_a\phi(v) = (0, 2\sqrt{r}, 0), \quad T_a\phi(Jv) = (0, 0, 2\sqrt{r}) = \tilde{J}T_a\phi(v)$$

and  $\tilde{\omega}(T_a\phi(v), T_a\phi(Jv)) = 4$ , while  $\omega(v, Jv) = 1$ .  $\square$

#### REMARKS

(4.6) The results of this section show that the spaces  $\mathcal{P}_+^3(\alpha)$  for generic  $\alpha$  are the symplectic leaves of the Poisson structure on the regular part of  ${}^m\mathcal{P}_+^3$ , or  ${}^m\mathcal{PP}_+^3$  given in (3.13) and (3.14).

(4.7) If one works in the pure quaternions  $I\mathbf{H}$ , the complex structure  $\tilde{J}$  on  $S_r^2$  becomes

$$\tilde{J}(v) = \frac{qv}{|q|}, \quad (v \in T_q S_r^2 = I\mathbf{H}).$$

The sphere  $S_r^2$  is a co-adjoint orbit of  $U_1(\mathbf{H})$  and the Hermitian form  $\tilde{\omega}$  is the Kirillov–Kostant form (see [Gu, Theorem 1.1]).

(4.8) The isomorphism between the symplectic reductions of the Grassmannian  $\mathbf{G}_2(\mathbf{C}^m)$  and the product of  $\mathbf{CP}^1$ 's that underlies our results 3.9, 4.4 and the proof of 4.5 is a symplectic version of the Gel'fand-MacPherson correspondence ([GM] and [GGMS]). The fact that this isomorphism comes from two reductions of  $\mathcal{M}$  is the philosophy of “dual pairs” (see [Mo] and the references therein).

## 5. THE GEL'FAND-CETLIN ACTION

On  ${}^m\mathcal{F}^k$  we have so far defined the length functions  $\tilde{\ell}$  measuring the distances between successive vertices. We now introduce  $\tilde{d} : {}^m\mathcal{F}^k \rightarrow \mathbf{R}^m$ ,  $\tilde{d}(\rho) = (|\rho(1)|, |\rho(1) + \rho(2)|, \dots, |\sum_{i=1}^m \rho(i)|)$ , the lengths of the diagonals connecting the vertices to the origin. (Only  $m-3$  of these functions are new, as  $\tilde{d}(\rho)_1 = \tilde{\ell}(\rho)_1$ ,  $\tilde{d}(\rho)_{m-1} = \tilde{\ell}(\rho)_m$ , and  $\tilde{d}(\rho)_m = 0$ . Hereafter we write only  $\ell_i, d_i$  and the  $\rho$  is to be understood.)