

3.2 Solutions of the Lagrange top

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

and hence $\Psi = \tilde{\Psi}$. Finally, the reader may check that the functions (46) and (47) have the analyticity properties from Proposition 3.2 and hence they coincide with the Baker-Akhiezer function defined in Proposition 3.1. \square

3.2 SOLUTIONS OF THE LAGRANGE TOP

Let $z = (z_1, z_2) \in J(C_h; \infty^\pm)$. It is easy to check that the functions

$$\frac{\theta_{11}(z_1 \pm \tau_2)}{\theta_{11}(z_1)} e^{\mp z_2}$$

live on $J(C_h; \infty^\pm)$. We shall see that they give solutions of the Lagrange top. By (16) we compute that $\frac{d}{dt}z = \text{constant}$, where

$$\begin{aligned} \frac{dz}{dt} &= \begin{pmatrix} V_1 \\ V_2 \end{pmatrix} = 2\pi i \begin{pmatrix} \int_{A_1} \frac{d\lambda}{\mu} & \int_{A_2} \frac{d\lambda}{\mu} \\ \int_{A_1} \frac{\lambda d\lambda}{\mu} & \int_{A_2} \frac{\lambda d\lambda}{\mu} \end{pmatrix}^{-1} \begin{pmatrix} -i \\ -ai \end{pmatrix}, \\ &\quad \int_{A_2} \frac{d\lambda}{\mu} = 0, \quad \int_{A_2} \frac{\lambda d\lambda}{\mu} = -2\pi i \end{aligned}$$

so

$$\begin{pmatrix} V_1 \\ V_2 \end{pmatrix} = \left(\int_{A_1} \frac{d\lambda}{\mu} \right)^{-1} \begin{pmatrix} 2\pi \\ -i \int_{A_1} \frac{\lambda d\lambda}{\mu} + ai \int_{A_1} \frac{d\lambda}{\mu} \end{pmatrix}, \quad a = -m\Omega_3.$$

THEOREM 3.4. *The following equations hold*

$$(48) \quad \bar{\epsilon}\Omega_1(t) + \epsilon\Omega_2(t) = \text{const}_3 \frac{\theta_{11}(z_1 - \tau_2)}{\theta_{11}(z_1)} e^{-z_2},$$

$$(49) \quad \epsilon\Omega_1(t) + \bar{\epsilon}\Omega_2(t) = \text{const}_4 \frac{\theta_{11}(z_1 + \tau_2)}{\theta_{11}(z_1)} e^{+z_2},$$

where

$$(50) \quad \begin{aligned} z_2 &= tV_2, \quad z_1 = tV_1 + \mathcal{A}(\infty^+ + \infty^- - P_1 - P_2), \\ \tau_2 &= \mathcal{A}(\infty^+ - \infty^-) = \int_{B_1} \omega_2 \end{aligned}$$

and

$$\text{const}_3 = \frac{2iV_1\theta'_{11}(0)}{\theta_{11}(\mathcal{A}(\infty^- - \infty^+))} \cdot \frac{\theta_{11}(\mathcal{A}(\infty^+ - P_1))}{\theta_{11}(\mathcal{A}(\infty^- - P_1))} \cdot \frac{\theta_{11}(\mathcal{A}(\infty^+ - P_2))}{\theta_{11}(\mathcal{A}(\infty^- - P_2))},$$

$$\text{const}_4 = \frac{2iV_1\theta'_{11}(0)}{\theta_{11}(\mathcal{A}(\infty^+ - \infty^-))} \cdot \frac{\theta_{11}(\mathcal{A}(\infty^- - P_1))}{\theta_{11}(\mathcal{A}(\infty^+ - P_1))} \cdot \frac{\theta_{11}(\mathcal{A}(\infty^- - P_2))}{\theta_{11}(\mathcal{A}(\infty^+ - P_2))}.$$

Let us denote

$$\begin{aligned}\omega_1 &= \pm(\omega_1^0 + O(\lambda^{-1})) d(\lambda^{-1}), & P = (\lambda, \mu) \rightarrow \infty^\pm, \\ \omega_2 &= \pm(\omega_2^1 \lambda + \omega_2^0 + O(\lambda^{-1})) d(\lambda^{-1}), & P = (\lambda, \mu) \rightarrow \infty^\pm.\end{aligned}$$

To prove Theorem 3.4 we shall need the following

LEMMA 3.5. *The above defined differentials are such that*

$$\begin{aligned}\omega_1^0 &= -i \int_{B_1} \Omega = -iV_1, & \omega_2^0 = i(c^+ - c^-), \\ V_2 &= -c^+ + c^- + i\Omega_3, & \mathcal{A}(\infty^+ - \infty^-) = \int_{B_1} \omega_2.\end{aligned}$$

Proof. The identity $\omega_1^0 = -i \int_{B_1} \Omega$ is a reciprocity law between the differential of the first kind ω_1 and the differential of the second kind Ω [13]. It is obtained by integrating $\pi(P)\omega_1$, where $\pi(P) = \int_{P_0}^P \Omega$, along the border of C_h cut along its homology basis A_1, B_1 . On the other hand

$$\omega_1 = 2\pi i \left(\int_{A_1} \frac{d\lambda}{\mu} \right)^{-1} \frac{d\lambda}{\mu}$$

and hence

$$\omega_1^0 = -2\pi i \left(\int_{A_1} \frac{d\lambda}{\mu} \right)^{-1} = -iV_1.$$

Similarly the identity $\omega_2^0 = i(c^+ - c^-)$ is a reciprocity law between the differential of the third kind ω_2 and the differential of the second kind Ω , and $\mathcal{A}(\infty^+ - \infty^-) = \int_{B_1} \omega_2$ is a reciprocity law between the differential of the third kind ω_2 and the differential of the first kind ω_1 . Finally, as

$$\omega_2 = \frac{\int_{A_1} \frac{\lambda d\lambda}{\mu}}{\int_{A_1} \frac{d\lambda}{\mu}} \frac{d\lambda}{\mu} - \frac{\lambda d\lambda}{\mu} \text{ we have } \omega_2^0 = -\frac{\int_{A_1} \frac{\lambda d\lambda}{\mu}}{\int_{A_1} \frac{d\lambda}{\mu}} - (1+m)\Omega_3 = -iV_1 - \Omega_3$$

and hence $V_2 = -c^+ + c^- + i\Omega_3$. \square

Proof of Theorem 3.4. According to (42), (43)

$$\bar{\epsilon} \Omega_1(t) + \epsilon \Omega_2(t) = -2 \lim_{P \rightarrow \infty^-} \frac{\lambda \Psi^1(t, P)}{\Psi^2(t, P)}$$

and

$$\epsilon \Omega_1(t) + \bar{\epsilon} \Omega_2(t) = +2 \lim_{P \rightarrow \infty^+} \frac{\lambda \Psi^2(t, P)}{\Psi^1(t, P)}.$$

To compute the limit we use (46), (47) and

$$\lim_{P \rightarrow \infty^-} \lambda(P) \theta_{11}(\mathcal{A}(P - \infty^-)) = \theta'_{11}(0) \frac{d}{ds} \Big|_{s=0} \int^s \omega_1 = \omega_1^0 \theta'_{11}(0)$$

$$\lim_{P \rightarrow \infty^+} \lambda(P) \theta_{11}(\mathcal{A}(P - \infty^+)) = \theta'_{11}(0) \frac{d}{ds} \Big|_{s=0} \int^s \omega_1 = \omega_1^0 \theta'_{11}(0)$$

(see Lemma 3.5). \square

3.3 EFFECTIVIZATION

Let \wp, ζ, σ be the Weierstrass functions related to the elliptic curve Γ defined by

$$(51) \quad \eta^2 = 4\xi^3 - g_2\xi - g_3$$

(we use the standard notations of [4]).

Consider also the *real* elliptic curve C with affine equation

$$(52) \quad \mu^2 + \lambda^4 + a_1\lambda^3 + a_2\lambda^2 + a_3\lambda + a_4 = 0$$

and natural anti-holomorphic involution $(\lambda, \mu) \rightarrow (\bar{\lambda}, \bar{\mu})$, and put

$$(53) \quad g_2 = a_4 + 3\left(\frac{a_2}{6}\right)^4 - 4\frac{a_1}{4}\frac{a_3}{4}, \quad g_3 = \det \begin{pmatrix} 1 & \frac{a_1}{4} & \frac{a_2}{6} \\ \frac{a_1}{4} & \frac{a_2}{6} & \frac{a_3}{4} \\ \frac{a_2}{6} & \frac{a_3}{4} & a_4 \end{pmatrix}.$$

It is well known that the curves C and Γ are isomorphic over \mathbf{C} and that under this isomorphism

$$(54) \quad \frac{d\lambda}{\mu} = \frac{d\xi}{\eta}.$$

Following Weil [25] we call Γ the Jacobian $J(C)$ of the elliptic curve C and we write $J(C) = \Gamma$. Note that $J(C)$ and Γ are real isomorphic and that $J(C)$ and C are not real isomorphic.

Further we make the substitution (23) and C becomes the spectral curve \tilde{C}_h of Adler and van Moerbeke $\{\mu^2 + f(\lambda) = 0\}$, where

$$f(\lambda) = \lambda^4 + 2(1+m)h_4\lambda^3 + (2h_3 + m(m+1)h_4^2)\lambda^2 - 2h_2\lambda + 1$$

and Γ becomes the Lagrange curve Γ_h . Recall that, as we explained at the end of Section 2, the curve C_h with an equation $\{\mu^2 = f(\lambda)\}$ and antiholomorphic involution $(\lambda, \mu) \rightarrow (\bar{\lambda}, -\bar{\mu})$, is isomorphic over \mathbf{R} to \tilde{C}_h , so we write $C_h = \tilde{C}_h$. The Jacobian curve $J(C_h) = \Gamma_h$ was computed by