

10. Direct products of graphs

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Proof. This follows from

$$\begin{aligned} 1/\rho(G_{\mathcal{X}}) &= \lim_{t \rightarrow \infty} \frac{1}{(tG_{\mathcal{X}})^{-1}} \\ &= \lim_{t \rightarrow \infty} \frac{1}{(tG_{\mathcal{E}})^{-1}} + \frac{1}{(tG_{\mathcal{F}})^{-1}} - \frac{1}{t} \quad \text{by (9.2)} \\ &= 1/\rho(G_{\mathcal{E}}) + 1/\rho(G_{\mathcal{F}}) - 0. \quad \square \end{aligned}$$

Note that the corollary does not extend to non-recurrent series; for instance, it fails if $\mathcal{E} = \mathcal{F} = \mathbf{Z}$. Indeed then

$$\begin{aligned} G_{\mathcal{E}} = G_{\mathcal{F}} &= \frac{1}{\sqrt{1-4t^2}}, & \rho(G_{\mathcal{E}}) = \rho(G_{\mathcal{F}}) &= 1/4, \\ G_{\mathcal{X}} &= \frac{3}{1+2\sqrt{1-12t^2}}, & \rho(G_{\mathcal{X}}) &= 1/\sqrt{12}. \end{aligned}$$

10. DIRECT PRODUCTS OF GRAPHS

There are two natural definitions for *direct products* of graphs; they correspond to direct products of groups with generating set either the union or cartesian product of the generating sets of the factors. A general treatment of products of graphs can be found in [CDS79, pages 65 and 203].

DEFINITION 10.1. If S is a set, the *stationing graph* on S is the graph $\mathcal{X} = \Sigma_S$ with $V(\mathcal{X}) = E(\mathcal{X}) = S$, where for the edges $s^\alpha = s^\omega = \bar{s} = s$ hold.

LEMMA 10.2. Let \mathcal{X} be a graph, and $\mathcal{E} = \mathcal{X} \sqcup \Sigma_{\mathcal{X}}$ be the graph obtained by adding a loop to every vertex in \mathcal{X} . Let $G_{\mathcal{X}}$ and $G_{\mathcal{E}}$ be the growth functions for circuits in \mathcal{X} and \mathcal{E} respectively. Then we have

$$G_{\mathcal{E}}(t) = \frac{1}{1-t} G_{\mathcal{X}}\left(\frac{t}{1-t}\right).$$

DEFINITION 10.3 (First Product). Let \mathcal{E} and \mathcal{F} be two graphs. Their *direct product* $\mathcal{X} = \mathcal{E} \times \mathcal{F}$ is defined by

$$V(\mathcal{X}) = V(\mathcal{E}) \times V(\mathcal{F})$$

and

$$E(\mathcal{X}) = (E(\mathcal{E}) \times \Sigma_{\mathcal{F}}) \sqcup (\Sigma_{\mathcal{E}} \times E(\mathcal{F})).$$

Note that if the graphs \mathcal{E} and \mathcal{F} have respectively adjacency matrices E and F , then their product has adjacency matrix $E \otimes \mathbf{1} + \mathbf{1} \otimes F$.

In that case we have

$$G_{\mathcal{X}} = \frac{1}{2i\pi} \oint_{S^1} \frac{G_{\mathcal{E}}((1+u)t) G_{\mathcal{F}}((1+u^{-1})t)}{u} du.$$

This is a simple application of the Laplace transform, that converts an exponential generating function into an ordinary one and vice versa [AS70, 29.3.3]. Indeed, if we had considered exponential generating functions, the formula would simply have been $G_{\mathcal{X}} = G_{\mathcal{E}}G_{\mathcal{F}}$, as is well known (see [Wil90] or [Sta78, page 102]).

As an example, let $\mathcal{E} = \mathcal{F} = \mathbf{Z}$, so $G_{\mathcal{E}} = G_{\mathcal{F}} = \frac{1}{\sqrt{1-t^2}}$. Then

$$\begin{aligned} G_{\mathcal{X}} &= \frac{1}{2i\pi} \oint_{S^1} \frac{du}{\sqrt{(1-4(1+u)^2t^2)(u^2-4(1+u)^2t^2)}} \\ &= \frac{2}{\pi} K(16t^2) = F\left(\begin{matrix} 1/2 & & 1/2 \\ & 1 & \end{matrix} \middle| 16t^2\right) \end{aligned}$$

where K is the complete elliptic function and F the hypergeometric series. These functions are known to be transcendental; thus the circuit series of \mathbf{Z}^2 is transcendental. This result appears in [GH97]. Numerical evidence suggests the growth function for \mathbf{Z}^3 is not even hypergeometric.

DEFINITION 10.4 (Second Product). Let \mathcal{E} and \mathcal{F} be two graphs, and suppose that for every vertex in \mathcal{E} and \mathcal{F} there is a loop at it. Then their *direct product* $\mathcal{X} = \mathcal{E} \times \mathcal{F}$ is defined by

$$V(\mathcal{X}) = V(\mathcal{E}) \times V(\mathcal{F})$$

and

$$E(\mathcal{X}) = E(\mathcal{E}) \times E(\mathcal{F}).$$

Note that if the graphs \mathcal{E} and \mathcal{F} have respectively adjacency matrices E and F , then their product has adjacency matrix $E \otimes F$.

In that case we have, again using Laplace transformations

$$G_{\mathcal{X}}(t) = \frac{1}{2i\pi} \oint_{S^1} \frac{G_{\mathcal{E}}(u)G_{\mathcal{F}}(t/u)}{u} du.$$

Note that with both definitions of products it is possible that the growth function for circuits in the product be transcendental even if the growth functions for circuits in the factors are algebraic.