

Proof of the main result

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Though we cannot solve the above problem, the purpose of this note is to show that, if one replaces \mathbf{R}^2 by \mathbf{Z}^2 , the new problem has a positive solution. In fact, we will prove the following result, which shows somewhat more, namely that the action on \mathbf{Z}^2 may be an action without non-trivial fixed points, rather than only locally commutative. We denote by $\text{SA}_2(\mathbf{Z})$ the group of all transformations $\vec{x} \mapsto A\vec{x} + \vec{a}$ of \mathbf{Z}^2 , with $A \in \text{SL}_2(\mathbf{Z})$ and $\vec{a} \in \mathbf{Z}^2$.

THEOREM. *The group $\text{SA}_2(\mathbf{Z})$ has a free subgroup F_2 of rank 2 which acts on \mathbf{Z}^2 without non-trivial fixed points, namely the subgroup generated by*

$$\zeta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

and

$$\eta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

The theorem implies the existence of a partition of \mathbf{Z}^2 into three pieces P , Q and R such that the six pieces P , Q , R , $P \sqcup Q$, $Q \sqcup R$, $R \sqcup P$ are pairwise F_2 -congruent, without the axiom of choice ([S0], and Corollary 4.12 in [W]).

As observed in the discussion which follows Proposition 7.1 in [W], it is known that the above theorem does not carry over to \mathbf{R}^2 ; more precisely, it is known that a subgroup of $\text{SA}_2(\mathbf{R})$ which acts on \mathbf{R}^2 without non-trivial fixed points is soluble, and consequently does not contain non-commutative free subgroups.

PROOF OF THE MAIN RESULT

Recall that a matrix in $\text{SL}_2(\mathbf{Z})$ is *hyperbolic* if the absolute value of its trace is strictly larger than 2, or equivalently if it has an eigenvalue of absolute value strictly larger than 1.

LEMMA 0. *The subgroup of $\text{SL}_2(\mathbf{Z})$ generated by*

$$\begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \text{ and } \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix}$$

is free of rank 2 and all its elements distinct from the identity are hyperbolic.

Proof. It is well-known that the subgroup of $\mathrm{SL}_2(\mathbf{Z})$ generated by

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$$

is free of rank 2, and that all its elements distinct from the identity are hyperbolic. (See Appendix B in [K], [Ma], [MW], [N], or the proof of Theorem 6.8 in [W].) The lemma follows, because

$$\begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^2 \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}^2$$

(see for example Exercice 12 of Section 1.4 in [MKS]). \square

The following is elementary linear algebra.

LEMMA 1. *For $A \in \mathrm{SL}_2(\mathbf{R})$ with $\mathrm{tr} A \neq 2$ and for $\vec{a} \in \mathbf{R}^2$, the affine transformation*

$$\begin{cases} \mathbf{R}^2 & \rightarrow \mathbf{R}^2 \\ \vec{x} & \mapsto A\vec{x} + \vec{a} \end{cases}$$

has a unique fixed point.

Our preparations are complete.

Proof of the main theorem. The two transformations ζ and η of our main result generate a group which is free of rank 2, by Lemma 0. As both these transformations fix the point

$$\begin{pmatrix} 2/3 \\ -5/3 \end{pmatrix} \in \mathbf{R}^2,$$

each element of the group they generate fix the same point. As this point is not in \mathbf{Z}^2 , the theorem follows by Lemma 1. \square

REMARK. Let $\alpha, \beta \in \mathrm{SA}_2(\mathbf{Z})$ be as in the introduction. Then we can prove the main theorem by using the group generated by $\alpha\beta^{-1}\alpha\beta^{-2}\alpha$ and $\beta\alpha^{-1}\beta\alpha^{-2}\beta$, because the transformations

$$\alpha\beta^{-1}\alpha\beta^{-2}\alpha : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -17 \\ -13 \end{pmatrix}$$

and

$$\beta\alpha^{-1}\beta\alpha^{-2}\beta : \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -13 \\ -17 \end{pmatrix}$$

have a common fixed point $\begin{pmatrix} 1/2 \\ 1/2 \end{pmatrix}$ in \mathbf{R}^2 and the subgroup of $\mathrm{SL}_2(\mathbf{Z})$ generated by

$$\begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix}$$

is free of rank 2 and all its elements distinct from the identity are hyperbolic. See [K] and the following calculations:

$$\begin{pmatrix} 13 & 22 \\ 10 & 17 \end{pmatrix} = tu(tu^{-1})^3(tu)^2tu^{-1}tu,$$

$$\begin{pmatrix} 17 & 10 \\ 22 & 13 \end{pmatrix} = tu^{-1}(tu)^3(tu^{-1})^2tutu^{-1},$$

where

$$t = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad u = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$$

with $\langle t, u \rangle / \{\pm 1\} \cong \mathbf{Z}_2 * \mathbf{Z}_3$.

The referee suggested to the author that the following could be shown (without the axiom of choice).

COROLLARY. *There exists a subset E_1 of \mathbf{Z}^2 such that, for every finite subset F of \mathbf{Z}^2 , the symmetric difference of E_1 and F is congruent to E_1 relative to the group $\mathrm{SA}_2(\mathbf{Z})$.*

Proof. This is a consequence of our main result and of Theorem 2 in [My] ($S = \mathbf{Z}^2$, $G = \langle \zeta, \eta \rangle$, $M = \{\zeta\eta, \zeta^2\eta^2, \zeta^3\eta^3, \dots\}$, $\mathbf{F} = \{F \subseteq \mathbf{Z}^2 \mid F \text{ is finite}\}$). \square

REFERENCES

- [BT] BANACH, S. and A. TARSKI. Sur la décomposition des ensembles de points en parties respectivement congruentes. *Fund. Math.* 6 (1924), 244–277.
- [B] BOREL, A. On free subgroups of semi-simple groups. *L'Enseignement Math.* 29 (1983), 151–164.
- [D] DEKKER, Th. J. Decompositions of sets and spaces I. *Indag. Math.* 18 (1956), 581–589.
- [DS] DELIGNE, P. and D. SULLIVAN. Division algebras and the Hausdorff-Banach-Tarski paradox. *L'Enseignement Math.* 29 (1983), 145–150.
- [K] KUROSH, A. G. *The Theory of Groups*, vol. 2. Chelsea Publishing Company, New York, 1956.