Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: p-ADIC L-FUNCTION OF TWO VARIABLES
Autor: Fox, Glenn J.
Kurzfassung
DOI: https://doi.org/10.5169/seals-64800

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.06.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

L'Enseignement Mathématique, t. 46 (2000), p. 225–278

A p-ADIC L-FUNCTION OF TWO VARIABLES

by Glenn J. FOX*)

ABSTRACT. For p prime and χ a primitive Dirichlet character, we derive a p-adic function $L_p(s,t;\chi)$, where $t \in \mathbf{C}_p$, $|t|_p \leq 1$, and $s \in \mathbf{C}_p$, $|s-1|_p < |p|_p^{1/(p-1)}|q|_p^{-1}$, $s \neq 1$ if $\chi = 1$, with $q = 4$ if $p = 2$ and $q = p$ if $p > 2$, that interpolates the values

$$L_p(1-n,t;\chi) = -\frac{1}{n}\Big(B_{n,\chi_n}(qt) - \chi_n(p)p^{n-1}B_{n,\chi_n}(p^{-1}qt)\Big),$$

for $n \in \mathbf{Z}$, $n \geq 1$. Here $B_{n,\chi}(t)$ is the nth generalized Bernoulli polynomial associated with the character χ, and $\chi_n = \chi\omega^{-n}$, where ω is the Teichmüller character. This function is then a two-variable analogue of the p-adic L-function $L_p(s;\chi)$, where $s \in \mathbf{C}_p$, $|s-1|_p < |p|_p^{1/(p-1)}|q|_p^{-1}$, $s \neq 1$ if $\chi = 1$, in that this function satisfies $L_p(s,0;\chi) = L_p(s;\chi)$. In addition to deriving this function, we establish several properties and applications of $L_p(s,t;\chi)$.

1. INTRODUCTION

Given a primitive Dirichlet character χ, having conductor f_χ (see Section 2 for definitions), the Dirichlet L-function associated with χ is defined by

$$L(s;\chi) = \sum_{b=1}^{\infty} \frac{\chi(b)}{b^s},$$

where $s \in \mathbf{C}$, $\Re(s) > 1$. This function can be continued analytically to the entire complex plane, except for a simple pole at $s = 1$ when $\chi = 1$, in which case we have the Riemann zeta function, $\zeta(s) = L(s;1)$. It is believed that the analysis of Dirichlet L-functions began with Euler's study of $\zeta(s)$, in which he considered the function only for real values of s. It was Riemann

*) A majority of these results were obtained while the author was a graduate student at the University of Georgia, Athens, under the direction of Andrew Granville.