2. Preliminaries

Objekttyp: Chapter

Zeitschrift: L'Enseignement Mathématique

Band (Jahr): 46 (2000)

Heft 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

PDF erstellt am: 25.05.2024

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek* ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

http://www.e-periodica.ch

THEOREM 4.12. Let n, c, and k be positive integers, and let $\tau \in \mathbb{Z}_p$ such that $|\tau|_p \leq |pq^{-1}F_0|_p$. Then the quantity

$$\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau) - \binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0) \in \mathbf{Z}_p[\chi],$$

and, modulo $q\mathbf{Z}_p[\chi]$, is independent of n.

These results show that if related congruences hold for

$$\beta_{n,\chi}(0) = -\frac{1}{n} (1 - \chi_n(p)p^{n-1})B_{n,\chi_n},$$

then they must also hold for $\beta_{n,\chi}(\tau)$, where τ is any element of \mathbb{Z}_p such that $|\tau|_p \leq |pq^{-1}F_0|_p$.

In [9] Granville defined ordinary Bernoulli numbers of negative index, B_{-n} , where $n \in \mathbb{Z}$, $n \ge 1$, in the field \mathbb{Q}_p according to

$$B_{-n} = \lim_{k \to \infty} B_{\phi(p^k) - n} \,,$$

where the limit is taken in the *p*-adic sense. In a similar manner we define generalized Bernoulli numbers of negative index, $B_{-n,\chi}$, $n \in \mathbb{Z}$, $n \ge 1$, and a collection of functions that correspond to generalized Bernoulli polynomials of negative index, $B_{-n,\chi}(t)$, $n \in \mathbb{Z}$, $n \ge 1$. As a result of our definitions, we show that the $B_{-n,\chi}(t)$ are actually power series that can be written in the form

$$B_{-n,\chi}(t) = \sum_{m=0}^{\infty} {\binom{-n}{m}} B_{-n-m,\chi} t^m,$$

converging for $t \in \mathbf{C}_p$, $|t|_p < 1$. We close out by considering some properties of these functions.

2. PRELIMINARIES

The *p*-adic *L*-functions, $L_p(s; \chi)$, were first generated by Kubota and Leopoldt for the purpose of finding functions that would serve as analogues of the Dirichlet *L*-functions in the *p*-adic number field [14]. They are characterized by the fact that they interpolate a specific expression involving generalized Bernoulli numbers when the variable *s* is a nonpositive integer. In the following, for each $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, we derive a *p*-adic function $L_p(s,\tau;\chi)$ that interpolates a specific expression involving generalized Bernoulli polynomials in τ for similar values of the variable *s*. These functions are designed so that $L_p(s, 0; \chi) = L_p(s; \chi)$. The method of derivation follows that found in [13], Chapter 3. However, this method will only account for those $\tau \in \overline{\mathbf{Q}}_p$ with $|\tau|_p \leq 1$. To complete the derivation we show that there exist functions $L_p(s, \tau; \chi)$ for all $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, such that for every sequence $\{\tau_i\}_{i=0}^{\infty}$ in $\overline{\mathbf{Q}}_p$, with $|\tau_i|_p \leq 1$, converging to some $\tau \in \mathbf{C}_p$, the sequence $\{L_p(1-n,\tau_i;\chi)\}_{i=0}^{\infty}$, with $n \in \mathbf{Z}$, $n \geq 1$, converges to $L_p(1-n,\tau;\chi)$. Thus for each $\tau \in \mathbf{C}_p$, $|\tau|_p \leq 1$, the function $L_p(s,\tau;\chi)$ must interpolate the appropriate expressions involving generalized Bernoulli polynomials for s = 1 - n, $n \in \mathbf{Z}$, $n \geq 1$.

Before we begin the derivation, we must first define the concepts that we shall need and review some of their resulting properties.

2.1 DIRICHLET CHARACTERS

For $n \in \mathbb{Z}$, $n \ge 1$, a Dirichlet character to the modulus n is a multiplicative map $\chi : \mathbb{Z} \to \mathbb{C}$ such that $\chi(a+n) = \chi(a)$ for all $a \in \mathbb{Z}$, and $\chi(a) = 0$ if and only if $(a, n) \ne 1$. Since $a^{\phi(n)} \equiv 1 \pmod{n}$ for all a such that (a, n) = 1, $\chi(a)$ must be a root of unity for such a.

If χ is a Dirichlet character to the modulus n, then for any positive multiple m of n we can induce a Dirichlet character ψ to the modulus m according to

$$\psi(a) = \begin{cases} \chi(a), & \text{if } (a,m) = 1\\ 0, & \text{if } (a,m) \neq 1. \end{cases}$$

The minimum modulus n for which a character χ cannot be induced from some character to the modulus m, m < n, is called the conductor of χ , denoted f_{χ} . We shall assume that each χ is defined modulo its conductor. Such a character is said to be primitive.

For primitive Dirichlet characters χ and ψ having conductors f_{χ} and f_{ψ} , respectively, we define the product, $\chi\psi$, to be the primitive character with $\chi\psi(a) = \chi(a)\psi(a)$ for all $a \in \mathbb{Z}$ such that $(a, f_{\chi}f_{\psi}) = 1$. Note that there may exist some values of a such that $\chi\psi(a) \neq \chi(a)\psi(a)$, due to the fact that our definition requires $\chi\psi$ to be a primitive character. The conductor $f_{\chi\psi}$ then divides $\operatorname{lcm}(f_{\chi}, f_{\psi})$. With this operation defined, we can then consider the set of primitive Dirichlet characters to form a group under multiplication. The identity of the group is the principal character $\chi = 1$, having conductor $f_1 = 1$. The inverse of the character χ is the character $\chi^{-1} = \overline{\chi}$, the map of complex conjugates of the values of χ .

Since any Dirichlet character χ is multiplicative, we must have $\chi(-1) = \pm 1$. A character χ is said to be odd if $\chi(-1) = -1$, and even if $\chi(-1) = 1$.

2.2 GENERALIZED BERNOULLI POLYNOMIALS

Let χ be a Dirichlet character with conductor f_{χ} . Then we define the functions, $B_{n,\chi}(t)$, $n \in \mathbb{Z}$, $n \ge 0$, by the generating function

(1)
$$\sum_{a=1}^{f_{\chi}} \frac{\chi(a) x e^{(a+t)x}}{e^{f_{\chi}x} - 1} = \sum_{n=0}^{\infty} B_{n,\chi}(t) \frac{x^n}{n!}, \quad |x| < \frac{2\pi}{f_{\chi}}$$

We define the generalized Bernoulli numbers associated with χ , $B_{n,\chi}$, $n \in \mathbb{Z}$, $n \ge 0$, by

$$\sum_{a=1}^{f_{\chi}} \frac{\chi(a) x e^{ax}}{e^{f_{\chi}x} - 1} = \sum_{n=0}^{\infty} B_{n,\chi} \frac{x^n}{n!}, \quad |x| < \frac{2\pi}{f_{\chi}},$$

so that $B_{n,\chi}(0) = B_{n,\chi}$. Note that

$$\sum_{a=1}^{f_{\chi}} \frac{\chi(a) x e^{(a+t)x}}{e^{f_{\chi}x} - 1} = e^{tx} \sum_{a=1}^{f_{\chi}} \frac{\chi(a) x e^{ax}}{e^{f_{\chi}x} - 1},$$

which implies that

$$\sum_{n=0}^{\infty} B_{n,\chi}(t) \frac{x^n}{n!} = e^{tx} \sum_{n=0}^{\infty} B_{n,\chi} \frac{x^n}{n!},$$

and from this we obtain

(2)
$$B_{n,\chi}(t) = \sum_{m=0}^{n} \binom{n}{m} B_{n-m,\chi} t^{m}.$$

Thus the functions $B_{n,\chi}(t)$, defined in (1), are actually polynomials, called the generalized Bernoulli polynomials associated with χ . Let $\mathbb{Z}[\chi]$ denote the ring generated over \mathbb{Z} by all the values $\chi(a)$, $a \in \mathbb{Z}$, and $\mathbb{Q}(\chi)$ the field generated over \mathbb{Q} by all such values. Then it can be shown that $f_{\chi}B_{n,\chi}$ must be in $\mathbb{Z}[\chi]$ for each $n \ge 0$ whenever $\chi \ne 1$. In general, we have $B_{n,\chi} \in \mathbb{Q}(\chi)$ for each $n \ge 0$, and so $B_{n,\chi}(t) \in \mathbb{Q}(\chi)[t]$. The polynomials $B_{n,\chi}(t)$ exhibit the property that, for all $n \ge 0$,

(3)
$$B_{n,\chi}(-t) = (-1)^n \chi(-1) B_{n,\chi}(t) ,$$

whenever $\chi \neq 1$. Thus $B_{n,\chi}(t)$, for $\chi \neq 1$, is either an even function or an odd function according to whether $(-1)^n \chi(-1)$ is 1 or -1. From (3) we obtain

$$B_{n,\chi}=(-1)^n\chi(-1)B_{n,\chi}\,,$$

and so $B_{n,\chi} = 0$ whenever *n* is even and χ is odd, or whenever *n* is odd and χ is even, $\chi \neq 1$. Another property that the polynomials satisfy is that for $m \in \mathbb{Z}$, $m \geq 1$,

(4)
$$B_{n,\chi}(mf_{\chi}+t) - B_{n,\chi}(t) = n \sum_{a=1}^{mf_{\chi}} \chi(a)(a+t)^{n-1},$$

for all $n \ge 0$. This can be derived from (1). Note that for $\chi = 1$ and t = 0 this becomes

$$\frac{1}{n} \left(B_{n,1}(m) - B_{n,1} \right) = \sum_{a=1}^{m} a^{n-1}.$$

If $\chi \neq 1$, then it can be shown that $\sum_{a=1}^{f_{\chi}} \chi(a) = 0$, and from the above relations we can derive

$$B_{0,\chi} = \frac{1}{f_{\chi}} \sum_{a=1}^{J_{\chi}} \chi(a)$$

for all χ . Therefore

$$B_{0,\chi} = \begin{cases} 0, & \text{if } \chi \neq 1 \\ 1, & \text{if } \chi = 1 \end{cases}$$

The ordinary Bernoulli polynomials, $B_n(t)$, $n \in \mathbb{Z}$, $n \ge 0$, are defined by

(5)
$$\frac{xe^{tx}}{e^{x}-1} = \sum_{n=0}^{\infty} B_{n}(t) \frac{x^{n}}{n!}, \quad |x| < 2\pi,$$

and the Bernoulli numbers, B_n , $n \in \mathbb{Z}$, $n \ge 0$,

$$\frac{x}{e^{x}-1} = \sum_{n=0}^{\infty} B_{n} \frac{x^{n}}{n!}, \quad |x| < 2\pi.$$

From this we obtain the values $B_0 = 1$, $B_1 = -1/2$, $B_2 = 1/6$, $B_4 = -1/30, \ldots$, with $B_n = 0$ for odd $n \ge 3$. For even $n \ge 2$, we have

$$B_n = -\frac{1}{n+1} \sum_{m=0}^{n-1} {\binom{n+1}{m}} B_m.$$

Note that we again have the relations $B_n(0) = B_n$ and

$$B_n(t) = \sum_{m=0}^n \binom{n}{m} B_{n-m} t^m,$$

as we did for the generalized Bernoulli polynomials.

Some of the more important properties of Bernoulli polynomials are that

(6)
$$B_n(t+1) - B_n(t) = nt^{n-1}$$

for all $n \ge 1$, and

$$B_n(1-t) = (-1)^n B_n(t)$$

for $n \ge 0$. Each of these results can be derived from the generating function (5) above.

Similar to (4) for the generalized Bernoulli polynomials, whenever $m, n \in \mathbb{Z}, m \ge 1, n \ge 1$,

$$\frac{1}{n}(B_n(m) - B_n) = \sum_{a=0}^{m-1} a^{n-1},$$

where we take 0^0 to be 1 in the case of a = 0 and n = 1. Note that this can be derived from (6) since

$$B_n(m) - B_n = \sum_{a=0}^{m-1} (B_n(a+1) - B_n(a))$$

The Bernoulli numbers are rational numbers, and, in fact, the von Staudt-Clausen theorem states that for even $n \ge 2$,

$$B_n + \sum_{\substack{p \text{ prime} \ (p-1)|n}} \frac{1}{p} \in \mathbb{Z}.$$

Thus the denominator of each B_n must be square-free.

The ordinary Bernoulli numbers are related to the generalized Bernoulli numbers in that for $\chi = 1$ we have

$$\frac{xe^{x}}{e^{x}-1} = \sum_{n=0}^{\infty} B_{n,1} \frac{x^{n}}{n!}, \quad |x| < 2\pi,$$

and since

$$\frac{xe^x}{e^x-1} = x + \frac{x}{e^x-1},$$

we see that $B_{n,1} = B_n$ for all $n \neq 1$, and $B_{1,1} = -B_1$. In fact, this can be written as $B_{n,1} = (-1)^n B_n$, and for the polynomials, $B_{n,1}(t) = (-1)^n B_n(-t)$.

2.3 DIRICHLET L-FUNCTIONS

For χ a Dirichlet character with conductor f_{χ} , the Dirichlet *L*-function for χ is defined by

$$L(s;\chi) = \sum_{b=1}^{\infty} \frac{\chi(b)}{b^s},$$

for $s \in \mathbb{C}$ such that $\Re(s) > 1$. Note that $L(s; \chi)$ can be continued analytically to all of \mathbb{C} , except for a pole of order 1 at s = 1 when $\chi = 1$.

Let $\tau(\chi)$ be a Gauss sum,

$$\tau(\chi) = \sum_{a=1}^{f_{\chi}} \chi(a) e^{2\pi i a/f_{\chi}},$$

where $i^2 = -1$, and let

$$\delta_{\chi} = \begin{cases} 0, & \text{if } \chi(-1) = 1 \\ 1, & \text{if } \chi(-1) = -1 \end{cases}$$

Then $L(s; \chi)$ satisfies the functional equation

(7)
$$\left(\frac{f_{\chi}}{\pi}\right)^{s/2} \Gamma\left(\frac{s+\delta_{\chi}}{2}\right) L(s;\chi) = W_{\chi}\left(\frac{f_{\chi}}{\pi}\right)^{(1-s)/2} \Gamma\left(\frac{1-s+\delta_{\chi}}{2}\right) L(1-s;\overline{\chi}),$$

where $\Gamma(s)$ is the gamma function, and $W_{\chi} = \frac{\tau(\chi)}{i^{\delta_{\chi}} \sqrt{f_{\chi}}}$, having the property that $|W_{\chi}| = 1$. Since $\Gamma(s)$ has simple poles at the negative integers, $L(s;\chi)$ must be zero for s = 1 - n, where $n \in \mathbb{Z}$, $n \ge 1$, such that $n \not\equiv \delta_{\chi} \pmod{2}$, except when $\chi = 1$ and n = 1. $L(s;\chi)$ can also be described by means of the Euler product $L(s;\chi) = \prod_{p \text{ prime}} (1 - \chi(p)p^{-s})^{-1}$, for $s \in \mathbb{C}$ such that $\Re(s) > 1$. Thus $L(s;\chi) \neq 0$ in this domain.

The generalized Bernoulli numbers, $B_{n,\chi}$, and the Dirichlet *L*-function, $L(s;\chi)$, share the following relationship, a proof of this being found in [13]:

THEOREM 2.1. Let χ be a Dirichlet character, and let $n \in \mathbb{Z}$, $n \ge 1$. Then $L(1-n;\chi) = -\frac{1}{n}B_{n,\chi}$.

Thus we have a way to express certain values of a function defined in terms of an infinite sum as quantities that can be found by a finite process.

2.4 The *p*-Adic number field

Let p be prime. We shall use \mathbb{Z}_p to represent the p-adic integers, and \mathbb{Q}_p the p-adic rationals. Let $|\cdot|_p$ denote the p-adic absolute value on \mathbb{Q}_p , normalized so that $|p|_p = p^{-1}$. Let $\overline{\mathbb{Q}}_p$ be the algebraic closure of \mathbb{Q}_p . The absolute value on \mathbb{Q}_p extends uniquely to $\overline{\mathbb{Q}}_p$, however $\overline{\mathbb{Q}}_p$ is not complete with respect to the absolute value. Let \mathbb{C}_p be the completion of $\overline{\mathbb{Q}}_p$ with respect to this absolute value. Then the absolute value extends to \mathbb{C}_p , and $\overline{\mathbb{Q}}_p$ is dense in \mathbb{C}_p . We also have \mathbb{C}_p algebraically closed. Furthermore, on \mathbb{C}_p

$$|a+b|_{p} \le \max\{|a|_{p}, |b|_{p}\}$$

for any $a, b \in \mathbf{C}_p$. Note that the two fields \mathbf{C} and \mathbf{C}_p are algebraically isomorphic, and any one of the two can be embedded in the other. We denote two particular subrings of \mathbf{C}_p in the following manner

$$\mathfrak{o} = \{ a \in \mathbf{C}_p : |a|_p \le 1 \}, \qquad \mathfrak{p} = \{ a \in \mathbf{C}_p : |a|_p < 1 \}.$$

Then \mathfrak{p} is a maximal ideal of \mathfrak{o} . If $\tau \in \mathbf{C}_p$ such that $|\tau|_p \leq |p|_p^s$, where $s \in \mathbf{Q}$, then $\tau \in p^s \mathfrak{o}$, and so we shall also write this as $\tau \equiv 0 \pmod{p^s \mathfrak{o}}$.

Any $n \in \mathbb{Z}$, n > 0, can be uniquely expressed in the form $n = \sum_{m=0}^{k} a_m p^m$, where $a_m \in \mathbb{Z}$, $0 \le a_m \le p-1$, for m = 0, 1, ..., k, and $a_k \ne 0$. For such n, we define

$$s_p(n) = \sum_{m=0}^k a_m,$$

the sum of the *p*-adic digits of *n*, and also define $s_p(0) = 0$. For any $n \in \mathbb{Z}$, let $v_p(n)$ be the highest power of *p* dividing *n*. This function is additive, and relates to the function $s_p(n)$ by means of the identity

(8)
$$v_p(n!) = \frac{n - s_p(n)}{p - 1},$$

which holds for all $n \ge 0$. Note that for $n \ge 1$ this implies that

$$v_p(n!) \leq \frac{n-1}{p-1}.$$

The definition of this function can be extended to all of **Q** by taking $v_p(1/n) = -v_p(n)$.

Throughout we let q = 4 if p = 2, and q = p otherwise. Note that there exist $\phi(q)$ distinct solutions, modulo q, to the equation $x^{\phi(q)} - 1 = 0$, and each solution must be congruent to one of the values $a \in \mathbb{Z}$, where $1 \le a \le q$,

(a,p) = 1. Thus, by Hensel's Lemma, given $a \in \mathbb{Z}$ with (a,p) = 1, there exists a unique $\omega(a) \in \mathbb{Z}_p$, where $\omega(a)^{\phi(q)} = 1$, such that

$$\omega(a) \equiv a \pmod{q\mathbf{Z}_p}.$$

Letting $\omega(a) = 0$ for $a \in \mathbb{Z}$ such that $(a, p) \neq 1$, we see that ω is actually a Dirichlet character, called the Teichmüller character, having conductor $f_{\omega} = q$. Let us define

$$\langle a \rangle = \omega^{-1}(a)a.$$

Then $\langle a \rangle \equiv 1 \pmod{q\mathbb{Z}_p}$. For $p \geq 3$, $\lim_{n \to \infty} a^{p^n} = \omega(a)$, since $a^{p^n} \equiv a \pmod{p}$ and $a^{p^n(p-1)} \equiv 1 \pmod{p^{n+1}}$.

For our purposes we shall need to make a slight extension of the definition of the Teichmüller character ω . If $t \in \mathbb{C}_p$ such that $|t|_p \leq 1$, then for any $a \in \mathbb{Z}$, $a + qt \equiv a \pmod{q0}$. Thus we define

$$\omega(a+qt) = \omega(a)$$

for these values of t. We also define

$$\langle a+qt\rangle = \omega^{-1}(a)(a+qt)$$

for such t.

Fix an embedding of the algebraic closure of \mathbf{Q} , $\overline{\mathbf{Q}}$, into \mathbf{C}_p . We may then consider the values of a Dirichlet character χ as lying in \mathbf{C}_p . For $n \in \mathbf{Z}$ we define the product $\chi_n = \chi \omega^{-n}$ in the sense of the product of characters. This implies that $f_{\chi_n} | f_{\chi}q$. However, since we can write $\chi = \chi_n \omega^n$, we also have $f_{\chi} | f_{\chi_n}q$. Thus f_{χ} and f_{χ_n} differ by a factor that is a power of p. In fact, either $f_{\chi_n}/f_{\chi} \in \mathbf{Z}$ and divides q, or $f_{\chi}/f_{\chi_n} \in \mathbf{Z}$ and divides q.

Let $\mathbf{Q}_p(\chi)$ denote the field generated over \mathbf{Q}_p by all values $\chi(a)$, $a \in \mathbf{Z}$. In this context we can state the following, found in [13] (pp. 14–15).

LEMMA 2.2. In the field $\mathbf{Q}_p(\chi)$, for all $n \in \mathbf{Z}$, $n \ge 0$,

$$B_{n,\chi} = \frac{1}{n+1} \lim_{h \to \infty} \frac{1}{p^h f_{\chi}} \left(B_{n+1,\chi} \left(p^h f_{\chi} \right) - B_{n+1,\chi}(0) \right) .$$

From this we can obtain

LEMMA 2.3. Let $\tau \in \mathbf{C}_p$. In the field $\mathbf{Q}_p(\chi, \tau)$, for all $n \in \mathbf{Z}$, $n \ge 0$,

$$B_{n,\chi_n}(\tau) = \lim_{h \to \infty} \frac{1}{p^h f_{\chi}} \sum_{a=1}^{p^h f_{\chi}} \chi_n(a) (a+\tau)^n \, .$$

Proof. By applying Lemma 2.2 to (4), we obtain

$$B_{n,\chi} = \lim_{h \to \infty} \frac{1}{p^h f_{\chi}} \sum_{a=1}^{p^h f_{\chi}} \chi(a) a^n \, .$$

Therefore, by (2),

$$B_{n,\chi_n}(\tau) = \sum_{m=0}^n \binom{n}{m} \tau^{n-m} \lim_{h \to \infty} \frac{1}{p^h f_{\chi_n}} \sum_{a=1}^{p^h f_{\chi_n}} \chi_n(a) a^m$$
$$= \lim_{h \to \infty} \frac{1}{p^h f_{\chi_n}} \sum_{a=1}^{p^h f_{\chi_n}} \chi_n(a) \sum_{m=0}^n \binom{n}{m} \tau^{n-m} a^m.$$

4 -

Since f_{χ} and f_{χ_n} differ by a factor that is a power of p, we must have

$$B_{n,\chi_n}(\tau) = \lim_{h \to \infty} \frac{1}{p^h f_{\chi}} \sum_{a=1}^{p^h f_{\chi}} \chi_n(a) (a+\tau)^n,$$

and the proof is complete. \Box

2.5 *p*-ADIC FUNCTIONS

Let K be an extension of \mathbf{Q}_p contained in \mathbf{C}_p . An infinite series $\sum_{n=0}^{\infty} a_n$, $a_n \in K$, converges in K if and only if $|a_n|_p \to 0$ as $n \to \infty$. Let K[[x]] be the algebra of formal power series in x. Then it follows that a power series

$$A(x) = \sum_{n=0}^{\infty} a_n x^n,$$

in K[[x]], converges at $x = \xi$, $\xi \in \mathbb{C}_p$, if and only if $|a_n\xi^n|_p \to 0$ as $n \to \infty$. Therefore whenever a power series A(x) converges at some $\xi_0 \in \mathbb{C}_p$, then it must converge at all $\xi \in \mathbb{C}_p$ such that $|\xi|_p \leq |\xi_0|_p$. The following result, for double series in K, can be found in [8]. PROPOSITION 2.4. Let $b_{n,m} \in K$, and suppose that for each $\epsilon > 0$ there exists $N \in \mathbb{Z}$, depending on ϵ , such that if $\max\{n,m\} \ge N$, then $|b_{n,m}|_p \le \epsilon$. Then both series

$$\sum_{n=0}^{\infty} \left(\sum_{m=0}^{\infty} b_{n,m} \right) \quad and \quad \sum_{m=0}^{\infty} \left(\sum_{n=0}^{\infty} b_{n,m} \right)$$

converge, and their sums are equal.

There are two power series that we wish to make note of in particular. First we define the *p*-adic exponential function, $\exp(x)$, in $\mathbb{Q}_p[[x]]$, by

(9)
$$\exp(x) = \sum_{n=0}^{\infty} \frac{1}{n!} x^n.$$

From (8) we can conclude that this power series converges in $\{x \in \mathbf{C}_p : |x|_p < p^{-1/(p-1)}\}$. The *p*-adic logarithm function, $\log(x)$, in $\mathbf{Q}_p[[x]]$, is defined by

(10)
$$\log(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} x^n,$$

the power series converging in the domain $\{x \in \mathbf{C}_p : |x|_p < 1\}$. For $|x|_p < p^{-1/(p-1)}$, we have $\log(\exp(x)) = x$ and $\exp(\log(1+x)) = 1 + x$.

The following property is a uniqueness property for power series, found in [13].

LEMMA 2.5. Let $A(x), B(x) \in K[[x]]$, such that each converges in a neighborhood of 0 in \mathbb{C}_p . If $A(\xi_n) = B(\xi_n)$ for a sequence $\{\xi_n\}_{n=0}^{\infty}, \xi_n \neq 0$, in \mathbb{C}_p , such that $\xi_n \to 0$, then A(x) = B(x).

Let U be an open subset of C_p , contained in the domain of the p-adic function f. We say that f is differentiable at $x \in U$ if the limit

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

exists. If this limit exists for each $x \in U$, then we say that f is differentiable in U.

The relationship between the derivatives of a function and its power series expansion is given in the following result, found in [8].

PROPOSITION 2.6. Let $\sum_{n=0}^{\infty} a_n x^n$ be a power series with coefficients in C_p , and suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n (x - \alpha)^n$$

converges on some closed ball B in C_p . Then

i) For each $x \in B$, the k^{th} derivative $f^{(k)}(x)$ exists, and is given by

$$f^{(k)}(x) = k! \sum_{n=k}^{\infty} {n \choose k} a_n (x - \alpha)^{n-k},$$

and we have

$$a_k = \frac{1}{k!} f^{(k)}(\alpha) \,.$$

ii) Let $\beta \in B$. Then there exists a series $\sum_{n=0}^{\infty} b_n x^n$ such that

$$f(x) = \sum_{n=0}^{\infty} b_n (x - \beta)^n$$

for any $x \in B$. Both series $\sum_{n=0}^{\infty} a_n x^n$ and $\sum_{n=0}^{\infty} b_n x^n$ have the same region of convergence.

Now let K be a finite extension of \mathbf{Q}_p . For $A(x) \in K[[x]]$, $A(x) = \sum_{n=0}^{\infty} a_n x^n$, where $a_n \in K$, define

$$||A|| = \sup_n |a_n|_p.$$

Let $P_K = \{A(x) \in K[[x]] : ||A|| < \infty\}$. Then $||\cdot||$ defines a norm on P_K , and so $K[x] \subset P_K \subset K[[x]]$. Furthermore P_K is complete in this norm.

Let $\{b_n\}_{n=0}^{\infty}$ be a sequence of elements of K, and let the sequence $\{c_n\}_{n=0}^{\infty}$ be defined by

(11)
$$c_n = \sum_{m=0}^n \binom{n}{m} (-1)^{n-m} b_m$$

for each $n \in \mathbb{Z}$, $n \ge 0$. Then $c_n \in K$ for each $n \ge 0$. Note that (11) implies that these sequences must satisfy

$$\sum_{n=0}^{\infty} c_n \frac{t^n}{n!} = e^{-t} \sum_{n=0}^{\infty} b_n \frac{t^n}{n!} \, .$$

This implies that

$$\sum_{n=0}^{\infty} b_n \frac{t^n}{n!} = e^t \sum_{n=0}^{\infty} c_n \frac{t^n}{n!},$$

and so we have the relationship

(12)
$$b_n = \sum_{m=0}^n \binom{n}{m} c_m$$

for each $n \in \mathbb{Z}$, $n \ge 0$. We can reverse this process to derive (11) given (12). Thus (11) and (12) must be equivalent. The following relate to sequences that satisfy (11) and (12), and are found in [13].

THEOREM 2.7. Let $\{b_n\}_{n=0}^{\infty}$ and $\{c_n\}_{n=0}^{\infty}$ be defined as in the above relation. Let $\rho \in \mathbf{R}$ such that $0 < \rho < |p|_p^{1/(p-1)}$. If $|c_n|_p \leq C\rho^n$ for all $n \geq 0$, where C > 0, then there exists a unique power series $A(x) \in P_K$ such that A(x) converges at every $\xi \in \mathbf{C}_p$ with $|\xi|_p < |p|_p^{1/(p-1)}\rho^{-1}$, and $A(n) = b_n$ for every $n \geq 0$.

COROLLARY 2.8. Let A(x) be the power series from the theorem. Then for each $\xi \in \mathbf{C}_p$ such that $|\xi|_p < |p|_p^{1/(p-1)}\rho^{-1}$, we have

$$A(\xi) = \sum_{n=0}^{\infty} c_n \binom{\xi}{n}.$$

Theorem 2.7 can be applied to the sequence $\{b_n\}_{n=0}^{\infty}$ in $K = \mathbf{Q}_p(\chi)$, where

$$b_n = \left(1 - \chi_n(p) p^{n-1}\right) B_{n,\chi_n}$$

in order to obtain a power series $A_{\chi}(s)$ satisfying $A_{\chi}(n) = b_n$, and converging on the domain $\{s \in \mathbf{C}_p : |s|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$. (Since $|p|_p^{1/(p-1)}|q|_p^{-1} > 1$ and $|n|_p \leq 1$ for each $n \in \mathbf{Z}$, all of \mathbf{Z} is contained in this domain.) From this a *p*-adic function, $L_p(s;\chi)$, can be derived that interpolates the values

$$L_p(1-n;\chi)=-\frac{1}{n}b_n\,,$$

and which converges in $\{s \in \mathbf{C}_p : |s-1|_p < |p|_p^{1/(p-1)}|q|_p^{-1}\}$, except $s \neq 1$ if $\chi = 1$. Note that if χ is odd, then χ_n is even when n is odd, and χ_n is odd when n is even. Thus the quantity $(1 - \chi_n(p)p^{n-1})B_{n,\chi_n} = 0$ for all $n \in \mathbf{Z}$, $n \geq 1$, as we saw from the properties of generalized Bernoulli numbers. Therefore $L_p(s;\chi)$ vanishes on a sequence such as $\{-p^m\}_{m=0}^{\infty}$, which has 0 as a limit point, implying that for such χ we must have $L_p(s;\chi) \equiv 0$.