

# **3.7 Links of 5-spheres in $S^8$**

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

By the transversality theorem ([17], IV.(2.4)), one sees that we may assume  $S_i^2 \cap g_j(S^3) = \emptyset$  for all  $i$  and  $j$ .

By Corollary 3.9, the ambient isotopy class of the embedding  $g_k$  is determined by the element  $\varphi_k := [g_k] \in \pi_3(W_k)$ ,  $W_k := W \setminus \bigcup_{j \neq k} g_j(S^3)$ ,  $k = 1, \dots, b'$ . We clearly have (compare [8])

$$\pi_3(W_k) = \pi_3\left(\underbrace{S^2 \vee \cdots \vee S^2}_{b \times} \vee \underbrace{S^3 \vee \cdots \vee S^3}_{(b'-1) \times}\right),$$

so that the Hilton-Milnor theorem yields

$$\pi_3(W_k) = \bigoplus_{i=1}^b \pi_3(S^2) \oplus \bigoplus_{1 \leq i < j \leq b} \pi_3(S^3) \oplus \bigoplus_{j \neq k} \pi_3(S^3).$$

Hence, we write  $\varphi_k$  as a tuple of integers:

$$\varphi_k = (l_i^k, i = 1, \dots, b; \quad l_{ij}^k, 1 \leq i < j \leq b; \quad \lambda_{kj}, j \neq k).$$

Observe that, for  $j \neq k$ ,  $\varphi_k$  is mapped under the natural homomorphism

$$\pi_3(W_k) \longrightarrow H_3(W_k, \mathbf{Z}) \longrightarrow H_3(W \setminus g_j(S^3), \mathbf{Z}) (\cong \mathbf{Z})$$

to the image of the fundamental class of  $S^3$  under  $g_{j*}$ . Thus,  $\lambda_{kj}$  is just the ‘usual’ linking number of the spheres  $g_k(S^3)$  and  $g_j(S^3)$  in  $W$  (compare [8]).

### 3.7 LINKS OF 5-SPHERES IN $S^8$

Let  $\mathcal{FC}_b^{\text{PL}(C^\infty)}$  be as before, and let  $\mathcal{C}_b^{\text{PL}(C^\infty)}$  be the group of isotopy classes of piecewise linear (smooth) embeddings of  $b$  disjoint copies of  $S^5$  into  $S^8$ . For  $b = 1$ , these groups are studied in [10], [19], and [20]. A brief summary with references of results in the case  $b > 1$  is contained in Section 2.6 of [11]. We will review some of this material below.

**PROPOSITION 3.15.** *We have  $\mathcal{FC}_1^{C^\infty} \cong \mathcal{FC}_1^{\text{PL}} \cong \mathbf{Z}_2$ .*

*Proof.* Since  $\pi_5(\text{SO}(3)) \cong \mathbf{Z}_2$ , the standard embedding of  $S^5$  into  $S^8$  with its two possible framings provides an injection of  $\mathbf{Z}_2$  into  $\mathcal{FC}_1^{\text{PL}(C^\infty)}$ . By Zeeman’s unknotting theorem 3.10, the map  $\mathbf{Z}_2 \longrightarrow \mathcal{FC}_1^{\text{PL}}$  is an isomorphism. As remarked in Section 2.6 of [11],  $\mathcal{FC}_1^{\text{PL}}$  is isomorphic to  $\mathcal{F}\vartheta$ , the group of h-cobordism classes of framed submanifolds of  $S^8$  which are homotopy 5-spheres. Moreover, by [10] and [19], there is an exact sequence

$$\cdots \longrightarrow \vartheta^6 \longrightarrow \mathcal{FC}_1^{C^\infty} \longrightarrow \mathcal{F}\vartheta \longrightarrow \vartheta^5 \longrightarrow \cdots.$$

As the groups  $\vartheta^5$  and  $\vartheta^6$  of exotic 5- and 6-spheres are trivial [17], our claim is settled.  $\square$

Let  $L_b \subset \mathcal{C}_b^{\mathcal{C}^\infty}$  be the subgroup of those embeddings for which the restriction to each component is isotopic to the standard embedding. As observed in Section 2.6 of [11], Zeeman's unknotting theorem 3.10 implies that  $L_b = \mathcal{C}_b^{\text{PL}}$ . The following result settles Proposition 2.3:

COROLLARY 3.16.  $\mathcal{FC}_b^{\mathcal{C}^\infty} \cong \mathcal{FC}_b^{\text{PL}} \cong L_b \oplus \bigoplus_{i=1}^b \mathbf{Z}_2$ .

For the group  $L_b$ , Theorem 1.3 of [11] provides a fairly explicit description as an extension of abelian groups. For this, consider the  $b$ -fold wedge product  $\bigvee_{i=1}^b S^2$  of 2-spheres together with its inclusion  $i: \bigvee_{i=1}^b S^2 \hookrightarrow X_{i=1}^b S^2$  into the  $b$ -fold product of 2-spheres. Finally, let  $p_i: \bigvee_{i=1}^b S^2 \rightarrow S^2$  be the projection onto the  $i^{\text{th}}$  factor,  $i = 1, \dots, b$ . Set, for  $m = 1, 2, \dots$ ,

$$\Lambda_{b,j}^m := \text{Ker}(\pi_m(p_j): \pi_m(\bigvee_{i=1}^b S^2) \rightarrow \pi_m(S^2)), \quad j = 1, \dots, b,$$

$$\Lambda_b^m := \bigoplus_{j=1}^b \Lambda_{b,j}^m$$

and

$$\Pi_b^m := \text{Ker}(\pi_m(i): \pi_m(\bigvee_{i=1}^b S^2) \rightarrow \bigoplus_{i=1}^b \pi_m(S^2)),$$

and define

$$w_b^m: \Lambda_b^m \rightarrow \Pi_b^{m+1}$$

on  $\Lambda_{b,j}^m$  by  $w_b^m(\alpha) := [\alpha, \iota_i]$ . Here,  $[., .]$  stands for the Whitehead product inside the homotopy groups of  $\bigvee_{i=1}^b S^2$  and  $\iota_i: S^2 \hookrightarrow \bigvee_{i=1}^b S^2$  for the inclusion of the  $i^{\text{th}}$  factor,  $i = 1, \dots, b$ . Theorem 1.3 of [11] yields in our situation

**THEOREM 3.17.** *There is an exact sequence of abelian groups*

$$0 \rightarrow \text{Coker}(w_b^6) \rightarrow L_b \rightarrow \text{Ker}(w_b^5) \rightarrow 0.$$

We remark that the formulas of Steer [33] might be used for the explicit computation of Whitehead products and thus for the determination

of  $\text{Coker}(w_b^6)$  and  $\text{Ker}(w_b^5)$ . The free part of  $\mathbf{L}_b$ , e.g., can be obtained quite easily. We confine ourselves to prove the following important fact.

**COROLLARY 3.18.** *The group  $\mathbf{L}_b$  has positive rank for  $b \geq 2$ .*

*Proof.* Let  $\mathbf{L}_b := \bigoplus_{l \geq 1} \mathbf{L}_{b,l}$  be the free graded Lie algebra with  $\mathbf{L}_{b,1} := \bigoplus_{i=1}^b \mathbf{Z} \cdot e_i$ . For  $l = 2, 3, \dots$ , let  $e_1^l, \dots, e_{d_l}^l$  be a basis for  $\mathbf{L}_{b,l}$  consisting of iterated commutators of the  $e_i$ . By assigning  $\iota_i$  to  $e_i$ , every iterated commutator  $c \in \mathbf{L}_{b,l}$  in the  $e_i$  defines an element  $\alpha(c) \in \pi_{l+1}(\bigvee_{i=1}^b S^2)$ .

To settle our claim, it is certainly sufficient to show that  $\text{Coker}(w_b^6)$  has positive rank. Now, by the Hilton-Milnor theorem

$$\Pi_b^7 \cong \bigoplus_{l=3}^7 \bigoplus_{k=1}^{d_{l-1}} \pi_7(S^l) \cdot \alpha(e_k^{l-1}).$$

Note that  $\pi_7(S^l)$  is finite for  $l \notin \{4, 7\}$  (see [32] and [35] for the explicit description of those groups). The Hopf fibration  $S^7 \rightarrow S^4$  [32], on the other hand, yields a decomposition  $\pi_7(S^4) \cong \pi_6(S^3) \oplus \pi_7(S^7) \cong \mathbf{Z}_{12} \oplus \mathbf{Z}$ . Therefore, it will suffice to show that the free part of  $\Lambda_b^6$  is mapped to  $\bigoplus_{j=1}^{d_6} \pi_7(S^7) \cdot \alpha(e_j^6)$ . For  $j = 1, \dots, b$ , we have

$$\Lambda_{b,j}^6 \cong \bigoplus_{i \neq j} \pi_6(S^2) \cdot \iota_i \oplus \bigoplus_{l=3}^6 \bigoplus_{k=1}^{d_{l-1}} \pi_6(S^l) \cdot \alpha(e_k^{l-1}).$$

The group  $\pi_6(S^l)$  is finite for  $l < 6$ , and we obviously have  $[\alpha(e_k^5), \iota_j] = \alpha([e_k^5, e_j])$ . If we expand the commutator  $[e_k^5, e_j]$  in the basis  $e_1^6, \dots, e_{d_6}^6$ , we find an expansion for  $[\alpha(e_k^6), \iota_j]$  in terms of the  $\alpha(e_k^6)$ .  $\square$

**COROLLARY 3.19.** *The set of  $\text{GL}_b(\mathbf{Z})$ -equivalence classes of elements in  $\mathbf{L}_b$  is infinite for  $b \geq 2$ .*

*Proof.* We have seen that the  $\text{GL}_b(\mathbf{Z})$ -set  $\mathbf{L}_{b,3}$  is contained in the  $\text{GL}_b(\mathbf{Z})$ -set  $\mathbf{L}_b$ . The  $\text{GL}_b(\mathbf{Z})$ -action on  $\mathbf{L}_{b,3}$  originates from a homomorphism  $\text{GL}_b(\mathbf{Z}) \rightarrow \text{GL}(\mathbf{L}_{b,3}) := \text{Aut}_{\mathbf{Z}}(\mathbf{L}_{b,3})$ . In particular, any matrix  $g \in \text{GL}_b(\mathbf{Z})$  preserves the absolute value of the determinant of any  $d_3$  elements in  $\mathbf{L}_{b,3}$ . This implies, for instance, that  $a \cdot e_1^3$  and  $b \cdot e_1^3$  cannot lie in the same  $\text{GL}_b(\mathbf{Z})$ -orbit, if  $0 \leq a < b$ .  $\square$