

5. Structure of the group

$\$Aut_0^{\{PL\}}(\sharp_{i=1}^b (S^2 \times S^5)) / Aut_0^{\{PL\}}(\sharp_{i=1}^b (S^2 \times D^6))\$$

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **05.06.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

In other words, a manifold X with $\delta_X \equiv 0$ is piecewise linearly (smoothly) isomorphic $X^\dagger \# X^*$ where X^* is the type of X and $b_4(X^\dagger) = 0$. As our surgery arguments above reveal, an isomorphism between $X^\dagger \# X^*$ and $X'^\dagger \# X'^*$ can be chosen of the form $\varphi^\dagger \# \varphi^*$ where $\varphi^\dagger: X^\dagger \rightarrow X'^\dagger$ and $\varphi^*: X^* \rightarrow X'^*$ are isomorphisms. Therefore, the set of isomorphy classes of based piecewise linear E-manifolds of type X^* with $b_2 = b$ is in bijection to the set of isomorphy classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$. The same goes for differentiable manifolds of type X^* , if X^* is not diffeomorphic to $X^* \# \Sigma$, Σ an exotic 8-sphere. Otherwise, we have to divide by the action of ϑ^8 . This observation together with Corollary 4.9 settles Theorem 2.4. \square

5. STRUCTURE OF THE GROUP $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$

In this section we prove that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is an abelian group which is, moreover, isomorphic to the group FL_b defined before. This result should be of some independent interest, especially because the group FL_b is quite well understood by Haefliger's work. For $b = 1$, we refer to [20] for more specific information.

We begin with the elementary

LEMMA 5.1. *Let $k \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ be a commutator. Then k extends to an automorphism of $\#_{i=1}^b(S^2 \times D^6)$.*

Proof. For the proof, we depict $\#_{i=1}^b(S^2 \times S^5)$ as follows: Let V_i , $i = 1, \dots, b$, be b copies of $S^2 \times D^6$, and we join V_i and V_{i+1} by a tube $T_i \cong [-1, 1] \times D^7$, $i = 1, \dots, b-1$. The result is a manifold W whose boundary is isomorphic to $\#_{i=1}^b(S^2 \times S^5)$. We make the following normalizations: Write ∂V_i as $(S^2 \times D_+^i) \cup (S^2 \times D_-^i)$, let n_i and s_i be the centers of D_+^i and D_-^i , respectively, and set $S_+^i := S^2 \times n_i$ and $S_-^i := S^2 \times s_i$, $i = 1, \dots, b$. Choose furthermore points $e_i \neq w_i$ in $(S^2 \times D_+^i) \cap (S^2 \times D_-^i)$, $i = 1, \dots, b$, and suppose that $\{-1\} \times D^7 \subset T_i$ is attached to a disc around w_i in ∂V_i and $\{1\} \times D^7 \subset T_i$ to a disc around e_{i+1} in ∂V_{i+1} , $i = 1, \dots, b-1$. Set $T := \bigsqcup_{i=1}^{b-1} T_i$.

Now, let $k = f \circ g \circ f^{-1} \circ g^{-1}$ with $f, g \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$. As $H_2(h, \mathbf{Z})$ is the identity for every element $h \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ and S_\pm^i , $i = 1, \dots, b$, both represent the same basis for $H_2(\partial W, \mathbf{Z})$, h is isotopic to a map h' which satisfies either assumption (A) or (B) below.

- (A) : h' is trivial on a tubular neighborhood of S_+^i which contains $(S^2 \times D_+^i) \setminus \text{Int}(T)$, $i = 1, \dots, b$.
- (B) : h' is trivial on a tubular neighborhood of S_-^i which contains $(S^2 \times D_-^i) \setminus \text{Int}(T)$, $i = 1, \dots, b$.

Next, replace f by an isotopic map f' satisfying (A), and g by an isotopic map g' satisfying (B). Then k' is isotopic to $f' \circ g' \circ f'^{-1} \circ g'^{-1}$. The map k' is the identity outside $\text{Int}(\partial T)$. It is, furthermore, the identity on a collar of $(\{-1\} \sqcup \{1\}) \times S^6$ in $R_i := [-1, 1] \times S^6 \subset \partial T_i$, $i = 1, \dots, b-1$. Let k'_i be the restriction of k' to R_i , $i = 1, \dots, b$. We know that each k'_i is the identity on a collar of $\{-1, 1\} \times S^6$ in R_i . Thus, we extend every k'_i to a PL automorphism \tilde{k}_i of $D^7 \times \{-1\} \cup R_i \cup D^7 \times \{1\} \cong S^7$ through $\text{id}_{D^7 \times \{-1\} \cup D^7 \times \{1\}}$. Now, by [27], Lemma 1.10, p. 8, \tilde{k}_i extends to an automorphism κ_i of $D^8 \cong D^7 \times [-1, 1]$, $i = 1, \dots, b$. Thus, the maps id_{V_i} and α_i , $i = 1, \dots, b$, glue to an automorphism of $\#_{i=1}^b(S^2 \times D^6)$ whose restriction to the boundary is just k' . \square

This lemma shows that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is a normal subgroup of $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$, and that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is abelian. Moreover, in Section 4.3, we have already defined a set theoretic bijection

$$\beta: \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6)) \longrightarrow \text{FL}_b.$$

THEOREM 5.2. *The map β is a group isomorphism.*

Proof. Since β is bijective, we have to verify that β is a homomorphism. In order to do so, we will construct a group \mathbf{G} together with surjective homomorphisms

$$\chi_1: \mathbf{G} \longrightarrow \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$$

and

$$\chi_2: \mathbf{G} \longrightarrow \text{FL}_b,$$

such that $\chi_2 = \beta \circ \chi_1$. This will clearly settle the claim.

Before we define \mathbf{G} , we recall some constructions and conventions from [11]. Let $S^8 = \{(x_0, \dots, x_9) \in \mathbf{R}^9 \mid x_0^2 + \dots + x_9^2 = 1\}$ be the unit sphere, write $S^8 = D_+^8 \cup D_-^8$, and let $\sigma: S^8 \rightarrow S^8$ be the reflection at $S^7 = D_+^8 \cap D_-^8$, interchanging the Northern and the Southern hemispheres. First, let $S_b := (S_1^5, \dots, S_b^5)$ be a ‘standard link’ in S^8 defined as follows: Fix real numbers $-1/2 < a_1 < \dots < a_b < 1/2$, and set

$$S_i^5 := \{ (x_0, \dots, x_9) \in S^8 \mid x_6 = x_7 = x_8 = 0, x_9 = a_i \}.$$

We choose, furthermore, framings $\tau_i: S_i^5 \times D^3 \rightarrow S^8$ which extend over D^6 , such that $\tau_i(D_{i,\pm}^5 \times D^3) \subset D_{\pm}^8$ and $\sigma \circ \tau_i = \tau_i \circ (\sigma|_{S_i^5} \times \text{id}_{D^3})$, $i = 1, \dots, b$. Let l_b^0 be the resulting framed link in S^8 with $l_{b,\pm}^0 := l_b^0 \cap D_{\pm}^8$. Recall from Section 1 of [11] that

1. Every framed link l of b five-dimensional spheres in S^8 is isotopic to a link l' , such that either (A) $l' \cap D_+^8 = l_{b,+}^0$ or (B) $l' \cap D_-^8 = l_{b,-}^0$.
2. If l_1 satisfies (A) and l_2 satisfies (B), then $l_1 + l_2$ is represented by the link l with $l \cap D_+^8 = l_2 \cap D_+^8$ and $l \cap D_-^8 = l_1 \cap D_-^8$.

Note that, if we perform surgery along l_b^0 , we get a manifold $W = W_+ \cup W_-$ which is isomorphic to $\#_{i=1}^b (S^2 \times S^6)$, and

$$W_{\pm} := (D_{\pm}^8 \setminus \text{Int}(l_b^0)) \cup \left(\bigsqcup_{i=1}^b (S_i^2 \times D_{\pm}^6) \right)$$

is canonically isomorphic to $\#_{i=1}^b (S^2 \times D^6)$. For the rest of the proof, we will use the description of $\#_{i=1}^b (S^2 \times S^5)$ as $\partial W_+ = \partial W_-$. Set

$$\mathbf{G} := \{ \text{PL-maps } f: S^7 \setminus \text{Int}(l_b^0) \rightarrow S^7 \setminus \text{Int}(l_b^0) : f|_{\text{boundary}} = \text{id} \}.$$

For every $f \in \mathbf{G}$, we define $\varphi(f): \#_{i=1}^b (S^2 \times S^5) \rightarrow \#_{i=1}^b (S^2 \times S^5)$, by extending f through the identity on $\bigsqcup_{i=1}^b (S_i^2 \times D^5)$. Similarly, define $\psi(f): S^7 \rightarrow S^7$. Obviously,

$$\begin{aligned} \chi_1: \mathbf{G} &\longrightarrow \text{Aut}_0^{\text{PL}}(\#_{i=1}^b (S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b (S^2 \times D^6)) \\ f &\longmapsto [\varphi(f)] \end{aligned}$$

is a surjective homomorphism.

Next, we associate to $f \in \mathbf{G}$ an element $\chi_2(f) \in \text{FL}_b$ as follows: First, we define $\Sigma(f) := D_+^8 \cup_{\psi(f)} D_-^8$ and the link $l'(f) := l_{b,+}^0 \cup_{\psi(f)} l_{b,-}^0$. Then we choose a piecewise linear homeomorphism $F: \Sigma(f) \rightarrow S^8$ and set $l_F(f) := F(l'(f))$. We have checked before that the isotopy class of $l_F(f)$ does not depend on the chosen homeomorphism, so that $\chi_2(f) := [l_F(f)] \in \text{FL}_b$ is well defined. To see that $\chi_2: \mathbf{G} \rightarrow \text{FL}_b$ is a homomorphism, let f, f' be in \mathbf{G} . Choose extensions $\bar{\psi}: D_+^8 \rightarrow D_+^8$ and $\bar{\psi}': D_-^8 \rightarrow D_-^8$ of $\psi(f)$ and $\psi(f')$, respectively. We then define $F: \Sigma(f) \rightarrow S^8$ as $\bar{\psi}$ on D_+^8 and as the identity on D_-^8 , $F': \Sigma(f') \rightarrow S^8$ as the identity on D_+^8 and $(\bar{\psi}')^{-1}$ on D_-^8 , and $F'': \Sigma(f' \circ f) \rightarrow S^8$ as $\bar{\psi}$ on D_+^8 and $(\bar{\psi}')^{-1}$ on D_-^8 . Then the link $l_F(f)$ satisfies (B), the link $l_{F'}(f')$ satisfies (A), and (2) above shows that $[l_{F''}(f' \circ f)] = [l_{F'}(f')] + [l_F(f)]$.

Finally, for given $f \in \mathbf{G}$, we can perform surgery on $\Sigma(f)$ along $l'(f)$. The result is $W_+ \cup_{\varphi(f)} W_-$. Reading this backwards means nothing else but $\beta(\chi_1(f)) = \chi_2(f)$ and we are done. \square