

# 4. Prime parts of torsion numbers

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## 4. PRIME PARTS OF TORSION NUMBERS

We recall Jensen's formula, a short argument for which can be found in [Yo86].

LEMMA 4.1 [Jensen's formula]. *For any complex number  $\alpha$ ,*

$$\int_0^1 \log |\alpha - e^{2\pi i\theta}| d\theta = \log \max\{1, |\alpha|\}.$$

By Lemma 4.1 the Mahler measure  $M(f)$  of a nonzero polynomial with complex coefficients can be computed as

$$\exp \int_0^1 \log |f(e^{2\pi i\theta})| d\theta.$$

This observation motivated the definition of Mahler measure for polynomials in several variables. (See [Bo81] or [EW99], for example.)

In [EF96], [Ev99] G.R. Everest and B. Ní Fhlathúin proved a  $p$ -adic analogue of Jensen's formula, which we describe. Assume that  $\alpha$  is an algebraic integer lying in a finite extension  $K$  of  $\mathbf{Q}$ . For every prime  $p$  there is a  $p$ -adic absolute value  $|\cdot|_p$ , the usual Archimedean absolute value corresponding to  $\infty$ . We recall the definition (see [La65] for more details): If  $p$  is a prime number, then  $|p^r m/n|_p = 1/p^r$ , where  $r$  is an integer, and  $m, n$  are nonzero integers that are not divisible by  $p$ . By convention,  $|0|_p = 0$ . Each  $|\cdot|_p$  extends to an absolute value  $|\cdot|_v$  on  $K$ . Let  $\Omega_v$  denote the smallest field which is algebraically closed and complete with respect to  $|\cdot|_v$ . Let  $\mathbf{T}_v$  denote the closure of the group of all roots of unity, which is in general locally compact. Note that if  $p = \infty$ , then  $\Omega_v = \mathbf{C}$  and  $\mathbf{T}_v = \mathbf{T}$ . Everest and Fhlathúin define

$$M_{\mathbf{T}_v}(t - \alpha) = \exp \int_{\mathbf{T}_v} \log |t - \alpha|_v d\mu = \exp \lim_{r \rightarrow \infty} \frac{1}{r} \sum_{\zeta^r=1} \log |\zeta - \alpha|_v.$$

Here  $\int$  denotes the Shnirelman integral, given by the limit of sums at the right, where one skips over the undefined summands. The above integral exists even if  $\alpha \in \mathbf{T}_v$ , in which case it can be shown to be zero. Moreover, one has

$$(4.1) \quad \int_{\mathbf{T}_v} \log |t - \alpha|_v d\mu = \log \max\{1, |\alpha|_v\},$$

which Everest and Fhlathúin refer to as a  $p$ -adic analogue of Jensen's formula.

Recall that  $b_r^{(p)}$  denotes the  $p$ -component of  $b_r$ , the largest power of  $p$  that divides  $b_r$ . The *content* of  $f \in \mathbf{Z}[t]$  is the greatest common divisor of the coefficients. Using (4.1) we will prove

**THEOREM 4.2.** *Let  $(G, \chi)$  be an augmented group, and let  $p$  be a prime.*

- (i) *If  $\mathcal{M}$  has a square matrix presentation and  $\Delta(t) \neq 0$ , then the sequence  $\{b_{r_k}\}$  of pure torsion numbers satisfies*

$$\lim_{r_k \rightarrow \infty} (b_{r_k}^{(p)})^{1/r_k} = (\text{content } \Delta)^{(p)}.$$

- (ii) *If  $\mathcal{M}$  is a direct sum of cyclic modules, then the sequence of all torsion numbers satisfies*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = (\text{content } \Delta)^{(p)}.$$

- (iii) *If  $\mathcal{M}$  is torsion free as an abelian group, then*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = 1.$$

**EXAMPLE 4.3.** For any positive integer  $m$ , consider the augmented group  $(G, \chi)$  where  $G$  is the Baumslag-Solitar group  $\langle x, y \mid y^m x = xy^m \rangle$  and  $\chi: G \rightarrow \mathbf{Z}$  maps  $x \mapsto 1$  and  $y \mapsto 0$ . One verifies that  $\mathcal{M} \cong \mathcal{R}_1/(m(t-1))$ . The quotient module  $\mathcal{M}_r$  is isomorphic to  $\mathbf{Z}^r/A_r\mathbf{Z}^r$ , where

$$A_r = \begin{pmatrix} m & 0 & 0 & 0 & \cdots & -m \\ -m & m & 0 & \cdots & & 0 \\ 0 & -m & m & 0 & \cdots & 0 \\ & & & \vdots & & \\ 0 & 0 & & \cdots & -m & m \end{pmatrix}.$$

The matrix is equivalent by elementary row and column operations to the diagonal matrix

$$\begin{pmatrix} m & & & & & \\ & \ddots & & & & \\ & & m & & & \\ & & & & & \\ & & & & & 0 \end{pmatrix}.$$

Hence  $\mathcal{M}_r \cong \mathbf{Z} \oplus (\mathbf{Z}/m)^{r-1}$ , and so  $b_r = m^{r-1}$  for all  $r$ . Consequently,

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = m^{(p)}.$$

The Alexander polynomial of any knot is nonzero, and its coefficients are relatively prime. Hence the following corollary is immediate from Theorem 4.2 (iii).

COROLLARY 4.4. *For any knot  $k$  and prime  $p$ ,*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = 1.$$

Theorem 2.10 and Corollary 4.4 imply that whenever the Alexander polynomial of  $k$  has Mahler measure greater than 1, infinitely many distinct primes occur in the factorization of the torsion numbers  $b_r$ . In other words, the homology groups  $H_1(M_r, \mathbf{Z})$  display nontrivial  $p$ -torsion for infinitely many primes  $p$ . Since the sequence  $\{b_r\}$  is a division sequence, the number of prime factors of  $b_r$  is unbounded.

What about the case in which the Alexander polynomial of  $k$  has Mahler measure equal to 1? The argument of Section 5.7 of [Go72] shows that the number of prime factors remains unbounded as long as the Alexander polynomial does not divide  $t^M - 1$  for any  $M$ . If it does divide, then the torsion numbers  $b_r$  are periodic by Section 5.3 of [Go72] (see also Corollary 2.2 of [SiWi00]). Hence we obtain

COROLLARY 4.5. *For any knot, either the torsion numbers  $b_r$  are periodic or else for any  $N > 0$  there exists an  $r$  such that the factorization of  $b_r$  has at least  $N$  distinct primes.*

The proof of Theorem 4.2 requires the following lemma.

LEMMA 4.6. *If  $f(t) = c_0 t^n + \cdots + c_{n-1} t + c_n$  is a nonzero polynomial in  $\mathbf{Z}[t]$  with roots  $\lambda_1, \dots, \lambda_n$  (not necessarily distinct) in  $\Omega_v$ , then*

$$|c_0|_v \prod_{i=1}^n \max\{1, |\lambda_i|_v\} = |\text{content } f|_v.$$

*Proof.* The argument that we present is found in [LW88]. Set  $a_j = c_j/c_0$  for  $0 \leq j \leq n$ , so  $f(t) = c_0(t^n + a_1 t^{n-1} + \cdots + a_n)$ . Each  $a_j$  is an elementary symmetric function of the roots  $\lambda_i$ , namely the sum of products of roots taken  $j$  at a time. Using the ultrametric property

$$|x + y|_v = \max\{|x|_v, |y|_v\},$$

we see that if exactly  $k$  values of  $|\lambda_i|_v$  are greater than 1, then

$$\max_j |a_j|_v = |a_k|_v = \prod_{j=1}^n \max\{1, |\lambda_j|_v\}.$$

But

$$\max_j |a_j|_v = \max \left\{ 1, \left| \frac{c_1}{c_0} \right|_v, \dots, \left| \frac{c_n}{c_0} \right|_v \right\} = \frac{|\text{content } f|_v}{|c_0|_v}.$$

Hence the lemma is proved.  $\square$

*Proof of Theorem 4.2.* In case (i), the pure torsion number  $b_{r_k}$  is equal to  $\left| \prod_{\zeta^{r_k}=1} \Delta(\zeta) \right|$ . We have

$$|b_{r_k}|_v = \left| \prod_{\zeta^{r_k}=1} \Delta(\zeta) \right|_v = |c_0|_v^{r_k} \prod_{\zeta^{r_k}=1} \prod_{j=1}^n |\zeta - \lambda_j|_v,$$

where  $c_0$  is the leading coefficient of  $\Delta$  and  $\lambda_1, \dots, \lambda_n$  are its roots. Hence

$$\begin{aligned} |b_{r_k}|_v^{1/r_k} &= |c_0|_v \prod_{\zeta^{r_k}=1} \prod_{j=1}^n |\zeta - \lambda_j|_v^{1/r_k} \\ &= |c_0|_v \prod_{j=1}^n \exp \left( \frac{1}{r_k} \sum_{\zeta^{r_k}=1} \log |\zeta - \lambda_j| \right), \end{aligned}$$

so that

$$\lim_{r_k \rightarrow \infty} |b_{r_k}|_v^{1/r_k} = |c_0|_v \prod_{j=1}^n \exp \int_{\mathbf{T}_v} \log |t - \lambda_j|_v \, d\mu,$$

which by equation (4.1) is equal to

$$|c_0|_v \prod_{j=1}^n \max \{1, |\lambda_j|_v\}.$$

By Lemma 4.6 this is equal to  $|\text{content } \Delta|_v$ . But for integers  $n$  we have  $n^{(p)} = |n|_v^{-1}$ .

Now suppose  $\mathcal{M}$  is cyclic. As in the proof of Theorem 3.8, we let  $\gamma$  be the cyclotomic order of  $\Delta$  and consider the subsequence of  $b_r$  with  $r$  in a fixed congruence class modulo  $\gamma$ . Then starting with Theorem 3.3 we may apply the argument above with  $\Delta/\Phi$  in place of  $\Delta$  to show that the limit of  $(|b_r|^{(p)})^{1/r}$  along this subsequence is the  $p$ -component of the content of  $\Delta/\Phi$ . But content is multiplicative and cyclotomic polynomials have content 1, so the limit along all congruence classes is  $(\text{content } \Delta)^{(p)}$ . The result is immediate for direct sums of cyclic modules.

Finally, we can extend the result when  $\mathcal{M}$  is torsion-free as an abelian group using Theorem 3.6. But for this case the content of  $\Delta$  is 1.  $\square$