

4. CONSEQUENCES OF GROMOV HYPERBOLICITY FOR THE SHAPE OF THE BOUNDARY

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **05.06.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

is bounded in view of the estimate $\kappa^{-1}(x^2 + y^2) < f(x, y) < \kappa(x^2 + y^2)$ for some universal $\kappa > 0$ and of the fact that $f(x, y) = \varepsilon$ on C_ε . \square

4. CONSEQUENCES OF GROMOV HYPERBOLICITY FOR THE SHAPE OF THE BOUNDARY

PROPOSITION 4.1. *Let D be a bounded convex domain in \mathbf{R}^n and let h be a Hilbert metric on D . If h is Gromov hyperbolic then the boundary ∂D is strictly convex, that is, it does not contain a line segment.*

This can be proven following the proof of N. Ivanov [Iv97] of Masur-Wolf's theorem [MW95] that the Teichmüller spaces (genus ≥ 2) are not Gromov hyperbolic. The proof makes use of Gromov's exponential divergence criterion, see [BH99, p. 412]. For another proof of the above proposition, see [SM00].

THEOREM 4.2. *Let D be a bounded convex domain in \mathbf{R}^n and let h be the Hilbert metric on D . If h is Gromov hyperbolic then the boundary ∂D is smooth of class C^1 .*

Proof. *2-dimensional case:* First, by the previous result, D is strictly convex. Let $y = f(x)$, $x \in (-a, a)$ be an equation of ∂D near some point. Then f is strictly convex and hence the one-sided derivatives $f'_-(x)$, $f'_+(x)$ exist and are strictly increasing on $(\varepsilon, \varepsilon)$, [RV73, § 11].

We prove that $f'_-(0) = f'_+(0)$. Suppose not, then by choosing appropriate Cartesian coordinates we may assume that $f'_-(0) < 0$ and $f'_+(0) > 0$. For each sufficiently small ε construct an ideal triangle $\Delta = \Delta(\varepsilon)$ in D with one vertex 0 and two other vertices corresponding to the intersection of the line $y = \varepsilon$ with ∂D . We assert that the slimness of $\Delta(\varepsilon)$ tends to ∞ when ε tends to zero. Namely we show that the Hilbert distance between the point $P = (0, \varepsilon)$ and any point Q of the side $[0, B]$ tends to ∞ . Let $f'_+(0) = \tan \alpha$, $0 < \alpha < \pi/2$. Let $x_1 < x_2$ be the points such that $f(x_1) = \varepsilon$ and $f'_+(0)x_2 = \varepsilon$. Then

$$PQ \geq \varepsilon \cos \alpha = f(x_1) \cos \alpha.$$

Let O, R be the intersection points of the line PQ with ∂D . We have therefore

$$QR \leq x_2 - x_1 = \frac{f(x_1)}{f'_+(0)} - x_1 = \frac{f(x_1) - f'_+(0)x_1}{f'_+(0)}$$

and hence, combining the last two inequalities,

$$\begin{aligned} \frac{PQ}{QR} &\geq \frac{f'_+(0)f(x_1)\cos\alpha}{f(x_1) - f'_+(0)x_1} \\ &= \frac{f'_+(0)\cos\alpha}{1 - f'_+(0)\frac{x_1}{f(x_1)}} \rightarrow \infty \text{ when } x_1 \rightarrow 0. \end{aligned}$$

It follows that

$$h(P, Q) = \ln \left(1 + \frac{PQ}{OP} \right) \left(1 + \frac{PQ}{QR} \right) \rightarrow \infty \text{ when } x_1 \rightarrow 0$$

and hence the slimness of $\Delta(\varepsilon)$ tends to ∞ when ε tends to zero.




FIGURE 4
Hyperbolicity implies C^1

It remains to show that f' is continuous. By [RV73, §14] we have

$$\begin{aligned} \lim_{x \rightarrow x_0+} f'_+(x) &= f'_+(x_0), \\ \lim_{x \rightarrow x_0-} f'_+(x) &= f'_-(x_0). \end{aligned}$$

From this we conclude that f'_+ is continuous at x_0 since $f'_+(x_0) = f'_-(x_0)$. But $f'(x_0) = f'_+(x_0)$ hence f' is also continuous at x_0 .

n-dimensional case: Recall the known result that if f is a differentiable convex function defined on an open convex set S in \mathbf{R}^{n+1} , then it is C^1 on S , see for example [RV73]. Let D be a bounded convex domain in \mathbf{R}^{n+1} , $n \geq 2$. It is enough to prove that ∂D is differentiable at any point. Given a point $p \in \partial D$, we can choose the coordinate axis of \mathbf{R}^{n+1} so that the origin O of the coordinates is at p , all of D lies in the halfspace $x_0 \geq 0$ and in a neighbourhood of p the surface ∂D can be represented as the graph of a nonpositive convex function $x_0 = f(x_1, x_2, \dots, x_n)$, $x = (x_1, x_2, \dots, x_n)$, $f(0) = 0$. Considering the 2-dimensional sections in the planes x_0 , x_i , $i = 1, \dots, n$, we obtain that the partial derivatives of f at 0 exist and $f_{x_i}(0) = 0$, $i = 1, \dots, n$. We have to prove that for each $\varepsilon > 0$ there is a

neighbourhood U_ε of 0 such that $f(x) < \varepsilon|x|$ in this neighbourhood. But in view of $f_{x_i}(0) = 0$, $i = 1, \dots, n$, we have $f(0, \dots, 0, x_i, 0, \dots, 0) < \varepsilon|x_i|$ for sufficiently small x_i and hence by convexity $f(x) < \varepsilon|x|$ for sufficiently small $|x|$. \square

REMARK 4.3. The following was announced in [B00]: *If a strictly convex domain D is divisible, that is, if it admits a proper cocompact group of isometries Γ , then D is Gromov hyperbolic if and only if ∂D is C^1 .* Our Theorem 4.2 shows that in the implication (Gromov hyperbolicity + divisibility $\Rightarrow C^1$) the condition of divisibility is superfluous.

5. NON-STRICELY CONVEX DOMAINS

This section owes much of its existence to [Be97] and [Be99]. Using a different argument, we prove certain extensions to arbitrary convex bounded domains of some of the results obtained in those papers.

LEMMA 5.1. *Let D be a bounded convex domain in \mathbf{R}^n . Let $\{x_n\}, \{y_n\}$ be two sequences of points in D . Assume that $x_n \rightarrow \bar{x} \in \partial D$, $y_n \rightarrow \bar{y} \in \overline{\partial D}$ and $[\bar{x}, \bar{y}] \not\subseteq \partial D$. Let x'_n and y'_n denote the endpoints of the chord through x_n and y_n as usual. Then x'_n converges to \bar{x} and y'_n converges to the endpoint \bar{y}' of the chord defined by \bar{x} and \bar{y} different from \bar{x} .*

Proof. Compare with Lemma 5.3. in [Be97]. Every limit point of chord endpoints must belong to the line through \bar{x} and \bar{y} . In addition, in the case of x'_n for example, any limit point must lie on the halfline from \bar{x} not containing \bar{y} . At the same time each limit point must belong to the boundary of D , and the statement follows since the line through \bar{x} and \bar{y} intersects ∂D only in \bar{x} and \bar{y}' . \square

THEOREM 5.2. *Let D be a bounded convex domain. Let $\{x_n\}$ and $\{z_n\}$ be two sequences of points in D . Assume that $x_n \rightarrow \bar{x} \in \partial D$, $z_n \rightarrow \bar{z} \in \partial D$ and $[\bar{x}, \bar{z}] \not\subseteq \partial D$. Then there is a constant $K = K(\bar{x}, \bar{z})$ such that for the Gromov product $(x_n \mid z_n)_y$ in Hilbert distances relative to some fixed point y in D we have*

$$\limsup_{n \rightarrow \infty} (x_n \mid z_n)_y \leq K.$$