

7. Appendix: further examples

Objekttyp: **Appendix**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

shows that \bar{x} commutes with $\bar{x}\bar{y}\bar{x}^{-1}\bar{y}^{-1}$ in the tile path group. Then Theorem 6.1(a) shows that $\pi(\mathcal{T}) \cong \mathbf{Z}/9\mathbf{Z}$. This means that the tile homotopy group only detects area, modulo 9.

On the other hand, we can easily show that if \mathcal{T} tiles a rectangle, then both sides must be even. Consider the ways that a tile can touch the edge of a rectangle.




FIGURE 6.9
Tiles along an edge of a rectangle

We see that the first two possibilities cannot occur, so each tile that touches the edge does so along an even length. Therefore, each edge of the rectangle has even length. In fact, it is not much harder to show that if \mathcal{T} tiles an $m \times n$ rectangle, then both m and n are multiples of 6. A straightforward argument shows that every tiling of a quadrant by \mathcal{T} is a union of 6×6 squares, which implies the result.

7. APPENDIX: FURTHER EXAMPLES

Here we give some more tiling restrictions we have found using the tile homotopy technique. In each case, there are signed tilings that show that the result cannot be obtained by tile homology methods, and there are tilings that show that the result is non-vacuous. Further details will be published elsewhere.

THEOREM 7.1. Let $\mathcal{T} = \{\begin{array}{c} \square \\ \square \end{array}, \begin{array}{c} \square \quad \square \\ \square \end{array}, \begin{array}{c} \square \quad \square \\ \square \quad \square \end{array}\}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then either m or n is a multiple of 4.
- (b) A 1×6 rectangle has a signed tiling by \mathcal{T} .


THEOREM 7.2. Let $\mathcal{T} = \{\begin{array}{c} \square \\ \square \end{array}, \begin{array}{c} \square \quad \square \\ \square \quad \square \end{array}, \begin{array}{c} \square \quad \square \\ \square \quad \square \\ \square \quad \square \end{array}\}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.
- (b) A 1×6 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.3. Let $\mathcal{T} = \{\square \begin{smallmatrix} & \\ & \vdots & \\ & & \end{smallmatrix} \}$, where rotations are permitted, but reflections are not.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is even.
- (b) A 1×5 rectangle has a signed tiling by \mathcal{T} .


REMARK 7.4. It is easy to show that if \mathcal{T} tiles a rectangle, then both sides are multiples of 5. Also, Yuri Aksyonov [1] has given a clever geometric proof that one side must be a multiple of 10.



THEOREM 7.5. Let $\mathcal{T} = \{\square \begin{smallmatrix} & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}\}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.
- (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .


THEOREM 7.6. Let



where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.
- (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .


THEOREM 7.7. Let $\mathcal{T} = \{\square \begin{smallmatrix} & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}\}$,



- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.
- (b) A 1×2 rectangle has a signed tiling by \mathcal{T} .

THEOREM 7.8. Let $\mathcal{T} = \{\square \begin{smallmatrix} & & \\ & \vdots & \\ & & \end{smallmatrix}, \square \begin{smallmatrix} & & & \\ & \vdots & & \\ & & & \end{smallmatrix}\}$, where all orientations are allowed.


- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.
- (b) A 2×2 square has a signed tiling by \mathcal{T} .



THEOREM 7.9. Let $\mathcal{T} = \{ \text{[T-shaped tile]}, \text{[cross-shaped tile]} \}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then either m is a multiple of 3 or n is a multiple of 6.


(b) A 1×1 square has a signed tiling by \mathcal{T} .



THEOREM 7.10. Let $\mathcal{T} = \{ \text{[U-shaped tile]}, \text{[cross-shaped tile]}, \text{[horizontal bar]} \}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 8.


(b) A 1×1 square has a signed tiling by \mathcal{T} .



THEOREM 7.11. Let $\mathcal{T} = \{ \text{[horizontal bar]}, \text{[cross-shaped tile]}, \text{[T-shaped tile]} \}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 5.


(b) A 1×1 square has a signed tiling by \mathcal{T} .



THEOREM 7.12. Let $\mathcal{T} = \{ \text{[horizontal bar]}, \text{[cross-shaped tile]} \}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 4.


(b) A 1×2 rectangle has a signed tiling by \mathcal{T} .



THEOREM 7.13. Let $\mathcal{T} = \{ \text{[horizontal bar]}, \text{[2x2 square]}, \text{[cross-shaped tile]} \}$, where all orientations are allowed.

(a) If \mathcal{T} tiles an $m \times n$ rectangle, then mn is a multiple of 4.


(b) A 1×2 rectangle has a signed tiling by \mathcal{T} .



THEOREM 7.14. Let $\mathcal{T} = \{ \text{[T-shaped tile]}, \text{[horizontal bar]}, \text{[cross-shaped tile]} \}$, where all orientations are allowed.


(a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.

(b) A 1×1 square has a signed tiling by \mathcal{T} .



THEOREM 7.15. Let $\mathcal{T} = \{ \text{[L-shaped polyominoes]} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles an $m \times n$ rectangle, then one of m or n is a multiple of 6.
- (b) A 2×3 rectangle has a signed tiling by \mathcal{T} .



THEOREM 7.16. Let $\mathcal{T} = \{ \text{[triangle divided into six smaller triangles]} \}$, where all orientations are allowed.

- (a) If \mathcal{T} tiles a triangle of side n , then n is a multiple of 8.
- (b) A triangle of side 4 has a signed tiling by \mathcal{T} .

REMARK 7.17. That \mathcal{T} tiles any triangle is quite interesting. Karl Scherer [15, 2.6 D] has found a tiling of a side 32 triangle by \mathcal{T} .

ACKNOWLEDGMENT. I thank Torsten Sillke for some interesting discussions.

REFERENCES

- [1] AKSYONOV, YU. E-mail communication to Torsten Sillke. March 1999 (<http://www.mathematik.uni-bielefeld.de/~sillke/PENTA/qu5-y-right>).
- [2] BERLEKAMP, E. R., J. H. CONWAY and R. K. GUY. *Winning Ways for Your Mathematical Plays*, vol. 2. Academic Press, London, 1982.
- [3] BLACK, M. *Critical Thinking*. Prentice-Hall, New York, 1946.
- [4] CONWAY, J. H. and J. C. LAGARIAS. Tiling with polyominoes and combinatorial group theory. *J. Combin. Theory Ser. A*, 53 (1990), 183–208.
- [5] THE GAP GROUP. GAP – Groups, Algorithms and Programming, version 4.3, 2002 (<http://www.gap-system.org>).
- [6] GAREY, M. R. and D. S. JOHNSON. *Computers and Intractability*. Freeman, San Francisco, 1979.
- [7] GOLOMB, S. W. Checker boards and polyominoes. *Amer. Math. Monthly* 61 (1954), 675–682.
- [8] —— Covering a rectangle with L -tetrominoes, Problem E1543. *American Mathematical Monthly* 69 (1962), 920.
Solution by D. A. KLARNER: *Amer. Math. Monthly* 70 (1963), 760–761.
- [9] HALL, M. JR. *The Theory of Groups*. Chelsea, New York, 1976.
- [10] KLARNER, D. A. Packing a rectangle with congruent N -ominoes. *J. Combin. Theory* 7 (1969), 107–115.
- [11] LANGMAN, H. *Play Mathematics*. Hafner, New York, 1962.