

5.2 The basic 3-form on G

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

5.2 THE BASIC 3-FORM ON G

Let $\theta^L, \theta^R \in \Omega^1(G, \mathfrak{g})$ be the left- and right-invariant Maurer-Cartan forms on G , respectively. The 3-form $\eta \in \Omega^3(G)$ given by³⁾

$$\eta = \frac{1}{12} \theta^L \cdot [\theta^L, \theta^L] = \frac{1}{12} \theta^R \cdot [\theta^R, \theta^R]$$

is closed, and has a closed equivariant extension $\eta_G \in \Omega_G^3(G)$ given by

$$\eta_G(\xi) := \eta - \frac{1}{2}(\theta^L + \theta^R) \cdot \xi.$$

Their cohomology classes represent generators of $H^3(G, \mathbf{Z}) = \mathbf{Z}$ and $H_G^3(G, \mathbf{Z}) = \mathbf{Z}$, respectively. The pull-back of η_G to any conjugacy class $\iota_C : C \hookrightarrow G$ is exact. In fact, let $\omega_C \in \Omega^2(C)^G \subset \Omega_G^2(C)$ be the invariant 2-form given on generating vector fields ξ_C, ξ'_C for $\xi, \xi' \in \mathfrak{g}$ by the formula

$$\omega_C(\xi_C(g), \xi'_C(g)) = \frac{1}{2}\xi \cdot (\text{Ad}_g - \text{Ad}_{g^{-1}})\xi'.$$

Then [1, 16]

$$d_G \omega_C + \iota_C^* \eta_G = 0.$$

We will now show that η_G is exact over each of the open subsets V_j . Let $C_j = q^{-1}(\mu_j) \subset V_j$ be the conjugacy classes corresponding to the vertices.

LEMMA 5.1. *The linear retraction*

$$[0, 1] \times \mathfrak{A}_j \rightarrow \mathfrak{A}_j, \quad (t, \mu_j + \zeta) \mapsto \mu_j + (1-t)\zeta$$

of \mathfrak{A}_j onto the vertex μ_j lifts uniquely to a smooth G -equivariant retraction from V_j onto C_j .

Proof. Recall that the slice S_j is an open neighborhood of $\exp(\mu_j)$ in G_j . Any G_j -equivariant retraction from S_j onto $\exp \mu_j$ extends uniquely to a G -equivariant retraction from $V_j = G \times_{G_j} S_j$ onto C_j . Note that $S'_j = G_j \cdot (\mathfrak{A}_j - \mu_j)$ is a star-shaped open neighborhood of 0 in \mathfrak{g}_j , and that $S'_j \rightarrow S_j$, $\zeta \mapsto \exp(\mu_j) \exp(\zeta)$ is a G_j -equivariant diffeomorphism. The linear retraction of S'_j onto the origin gives the desired retraction of S_j . Uniqueness is clear, since the retraction has to preserve $\exp(\mathfrak{A}_j) \subset V_j$, by equivariance.

³⁾ For \mathfrak{g} -valued forms β_1, β_2 , the bracket $[\beta_1, \beta_2]$ denotes the \mathfrak{g} -valued form obtained by applying the Lie bracket $[\cdot, \cdot] : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ to the $\mathfrak{g} \otimes \mathfrak{g}$ -valued form $\beta_1 \wedge \beta_2$.

Let

$$\mathbf{h}_j: \Omega^p(V_j) \rightarrow \Omega^p([0, 1] \times V_j) \rightarrow \Omega^{p-1}(V_j)$$

be the de Rham homotopy operator for this retraction, given (up to a sign) by pull-back under the retraction, followed by integration over the fibers of $[0, 1] \times V_j \rightarrow V_j$. It has the property

$$(5.1) \quad d_G \mathbf{h}_j + \mathbf{h}_j d_G = \text{Id} - \pi_j^* \iota_j^*$$

where $\iota_j: \mathcal{C}_j \rightarrow V_j$ is the inclusion and $\pi_j: V_j = G \times_{G_j} S_j \rightarrow G/G_j = \mathcal{C}_j$ the projection. Let $(\varpi_j)_G = \mathbf{h}_j \eta_G - \pi_j^* \omega_{\mathcal{C}_j} \in \Omega_G^2(V_j)$, and write $(\varpi_j)_G = \varpi_j - \Psi_j$ where $\varpi_j \in \Omega^2(V_j)$ and $\Psi_j \in \Omega^0(V_j, \mathfrak{g})$.

PROPOSITION 5.2. *The equivariant 2-form $(\varpi_j)_G = \varpi_j - \Psi_j$ has the following properties.*

- (a) $d_G(\varpi_j)_G = \eta_G$.
- (b) *The pull-back of $(\varpi_j)_G$ to a conjugacy class $\mathcal{C} \subset V_j$ is given by*

$$\iota_{\mathcal{C}}^*(\varpi_j)_G = \Psi_j^*(\omega_{\mathcal{O}})_G - \omega_{\mathcal{C}},$$

where $(\omega_{\mathcal{O}})_G$ is the equivariant symplectic form on the adjoint orbit $\mathcal{O} = \Psi_j(\mathcal{C})$,

- (c) *The pull-back of Ψ_j to the conjugacy class \mathcal{C}_j vanishes. In fact, $\Psi_j(\exp \xi) = \xi - \mu_j$ for all $\xi \in \mathfrak{A}_j$.*
- (d) *Over each intersection $V_{ij} = V_i \cap V_j$, the difference $\Psi_i - \Psi_j$ takes values in the adjoint orbit \mathcal{O}_{ij} through $\mu_j - \mu_i \in \mathfrak{g} \cong \mathfrak{g}^*$. Furthermore,*

$$(\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G$$

where $p_{ij}: V_{ij} \rightarrow \mathcal{O}_{ij}$ is the map defined by $\Psi_i - \Psi_j$, and $(\omega_{\mathcal{O}_{ij}})_G$ is the equivariant symplectic form on the orbit.

Proof. (a) holds by construction. (b) follows from the observation that $\iota_{\mathcal{C}}^*(\varpi_j)_G + \omega_{\mathcal{C}}$ is an equivariantly closed 2-form on \mathcal{C}_j , with Ψ_j as its moment map. To prove (c) we note that since the retraction is equivariant, we have $\tilde{\mathbf{h}}_j \circ (\exp|_{\mathfrak{A}_j})^* = (\exp|_{\mathfrak{A}_j})^* \circ \mathbf{h}_j$ where $(\exp|_{\mathfrak{A}_j})^*$ is pull-back to $\mathfrak{A}_j \subset \mathfrak{t}$ and where $\tilde{\mathbf{h}}_j$ is the homotopy operator for the linear retraction of \mathfrak{t} onto $\{\mu_j\}$. Let $\nu: \mathfrak{A}_j \rightarrow \mathfrak{t}$ be the coordinate function (inclusion). Then

$$\tilde{\mathbf{h}}_j \circ (\exp|_{\mathfrak{A}_j})^* \frac{1}{2}(\theta^L + \theta^R) = \tilde{\mathbf{h}}_j \circ d\nu = \nu - \mu_j,$$

proving that $(\exp|_{\mathfrak{A}_j})^* \Psi_j = \nu - \mu_j$. This yields (c), by equivariance. For $\nu \in \mathfrak{A}_{ij}$ we have, using (c),

$$(\Psi_i - \Psi_j)(\exp \nu) = (\nu - \mu_i) - (\nu - \mu_j) = \mu_j - \mu_i.$$

By equivariance, it follows that $\Psi_i - \Psi_j$ takes values in the adjoint orbit through $\mu_j - \mu_i$. The difference $\varpi_i - \varpi_j$ vanishes on the maximal torus T , and is therefore determined by its contractions with generating vector fields. Since $\Psi_i - \Psi_j$ is a moment map for $\varpi_i - \varpi_j$, it follows that $\varpi_i - \varpi_j$ equals the pull-back of the symplectic form on $G \cdot (\mu_j - \mu_i)$.

5.3 THE SPECIAL UNITARY GROUP

For the special unitary group $G = \mathrm{SU}(d+1)$, the construction of the basic gerbe simplifies due to the fact that in this case all vertices μ_j of the alcove are contained in the weight lattice. In fact the gerbe is presented as a Chatterjee-Hitchin gerbe for the cover $\mathcal{V} = \{V_i, i = 0, \dots, d\}$.

For each weight $\mu \in \Lambda^* \subset \mathfrak{t} \subset \mathfrak{g}$, let G_μ be its stabilizer for the adjoint action and let \mathbf{C}_μ the 1-dimensional G_μ -representation with infinitesimal character μ . Let the line bundle $L_\mu = G \times_{G_\mu} \mathbf{C}_\mu$ equipped with the unique left-invariant connection ∇ . Then L_μ is a G -equivariant pre-quantum line bundle for the orbit $\mathcal{O} = G \cdot \mu$. That is,

$$\frac{i}{2\pi} \mathrm{curv}_G(\nabla) = (\omega_{\mathcal{O}})_G := \omega_{\mathcal{O}} - \Phi_{\mathcal{O}}$$

where $\omega_{\mathcal{O}}$ is the symplectic form and $\Phi_{\mathcal{O}}: \mathcal{O} \hookrightarrow \mathfrak{g}^*$ is the moment map given as inclusion.

In particular, in the case of $\mathrm{SU}(d+1)$ all orbits $\mathcal{O}_{ij} = G \cdot (\mu_j - \mu_i)$ carry G -equivariant pre-quantum line bundles. Recall the fibrations $p_{ij}: V_{ij} \rightarrow \mathcal{O}_{ij}$ defined by $\Psi_i - \Psi_j$, and let

$$L_{ij} = p_{ij}^*(L_{\mu_j - \mu_i}),$$

equipped with the pull-back connection. For any triple intersection $V_{ijk} = G \times_{G_{ijk}} S_{ijk}$, the tensor product $(\delta L)_{ijk} = L_{jk} L_{ik}^{-1} L_{ij}$ is the pull-back of the line bundle over G/G_{ijk} , defined by the zero weight

$$(\mu_k - \mu_j) - (\mu_k - \mu_i) + (\mu_j - \mu_i) = 0$$

of G_{ijk} . It is hence canonically trivial, with $(\delta \nabla)_{ijk}$ the trivial connection. The trivializing section $t_{ijk} = 1$ satisfies $\delta t = 1$ and $(\delta \nabla)t = 0$. Take $(B_j)_G = (\varpi_j)_G$. Then

$$(B_j)_G - (B_i)_G = (\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G = \frac{1}{2\pi i} \mathrm{curv}_G(\nabla^{L_{ij}}).$$

Thus $\mathcal{G} = (\mathcal{V}, L, t)$ is a equivariant gerbe with connection (∇, B) . Since