

# 1.4 FURTHER PROPERTIES OF $X_m$

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **05.06.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$W_z$ -invariant, we deduce that  $p(x) - p(sx) = 0$ , so that in this case  $p(x) - p(sx)$  also is divisible by  $\alpha_s(x)^{2m_s+1}$ .

To conclude, notice that  $p(z) \neq 0$ . Indeed, for a reflection  $s$ ,  $\alpha_s$  vanishes exactly on the fixed points of  $s$ , so that  $\prod_{s \in \Sigma, sz \neq z} \alpha_s(z)^{2m_s+1} \neq 0$ . Also for all  $w \in W_z$ ,  $f(wz) = f(z) \neq 0$ . On the other hand, it is clear that  $p(y) = 0$ .  $\square$

EXAMPLE 1.5. Take  $W = \mathbf{Z}/2$ . As we have already seen,  $Q_m$  has a basis given by the monomials  $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$ . From this we deduce that setting  $z = x^2$  and  $y = x^{2m+1}$ ,  $Q_m = \mathbf{C}[y, z]/(y^2 - z^{2m+1}) = \mathbf{C}[K]$ , where  $K$  is the plane curve with a cusp at the origin, given by the equation  $y^2 = z^{2m+1}$ . The map  $\pi: \mathbf{C} \rightarrow K$  is given by  $\pi(t) = (t^{2m+1}, t^2)$ , which is clearly bijective.

#### 1.4 FURTHER PROPERTIES OF $X_m$

Let us get to some deeper properties of quasi-invariants. Let  $X$  be an irreducible affine variety over  $\mathbf{C}$  and  $A = \mathbf{C}[X]$ . Recall that, by the Noether Normalization Lemma, there exist  $f_1, \dots, f_n \in \mathbf{C}[X]$  which are algebraically independent over  $\mathbf{C}$  and such that  $\mathbf{C}[X]$  is a finite module over the polynomial ring  $\mathbf{C}[f_1, \dots, f_n]$ . This means that we have a finite morphism of  $X$  onto an affine space.

DEFINITION 1.6.  $A$  (and  $X$ ) is said to be *Cohen-Macaulay* if there exist  $f_1, \dots, f_n$  as above, with the property that  $\mathbf{C}[X]$  is a locally free module over  $\mathbf{C}[f_1, \dots, f_n]$ . (Notice that by the Quillen-Suslin theorem, this is equivalent to saying that  $A$  is a free module.)

REMARK. If  $A$  is Cohen-Macaulay, then for any  $f_1, \dots, f_n$  which are algebraically independent over  $\mathbf{C}$  and such that  $A$  is a finite module over the polynomial ring  $\mathbf{C}[f_1, \dots, f_n]$ , we have that  $A$  is a locally free  $\mathbf{C}[f_1, \dots, f_n]$ -module, see [Eis], Corollary 18.17.

THEOREM 1.7 ([EG2], [BEG], conjectured in [FV]).  $Q_m$  is Cohen-Macaulay.

Notice that, using Chevalley's result that  $\mathbf{C}[\mathfrak{h}]^W$  is a polynomial ring, it will suffice, in order to prove Theorem 1.7, to prove:

THEOREM 1.8 ([EG2, BEG], conjectured in [FV]).  $Q_m$  is a free  $\mathbf{C}[\mathfrak{h}]^W$ -module.

We show how one can prove this Theorem in 3.10. This proof follows [BEG] (the original proof of [EG2] is shorter but somewhat less conceptual). The main idea of the proof is to show that the  $\mathbf{C}[\mathfrak{h}]^W$ -module  $Q_m$  can be extended to a module over a bigger (noncommutative) algebra, namely the spherical subalgebra of the rational Cherednik algebra. Furthermore, this module belongs to an appropriate category of representations of this algebra, called category  $\mathcal{O}$ . On the other hand, it can be shown that any module over the spherical subalgebra that belongs to this category is free when restricted to the commutative algebra  $\mathbf{C}[\mathfrak{h}]^W$ .

### 1.5 THE POINCARÉ SERIES OF $Q_m$

Consider now the Poincaré series

$$h_{Q_m}(t) = \sum_{r \geq 0} \dim Q_m[r] t^r,$$

where  $Q_m[r]$  denotes the graded component of  $Q_m$  of degree  $r$ . For every irreducible representation  $\tau \in \widehat{W}$ , define

$$\chi_\tau(t) = \sum_{r \geq 0} \dim \text{Hom}_W(\tau, \mathbf{C}[\mathfrak{h}][r]) t^r.$$

Consider the element in the group ring  $\mathbf{Z}[W]$

$$\mu_m = \sum_{s \in \Sigma} m_s (1 - s).$$

The  $W$ -invariance of  $m$  implies that  $\mu_m$  lies in the center of  $\mathbf{Z}[W]$ . Hence it is clear that  $\mu_m$  acts as a scalar,  $\xi_m(\tau)$ , on  $\tau$ . Let  $d_\tau$  be the degree of  $\tau$ .

**LEMMA 1.9.** *The scalar  $\xi_m(\tau)$  is an integer.*

*Proof.*  $\mathbf{Z}[W]$  and hence also its center, is a finite  $\mathbf{Z}$ -module. This clearly implies that  $\xi_m(\tau)$  is an algebraic integer. Thus to prove that  $\xi_m(\tau)$  is an integer, it suffices to see that  $\xi_m(\tau)$  is a rational number. Let  $d_{\tau,s}$  be the dimension of the space of  $s$ -invariants in  $\tau$ . Taking traces we get

$$d_\tau \xi_m(\tau) = \sum_{s \in \Sigma} 2m_s (d_\tau - d_{\tau,s}),$$

which gives the rationality of  $\xi_m(\tau)$ .  $\square$