

The Hopf conjecture and the Singer conjecture

Autor(en): **Davis, Michael W.**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109896>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

27

THE HOPF CONJECTURE AND THE SINGER CONJECTURE

by Michael W. DAVIS

CONJECTURE 27.1. *Suppose M^{2k} is a closed, aspherical manifold of dimension $2k$. Then $(-1)^k \chi(M^{2k}) \geq 0$.*

The conjecture is true in dimension 2 since the only surfaces which have positive Euler characteristic are S^2 and \mathbf{RP}^2 and they are the only two which are not aspherical. In the special case where M^{2k} is a nonpositively curved Riemannian manifold this conjecture is usually attributed to Hopf by topologists and either to Chern or to both Chern and Hopf by differential geometers.

When I first heard about this conjecture in 1981, I thought I could come up with a counterexample by using right-angled Coxeter groups. Given a finite simplicial complex L which is a flag complex, there is an associated right-angled Coxeter group W . Its Euler characteristic is given by the formula

$$(27.1) \quad \chi(W) = 1 + \sum_{i=0}^{\dim L} \left(-\frac{1}{2}\right)^{i+1} f_i,$$

where f_i denotes the number of i -simplices in L . If L is a triangulation of S^{n-1} , then W acts properly and cocompactly on a contractible n -manifold. The quotient of this contractible manifold by any finite index, torsion-free subgroup $\Gamma \subset W$ is a closed aspherical n -manifold M^n . Since $\chi(M^n)$ is a positive multiple of $\chi(W)$ (by $[W : \Gamma]$), they have the same sign. So, this looked like a good way to come up with counterexamples to Conjecture 27.1. Conversely, if you believe Conjecture 27.1, then you must also believe the following

CONJECTURE 27.2. *If L is any flag triangulation of S^{2k-1} then*

$$(-1)^k \kappa(L) \geq 0,$$

where $\kappa(L)$ is the quantity defined by the right-hand side of (27.1).

Ruth Charney and I published this conjecture in [2]. It is sometimes called the Charney–Davis Conjecture.

In the 1970's Atiyah [1] introduced L^2 methods into topology. If a discrete group Γ acts properly and cocompactly on a smooth manifold or a CW-complex Y , then one can define the reduced L^2 -cohomology spaces of Y and their “dimensions” with respect to Γ , the so-called “ L^2 -Betti numbers”. Let $L^2 b_i(Y; \Gamma)$ be the Γ -dimension of the L^2 -cohomology of Y in dimension i . It is a *nonnegative* real number. If $Y \rightarrow X$ is a regular covering of a finite CW-complex X with group of deck transformations Γ , the Euler characteristic of X can be calculated from the L^2 -Betti numbers of Y by the formula

$$(27.2) \quad \chi(X) = \sum (-1)^i L^2 b_i(Y; \Gamma).$$

Shortly after Atiyah described this formula in [1], Dodziuk [4] and Singer realized that there is a conjecture about L^2 -Betti numbers which is stronger than Conjecture 27.1. It is usually called the Singer Conjecture. Beno Eckmann [5] also discusses it in this volume.

CONJECTURE 27.3 ([4]). *Suppose M^n is a closed, aspherical manifold with fundamental group π and universal cover \widetilde{M}^n . Then $L^2 b_i(\widetilde{M}^n; \pi) = 0$ for all $i \neq \frac{n}{2}$. (In particular, when n is odd this means all its L^2 -Betti numbers vanish.)*

This implies Conjecture 27.1 since, when $n = 2k$, formula (27.2) gives: $(-1)^k \chi(M^{2k}) = L^2 b_k(\widetilde{M}^{2k}; \pi) \geq 0$.

Of course, there is also the following version of Conjecture 27.3 for Coxeter groups.

CONJECTURE 27.4. *Suppose that L is a triangulation of S^{n-1} as a flag complex, that W is the associated right-angled Coxeter group and that Σ is the contractible n -manifold on which W acts. Then $L^2 b_i(\Sigma; W) = 0$ for all $i \neq \frac{n}{2}$.*

Boris Okun and I discussed this conjecture in [3] and we proved it for $n \leq 4$. The result for $n = 4$ implies Conjecture 27.2 when L is a flag triangulation of S^3 . So, Conjecture 27.2 is true in the first dimension for which it is not obvious.

REFERENCES

- [1] ATIYAH, M. F. Elliptic operators, discrete groups and von Neumann algebras. In: *Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974)*, 43–72. Astérisque 32–33. Soc. Math. France, 1976.
- [2] CHARNEY, R. and M. W. DAVIS. The Euler characteristic of a nonpositively curved, piecewise Euclidean manifold. *Pacific J. Math.* 171 (1995), 117–137.
- [3] DAVIS, M. W. and B. OKUN. Vanishing theorems and conjectures for the L^2 -homology of right-angled Coxeter groups. *Geom. Topol.* 5 (2001), 7–74.
- [4] DODZIUK, J. L^2 harmonic forms on rotationally symmetric Riemannian manifolds. *Proc. Amer. Math. Soc.* 77 (1979), 395–400.
- [5] ECKMANN, B. The Singer conjecture. (*This volume.*)

M. Davis

The Ohio-State University
231 West 18th Avenue
43210 Columbus, OH
USA
e-mail: mdavis@math.ohio-state.edu