

Alperin's weight conjecture

Autor(en): **Kessar, Radha / Linckelmann, Markus**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109911>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

42

ALPERIN'S WEIGHT CONJECTURE

by Radha KESSAR and Markus LINCKELMANN

Let p be a prime number.

CONJECTURE 42.1. *Let G be a finite group and let P be a Sylow p -subgroup of G .*

- (i) *The number of conjugacy classes of p' -elements of G is greater than or equal to the number of conjugacy classes of $N_G(P)/P$.*
- (ii) *If P is abelian, then the number of conjugacy classes of G is greater than or equal to the number of conjugacy classes of $N_G(P)$.*

The above inequalities would follow from Alperin's weight conjecture [1] which we describe now.

Let k be an algebraically closed field of characteristic p . For a finite group H denote by $l(kH)$ the number of isomorphism classes of simple kH -modules, and by $w(kH)$ the number of isomorphism classes of simple projective kH -modules. The weight conjecture predicts the following

CONJECTURE 42.2. *Let G be a finite group. Then*

$$l(kG) = \sum_{Q \in \mathcal{I}} w(kN_G(Q)/Q),$$

where \mathcal{I} denotes a set of representatives of G -conjugacy classes of p -subgroups of G .

Conjecture 42.2 comes in a block-wise version as well. The reformulation of this conjecture in terms of alternating chains in [13] paved the way for many extensions (see for instance [9], [10], [11], [16]). Despite having been verified for many families of finite groups, including finite p -solvable groups,

symmetric groups, and in some cases for finite groups of Lie type and some sporadic simple groups (see for instance [2], [3], [4], [5], [6], [8], [12]), a true understanding of Conjecture 42.2 or indeed of Conjecture 42.1 remains elusive.

In its original form stated above, Alperin's weight conjecture is a numerical equality interpreting the number of simple modules of a finite group or a p -block in terms of the involved p -local structure. In subsequent years structural approaches to this and related conjectures in terms of linear source modules [7], fusion category algebras [14], and cohomological invariants of functors over certain finite categories [15], have emerged.

We briefly explain how 42.1 would follow from 42.2. By a theorem of Brauer, $l(kG)$ is equal to the number of conjugacy classes of p' -elements in G , and the summand for $Q = P$ on the right side of the equality 42.2 is equal to the number of conjugacy classes of $N_G(P)/P$. Thus 42.2 implies the inequality 42.1 (i). The inequality 42.1 (ii) follows from 42.1 (i) applied to centralizers of p -elements.

REFERENCES

- [1] ALPERIN, J. L. Weights for finite groups. In: *The Arcata Conference on Representations of Finite Groups*, 369–379. Proc. Sympos. Pure Math. 47. Amer. Math. Soc., Providence, R.I., 1987.
- [2] ALPERIN, J. L. and P. FONG. Weights for symmetric and general linear groups. *J. Algebra* 131 (1990), 2–22.
- [3] AN, J. 2-weights for general linear groups. *J. Algebra* 149 (1992), 500–527.
- [4] —— Weights for classical groups. *Trans. Amer. Math. Soc.* 342 (1994), 1–42.
- [5] —— The Alperin and Uno conjectures for the Fischer simple group Fi_{22} . *Comm. Algebra* 33 (2005), 1529–1557.
- [6] AN, J. and R.A. WILSON. The Alperin weight conjecture and Uno's conjecture for the Baby Monster \mathbf{B} , p odd. *London Math. Soc. J. Comput. Math.* 7 (2004), 120–166.
- [7] BOLTJE, R. Alperin's weight conjecture in terms of linear source modules and trivial source modules. In: *Modular Representation Theory of Finite Groups* (Charlottesville, VA, 1998), 147–155. Walter de Gruyter, Berlin, 2001.
- [8] CABANES, M. and M. ENGUEHARD. On fusion in unipotent blocks. *Bull. London Math. Soc.* 31 (1999), 143–148.
- [9] DADE, E.C. Counting characters in blocks, I. *Invent. Math.* 109 (1992), 187–210.
- [10] —— Counting characters in blocks, II. *J. Reine Angew. Math.* 448 (1994), 97–190.

- [11] —— Counting characters in blocks, II.9. In: *Representation Theory of Finite Groups*, 45–59. Ohio State Univ. Math. Res. Inst. Publ. 6. Walter de Gruyter, Berlin, 1997.
- [12] ISAACS, I.M. and G. NAVARRO. Weights and vertices for characters of π -separable groups. *J. Algebra* 177 (1995), 339–366.
- [13] KNÖRR R. and G.R. ROBINSON. Some remarks on a conjecture of Alperin. *J. London Math. Soc.* (2) 39 (1989), 48–60.
- [14] LINCKELMANN, M. Fusion category algebras. *J. Algebra* 277 (2004), 222–235.
- [15] —— Alperin’s weight conjecture in terms of equivariant Bredon cohomology. *Math. Z.* 250 (2005), 495–513.
- [16] ROBINSON, G.R. Local structure, vertices and Alperin’s conjecture. *Proc. London Math. Soc.* (3) 72 (1996), 312–330.

Radha Kessar, Markus Linckelmann

Department of Mathematical Sciences
University of Aberdeen
Meston Building
Aberdeen AB24 3UE
United Kingdom
e-mail: kessar@maths.abdn.ac.uk, m.linckelmann@maths.abdn.ac.uk