

On algebraic characterizations for the finiteness of the dimension of EG

Autor(en): **Talelli, Olympia**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **54 (2008)**

Heft 1-2

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-109930>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

61

ON ALGEBRAIC CHARACTERIZATIONS FOR THE FINITENESS OF THE DIMENSION OF $\underline{E}G$

by Olympia TALELLI

In [5] the following theorem was proved :

THEOREM 61.1. *If G is an $\text{H}\mathfrak{F}$ -group of type FP_∞ then G admits a finite dimensional model for $\underline{E}G$.*

The class $\text{H}\mathfrak{F}$ was introduced by P. H. Kropholler in [4] and it is defined as the smallest class of groups containing the class of finite groups, with the property : if a group G admits a finite dimensional contractible G -CW-complex with all cell stabilizers in $\text{H}\mathfrak{F}$ then G is in $\text{H}\mathfrak{F}$.

This theorem, especially its proof, was the motivation for defining groups of type Φ in [7] and for proposing those as the ones which admit a finite dimensional model for $\underline{E}G$.

DEFINITION 61.2 ([7]). A group G is said to be of type Φ if it has the property that for every $\mathbf{Z}G$ -module M , $\text{projdim}_{\mathbf{Z}G} M < \infty$ if and only if $\text{projdim}_{\mathbf{Z}H} M < \infty$ for every finite subgroup H of G .

CONJECTURE 61.3. *The following statements are equivalent for a group G :*

- (1) *G admits a finite dimensional model for $\underline{E}G$.*
- (2) *G admits a finite dimensional contractible G -CW-complex with finite cell stabilizers.*
- (3) *G is of type Φ .*
- (4) *$\text{spli } \mathbf{Z}G < \infty$.*
- (5) *$\text{silp } \mathbf{Z}G < \infty$.*
- (6) *$\text{findim } \mathbf{Z}G < \infty$.*

The algebraic invariants $\text{spli } \mathbf{Z}G$ and $\text{silp } \mathbf{Z}G$ were defined in [3]: $\text{silp } \mathbf{Z}G$ is the supremum of the injective lengths of the projective $\mathbf{Z}G$ -modules and $\text{spli } \mathbf{Z}G$ is the supremum of the projective lengths of the injective $\mathbf{Z}G$ -modules. It was shown in [3] that $\text{silp } \mathbf{Z}G \leq \text{spli } \mathbf{Z}G$, and that if $\text{spli } \mathbf{Z}G < \infty$ then $\text{spli } \mathbf{Z}G = \text{silp } \mathbf{Z}G$. The *finitistic dimension* of $\mathbf{Z}G$, $\text{findim } \mathbf{Z}G$, is the supremum of the projective dimensions of the $\mathbf{Z}G$ -modules of finite projective dimension.

It is not very difficult to show that $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (6)$, see [7].

If G is an HF -group of type FP_∞ then there is a bound on the orders of the finite subgroups of G [4], and $\text{projdim}_{\mathbf{Z}G} \mathbf{B}(G, \mathbf{Z})$ is finite, where $\mathbf{B}(G, \mathbf{Z})$ is the $\mathbf{Z}G$ -module of the bounded functions from G to \mathbf{Z} [1].

Theorem 61.1 is now an immediate consequence of the following theorem which was proved in [5]:

THEOREM 61.4. *If G is an HF -group such that there is a bound on the orders of the finite subgroups and $\text{projdim}_{\mathbf{Z}G} \mathbf{B}(G, \mathbf{Z})$ is finite, then G admits a finite dimensional model for $\underline{\mathbf{E}}G$.*

Since if G is an HF -group, $\text{projdim}_{\mathbf{Z}G} \mathbf{B}(G, \mathbf{Z})$ is finite if and only if $\text{findim } \mathbf{Z}G$ is finite [1], it follows from Theorem 61.4 that $(6) \Rightarrow (1)$ in Conjecture 61.3 if G is an HF -group with a bound on the orders of the finite subgroups. Moreover in [8] it is shown that $(6) \Rightarrow (1)$ if G is a torsion-free elementary amenable group.

In [6] it was shown that if a group G admits a finite dimensional model for $\underline{\mathbf{E}}G$ then for every finite subgroup H of G , $W(H)$ admits a finite dimensional model for $\underline{\mathbf{E}}W(H)$, where $W(H) = N_G(H)/H$. In [7] we show that if $\text{spli } \mathbf{Z}G$ is finite, then $\text{spli } \mathbf{Z}W(H)$ is finite for every finite subgroup H of G .

In support of Conjecture 61.3 is also the following characterization of finite groups, which we obtained in [2]: a group G is finite if and only if $\text{spli } \mathbf{Z}G = 1$.

REFERENCES

- [1] CORNICK, J. and P.H. KROPHOLLER. Homological finiteness conditions for modules over group algebras. *J. London Math. Soc.* (2) 58 (1998), 49–62.
- [2] DEMBEGIOTI, F. and O. TALELLI. An integral homological characterization of finite groups. *J. Algebra* 319 (2008), 267–271.

- [3] GEDRICH, T. V. and K. W. GRUENBERG. Complete cohomological functors on groups. *Topology Appl.* 25 (1987), 203–223.
- [4] KROPHOLLER, P. H. On groups of type FP_∞ . *J. Pure Appl. Algebra* 90 (1993), 55–67.
- [5] KROPHOLLER, P. H. and G. MISLIN. Groups acting on finite dimensional spaces with finite stabilizers. *Comment. Math. Helv.* 73 (1998), 122–136.
- [6] LÜCK, W. The type of the classifying space for a family of subgroups. *J. Pure Appl. Algebra* 149 (2000), 177–203.
- [7] TALELLI, O. On groups of type Φ . *Archiv der Math.* 89 (2007), 24–32.
- [8] TALELLI, O. A characterization for cohomological dimension for a big class of groups. (Preprint, 2007.)

Olympia Talelli

Department of Mathematics
University of Athens
Athens 15784
Greece
e-mail: otalelli@math.uoa.gr