Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	65 (2019)
Heft:	1-2
Artikel:	Nonhyperbolic Coxeter groups with Menger boundary
Autor:	Haulmark, Matthew / Hruska, G. Christopher / Sathaye, Bakul
DOI:	https://doi.org/10.5169/seals-869349

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 12.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Nonhyperbolic Coxeter groups with Menger boundary

Matthew HAULMARK, G. Christopher HRUSKA and Bakul SATHAYE

Abstract. A generic finite presentation defines a word hyperbolic group whose boundary is homeomorphic to the Menger curve. In this article we produce the first known examples of non-hyperbolic CAT(0) groups whose visual boundary is homeomorphic to the Menger curve. The examples in question are the Coxeter groups whose nerve is a complete graph on *n* vertices for  $n \ge 5$ . The construction depends on a slight extension of Sierpiński's theorem on embedding 1-dimensional planar compacta into the Sierpiński carpet. We give a simplified proof of this theorem using the Baire category theorem.

Mathematics Subject Classification (2010). Primary: 20F67; Secondary: 20F55, 54F50.

Keywords. Nonpositive curvature, Menger curve, Coxeter group.

#### 1. Introduction

Many word hyperbolic groups have Gromov boundary homeomorphic to the Menger curve. Indeed random groups have Menger boundary with overwhelming probability [Cha, DGP]. Therefore, in a strong sense, Menger boundaries are ubiquitous among hyperbolic groups. This phenomenon depends heavily on the fact that the boundary of a one-ended hyperbolic group is always locally connected, a necessary condition since the Menger curve is a locally connected compactum.

However in the broader setting of CAT(0) groups, the visual boundary often fails to be locally connected, especially in the case when the boundary is onedimensional. For instance the direct product  $F_2 \times \mathbb{Z}$  of a free group with the integers has boundary homeomorphic to the suspension of the Cantor set, which is one-dimensional but not locally connected. The CAT(0) groups with isolated flats are, in many ways, similar to hyperbolic groups, and are often viewed as the simplest nontrivial generalization of hyperbolicity. However, even in that setting many visual boundaries are not locally connected. For example if X is formed by gluing a closed hyperbolic surface to a torus along a simple closed geodesic loop, then its fundamental group G is a CAT(0) group with isolated flats that has non-locally connected boundary [MR].

One might wonder whether the Menger curve boundary is a unique feature of the hyperbolic setting. Indeed, recently Kim Ruane observed that not a single example was known of a nonhyperbolic CAT(0) group with a visual boundary homeomorphic to the Menger curve, posing the following question.

**Question 1.1** (Ruane). Does there exist a nonhyperbolic group G acting properly, cocompactly, and isometrically on a CAT(0) space X such that the visual boundary of X is homeomorphic to the Menger curve?

In this article we provide the first explicit examples of nonhyperbolic CAT(0) groups with Menger visual boundary.

**Theorem 1.2.** Let W be the Coxeter group defined by a presentation with n generators of order two such that the order  $m_{st}$  of st satisfies  $3 \le m_{st} < \infty$  for all generators  $s \ne t$  (or more generally let W be any Coxeter group whose nerve is 1-dimensional and equal to the complete graph  $K_n$ ).

- (1) If n = 3 the group W has visual boundary homeomorphic to the circle and acts as a reflection group on the Euclidean or hyperbolic plane.
- (2) If n = 4 the group W has visual boundary homeomorphic to the Sierpiński carpet and acts as a reflection group on a convex subset of  $\mathbb{H}^3$  with fundamental chamber a (possibly ideal) convex polytope.
- (3) For each  $n \ge 5$ , the group W has visual boundary homeomorphic to the Menger curve.

The nerve of a Coxeter system is defined in Definition 3.1. The nerve is 1-dimensional when all three-generator special subgroups are finite (see Remark 3.3). We note that a 1-dimensional nerve L is a complete graph if and only if every  $m_{st}$  is finite.

The proof of this theorem depends on work of Hruska–Ruane determining which CAT(0) groups with isolated flats have locally connected visual boundary [HR1] and subsequent work of Haulmark on the existence of local cut points in boundaries [Hau]. In particular, [Hau] gives a criterion that ensures the visual boundary of a CAT(0) group with isolated flats will be either the circle, the Sierpiński carpet, or the Menger curve (extending a theorem of Kapovich–Kleiner from the word hyperbolic setting [KK]). The circle occurs only for virtual surface groups. In order to distinguish between the other two possible boundaries, one needs to determine whether the boundary is planar. In general the nerve of a Coxeter group does not have an obvious natural embedding into the visual

boundary. However we show in this article that Coxeter groups with nerve  $K_n$  do admit an embedding of  $K_5$  in the boundary and hence have non-planar boundary whenever  $n \ge 5$ .

By Bestvina-Kapovich-Kleiner [BKK], we get the following corollary.

**Corollary 1.3.** Let W be a Coxeter group with at least 5 generators such that every  $m_{st}$  satisfies  $3 \le m_{st} < \infty$ . Then W acts properly on a contractible 4manifold but does not admit a coarse embedding into any contractible 3-manifold. In particular, W is not virtually the fundamental group of any 3-manifold.

**1.1. Related problems and open questions.** A word hyperbolic special case of Theorem 1.2 (when all  $m_{st}$  are equal and are strictly greater than 3) is due to Benakli [Ben]. Related results of Bestvina–Mess, Champetier, and Bonk–Kleiner [BM, Cha, BK] provide various methods for constructing embedded arcs and graphs in boundaries of hyperbolic groups.

In principle, any of the well-known hyperbolic techniques could be expected to generalize to some families of CAT(0) spaces with isolated flats, although the details of such extensions would necessarily be more subtle than in the hyperbolic case. For example, as mentioned above many groups with isolated flats have nonlocally connected boundary, and thus are not linearly connected with respect to any metric.

We note that the proof of Theorem 1.2 given here is substantially different from the methods used by Benakli in the hyperbolic setting. The proof here is quite short and uses a slight extension of Sierpiński's classical embedding theorem to produce arcs in the boundary. (We provide a new proof of this embedding theorem.) Unlike in the hyperbolic setting, these arcs do not arise as boundaries of quasi-isometrically embedded hyperbolic planes.

Nevertheless it seems likely that many of the hyperbolic techniques mentioned above could also be extended to the present setting, which suggests the following natural questions.

**Question 1.4.** What conditions on the nerve of a Coxeter group W are sufficient to ensure that the open cone on the nerve L admits a proper, Lipschitz, expanding map into the Davis–Moussong complex of W? When does the nerve L embed in the visual boundary?

**Question 1.5.** Let *G* be a one-ended CAT(0) group with isolated flats. Does the visual boundary of *G* have the doubling property? If the boundary is locally connected, is it linearly connected? Note that the usual visual metrics on Gromov boundaries do not exist in the CAT(0) setting, so a different metric must be used – such as the metric studied in [OS].

**Question 1.6.** Let W be the family of all nonhyperbolic Coxeter groups W with nerve a complete graph  $K_n$  where  $n \ge 5$  and all labels  $m_{st} = 3$ . Are all groups in W quasi-isometric? Can conformal dimension be used to distinguish quasi-isometry classes of groups in W? As above, one would need to select an appropriate metric on the boundary in order to make this question more precise.

It is known that vanishing of the  $\ell^2$ -Betti number in dimension *i* is a quasiisometry invariant for each *i* [Gro, Pan]. Mogilski has computed the  $\ell^2$ -Betti numbers of the groups mentioned in the previous question: for each  $W \in W$ they are nontrivial in dimension two and vanish in all other dimensions [Mog, Cor 5.7]. However, this computation does not give any information about the quasi-isometry classification of the family W. Thus different techniques would be needed to address Question 1.6.

# 2. Arcs in the Sierpiński carpet

In 1916, Sierpiński proved that every planar compactum of dimension at most one embeds in the Sierpiński carpet [Sie]. The main result of this section is Proposition 2.3 - a slight extension of Sierpiński's theorem – which establishes the existence of embedded graphs in the Sierpiński carpet that connect an arbitrary finite collection of points lying on peripheral circles.

Although Sierpiński's proof of the embedding theorem was rather elaborate, we present here a simplified proof using the Baire Category Theorem. The general technique of applying the Baire Category Theorem to function spaces in order to prove embedding theorems is well-known in dimension theory and appears to originate in work of Hurewicz from the 1930s. The conclusion of Proposition 2.3 may not be surprising to experts, but we have provided the proof for the benefit of the reader.

We begin our discussion with a brief review of Whyburn's topological characterization of planar embeddings of the Sierpiński carpet.

**Definition 2.1** (Null family of subspaces). Let M be a compact metric space. A collection  $\mathcal{A}$  of subspaces of M is a *null family* if for each  $\epsilon > 0$  only finitely many members of  $\mathcal{A}$  have diameter greater than  $\epsilon$ . If  $\mathcal{A}$  is a null family of closed, pairwise disjoint subspaces, the quotient map  $\pi: M \to M/\mathcal{A}$ , which collapses each member of  $\mathcal{A}$  to a point, is *upper semicontinuous* in the sense that  $\pi$  is a closed map (see for example Proposition I.2.3 of [Dave]). **Remark 2.2** (Planar Sierpiński carpets). A Jordan region in the sphere  $S^2$  is a closed disc bounded by a Jordan curve. By a theorem of Whyburn [Why], a subset  $S \subset S^2$  is homeomorphic to the Sierpiński carpet if and only if it can be expressed as  $S = S^2 - \bigcup \operatorname{int}(D_i)$  for some null family of pairwise disjoint Jordan regions  $\{D_1, D_2, \ldots\}$  such that  $\bigcup D_i$  is dense in  $S^2$ . A peripheral circle of S is an embedded circle whose removal does not disconnect S. Equivalently the peripheral circles in a planar Sierpiński carpet  $S \subset S^2$  are precisely the boundaries of the Jordan regions  $D_i$ . We will denote the collection of peripheral circles in S by  $\mathcal{P}$ .

Let  $E_k$  be the k-pointed star, i.e., the cone on a set of k points. Let  $e_1, \ldots, e_k$  be the edges of  $E_k$ , which we will think of as embeddings of [0, 1] into E parametrized such that  $e_i(0) = e_j(0)$  for all  $i, j \in \{1, \ldots, k\}$ .

**Proposition 2.3.** Let  $P_1, \ldots, P_k \in \mathcal{P}$  be distinct peripheral circles in the Sierpiński carpet S, and fix points  $p_i \in P_i$ . There is a topological embedding  $h: E_k \hookrightarrow S$  such that  $h \circ e_i(1) = p_i$  for each  $i \in \{1, \ldots, k\}$ . Furthermore the image of  $E_k$  intersects the union of all peripheral circles precisely in the given points  $p_1, \ldots, p_k$ .

*Proof.* Let Q be the quotient space  $S/\sim$  formed by collapsing each peripheral circle  $P \in \mathcal{P} - \{P_1, \ldots, P_k\}$  to a point. Our strategy is to first show that Q is an orientable, genus zero surface with k boundary curves. Then we apply the Baire Category Theorem and the fact that  $E_k$  is 1-dimensional to find embeddings of  $E_k$  that avoid the countably many peripheral points of Q. The conclusion of the Proposition is illustrated in Figure 1.

We first check that Q is a surface. Fix an embedding  $S \hookrightarrow S^2$  as in Remark 2.2. We may form Q from  $S^2$  in two steps as follows. First collapse each peripheral Jordan region to a point except for those bounded by the curves  $P_1, \ldots, P_k$ . By a theorem of R.L. Moore [Moo], this upper semicontinuous quotient of  $S^2$  is again homeomorphic to  $S^2$ . (In particular, the quotient is Hausdorff.) The space Q may be recovered from this quotient by removing the interiors of the regions bounded by  $P_1, \ldots, P_k$ . Therefore Q may be obtained from a 2-sphere by removing the interiors of k pairwise disjoint Jordan regions.

Let  $\pi: S \to Q$  be the associated quotient map. By a slight abuse of notation we let  $P_i \subset Q$  and  $p_i \in Q$  denote  $\pi(P_i)$  and  $\pi(p_i)$ . A *peripheral point* of Q is the image of a peripheral circle  $P \in \mathcal{P} - \{P_1, \ldots, P_k\}$ . Observe that the peripheral points are a countable dense set in Q.

Let  $\mathcal{E}$  be the space of all embeddings  $\iota: E_k \hookrightarrow Q$  such that for each *i* we have  $\iota \circ e_i(1) = p_i$  and the image of  $\iota$  intersects  $P_1 \cup \cdots \cup P_k$  only in the *k* points  $p_1, \ldots, p_k$ . We fix a metric  $\rho$  on Q and equip  $\mathcal{E}$  with the complete



FIGURE 1 An embedding of the graph  $E_5$  that intersects the union of all peripheral circles precisely in the given points  $p_1, \ldots, p_5$ 

metric given by  $d(f,g) = \sup \{ \rho(f(x), g(x)) \mid x \in E_k \}$ . Our strategy is to show that for each peripheral point p, the set of embeddings avoiding p is open and dense in  $\mathcal{E}$ . It then follows by the Baire Category Theorem that there exists an embedding  $\iota \in \mathcal{E}$  whose image contains no peripheral points.

Toward this end, we fix an arbitrary peripheral point  $p \in Q$ . Since  $E_k$  is compact, the set of embeddings avoiding p is open, so we only need to prove that it is dense. Suppose  $f \in \mathcal{E}$  and p is in the image of f. Let  $\epsilon$  be any positive number small enough that the ball  $B(p,\epsilon)$  lies in the interior of Q. Since f is a homeomorphism onto its image, the image  $f(E_k)$  is 1-dimensional and thus does not contain any 2-dimensional disc. In particular, there is at least one point  $q \in B(p,\epsilon)$  not in the image  $f(E_k)$ . Apply an isotopy  $\Phi_t$  to Qkeeping  $Q - B(p,\epsilon)$  fixed and such that  $\Phi_1(q) = p$ . Then  $\Phi_1 \circ f : E_k \hookrightarrow Q$ is an element of  $\mathcal{E}$  which misses p. Furthermore its distance from f is less than  $2\epsilon$ . Since  $\epsilon$  may be chosen arbitrarily small, we conclude that the set of embeddings avoiding p is open and dense in  $\mathcal{E}$ .

Since the quotient map  $\pi: S \to Q$  is one-to-one on the complement of the peripheral circles, we may lift any embedding  $f \in \mathcal{E}$  that avoids peripheral points to an embedding  $E_k \hookrightarrow S$  satisfying the conclusion of the proposition. Indeed by compactness,  $f(E_k)$  is closed in Q, so its preimage  $\pi^{-1}f(E_k)$  in S is closed. The restriction of  $\pi$  to this compact preimage is a continuous bijection onto the Hausdorff space  $f(E_k)$ , so there is a continuous inverse function  $\pi^{-1}$  defined on  $f(E_k)$ . The composition  $\pi^{-1}f$  is the desired lift.

#### 3. Coxeter groups and the Davis–Moussong complex

Let  $\Upsilon$  be a finite simplicial graph with vertex set S whose edges are labeled by integers  $\geq 2$ . Let  $m_{st}$  denote the label on the edge  $\{s,t\}$ . If s and t are distinct vertices not joined by an edge, we let  $m_{st} = \infty$ . The Coxeter group determined by  $\Upsilon$  is the group

 $W = \langle S \mid s^2, (st)^{m_{st}}$  for all s, t distinct elements of  $S \rangle$ .

A Coxeter system (W, S) is a Coxeter group W with generating set S as above.

**Definition 3.1.** The *nerve* of a Coxeter system (W, S) is a metric simplicial complex with a 0-simplex for each generator  $s \in S$  and a higher simplex for each subset  $T \subseteq S$  such that T generates a finite subgroup of W.

If (W, S) is any Coxeter system, the Coxeter group W acts properly, cocompactly, and isometrically on the associated *Davis–Moussong complex*  $\Sigma(W, S)$ , a piecewise Euclidean CAT(0) complex such that the link L of each vertex is equal to the nerve of (W, S) [Dav1, Mou, Dav2].

We state here a result regarding limit sets of special subgroups. The first part is a folklore result (see, for example, Świątkowski [Świ]). The second part holds for all convex subgroups of CAT(0) groups (see, for instance, Swenson [Swe]).

**Proposition 3.2.** Let (W, S) be any Coxeter system and let  $W_T$  denote the special subgroup of W generated by a subset  $T \subset S$ .

- (1) The Davis-Moussong complex  $\Sigma(W_T, T)$  is a convex subspace of  $\Sigma(W, S)$ whose limit set  $\Lambda \Sigma(W_T, T)$  is naturally homeomorphic to the visual boundary of  $\Sigma(W_T, T)$ .
- (2) For any two subsets T and T' of S, we have

$$\Lambda\Sigma(W_T,T) \cap \Lambda\Sigma(W_{T'},T') = \Lambda\Sigma(W_{T\cap T'},T\cap T').$$

**Remark 3.3** (1-dimensional nerves). A Coxeter group has a 1-dimensional nerve L if and only if L does not contain a 2-simplex. A set of three generators  $\{r, s, t\}$  bounds a 2-simplex in L precisely when it generates a finite subgroup, i.e., when  $1/m_{rs} + 1/m_{st} + 1/m_{rt} > 1$ . (We follow the usual convention regarding  $\infty$  by considering  $1/m_{st}$  to equal zero when  $m_{st} = \infty$ .) Therefore the nerve L is 1-dimensional if for each triangle, the sum above is  $\leq 1$ . For example a Coxeter group has *large type* if all  $m_{st}$  satisfy  $3 \leq m_{st} \leq \infty$ . Evidently all large type Coxeter groups have 1-dimensional nerve.

In the 1-dimensional case, the nerve L is equal to the graph  $\Upsilon$ , the Davis– Moussong complex is 2-dimensional, each face is isometric to a regular Euclidean  $(2m_{st})$ -sided polygon, and the nerve L has a natural angular metric in which each edge  $\{s,t\}$  has length  $\pi - (\pi/m_{st})$ . We refer the reader to [Dav2] for more background on Coxeter groups from the CAT(0) point of view.

Coxeter groups of large type always have isolated flats – even when the nerve is not complete – by an observation of Wise (see [Hru] for details). The following analogous result for Coxeter groups with nerve a complete graph follows immediately from Corollary D of [Cap], since two adjacent edges in the nerve cannot both have label 2.

**Proposition 3.4.** Coxeter groups whose nerve is a complete graph always have isolated flats.

By Hruska-Kleiner [HK], the groups acting geometrically on CAT(0) spaces with isolated flats have a well-defined boundary in the following sense: If G acts geometrically on two CAT(0) spaces X and Y with isolated flats, then there exists a G-equivariant homeomorphism between their visual boundaries  $\partial X$  and  $\partial Y$ . This common boundary will be denoted  $\partial G$ .

**Proposition 3.5.** Let W be a Coxeter group whose nerve is a complete graph  $K_n$  with  $n \ge 3$ . The boundary  $\partial W$  of W is homeomorphic to either the circle, the Sierpiński carpet, or the Menger curve.

*Proof.* A theorem due to Serre [Ser, §I.6.5] states that if G is generated by a finite number of elements  $s_1, \ldots, s_n$  such that each  $s_i$  and each product  $s_i s_j$  has finite order, then G has Serre's Property FA. In other words, every action of G on a simplicial tree has a global fixed point. Evidently W satisfies Serre's criterion, and hence W does not split as a nontrivial graph of groups.

Since W acts geometrically on a 2-dimensional CAT(0) space, its boundary has dimension at most 1 by [Bes]. As W is infinite and not virtually free the dimension of the boundary must be exactly 1, provided that  $n \ge 3$ .

The first author proves in [Hau] that a CAT(0) group with isolated flats with 1-dimensional boundary that does not split over a virtually cyclic subgroup must have visual boundary homeomorphic to either the circle, the Sierpiński carpet, or the Menger curve.

Infinite Coxeter triangle groups always act as reflection groups on either the Euclidean plane or the hyperbolic plane. In particular they have circle boundary. The following proposition examines the case of Coxeter groups with nerve  $K_4$ .

**Proposition 3.6.** Let W be a Coxeter group whose nerve L is a complete graph  $K_4$  on 4 vertices. Then the boundary  $\partial W$  of W is homeomorphic to the Sierpiński carpet, and the limit set of each three generator special subgroup of W is a peripheral circle.

*Proof.* The nerve L of W is planar, so W embeds as a special subgroup of a Coxeter group with visual boundary  $S^2$  by a well-known doubling construction. (See, for example, [DO].) Indeed, one embeds L into  $S^2$ , and then fills each complementary region in the sphere with 2-simplices by adding a vertex in the interior of the region and coning off the boundary of the region to the new vertex. Each such cone is "right-angled" in the sense that each added edge  $\{s,t\}$  is assigned the label  $m_{st} = 2$ . This procedure produces a metric flag triangulation  $\hat{L}$  of  $S^2$ , which has L as a full subcomplex. Let  $W_{\hat{L}}$  be the Coxeter group determined by the 1-skeleton of  $\hat{L}$ , and having the triangulated 2-sphere  $\hat{L}$  as its nerve. Then  $\partial W_{\hat{L}}$  is homeomorphic to  $S^2$ . By Proposition 3.2(1), it follows that  $\partial W$  is planar.

Let  $\mathcal{T}$  be the collection of three generator special subgroups of W. Each element  $W' \in \mathcal{T}$  is an infinite triangle reflection group, i.e., either Euclidean or hyperbolic type. By Proposition 3.2 the circle boundary of each  $W' \in \mathcal{T}$  embeds in  $\partial W$ , and these circles are pairwise disjoint. Since  $\partial W$  is planar and contains more than one circle, it must be homeomorphic to the Sierpiński carpet by Proposition 3.5.

The group W is hyperbolic relative to  $\mathcal{T}$  by [Cap]. Hung Cong Tran has shown that the Bowditch boundary is the quotient space obtained from the visual boundary  $\partial W$  by collapsing the limit sets of the three generator special subgroups and their conjugates to points [Tra]. Since W has Property FA, its Bowditch boundary  $\partial(W, \mathcal{T})$  has no cut points [Bow]. It follows that the limit set of a three-generator special subgroup (or any of its conjugates) is always a peripheral circle of the Sierpiński carpet.

In fact, the group W in the preceding proposition acts on  $\mathbb{H}^3$  as a geometrically finite reflection group, as described below.

**Remark 3.7.** In the special case where the nerve is  $K_4$  and every  $m_{st} = 3$ , the group W is an arithmetic nonuniform lattice acting on  $\mathbb{H}^3$  as the group generated by the reflections in the faces of a regular ideal tetrahedron and is commensurable with the fundamental group of the figure eight knot complement and the Bianchi group PGL(2,  $\mathcal{O}_3$ ). The relationship between W and the figure eight knot group is discussed, for example, by Maclachlan–Reid (see Section 4.7.1 and Figure 13.2 of [MRe]).

More generally each Coxeter group with nerve  $K_4$  acts as a reflection group on a convex subset of  $\mathbb{H}^3$  with fundamental chamber a possibly ideal convex polytope. Start with the triangulation  $\hat{L}$  of  $S^2$  described in the proof of Proposition 3.6, and replace each right-angled cone on a Euclidean triangle with a single 2-simplex. The dual polytope K has a (possibly ideal) hyperbolic metric by Andreev's theorem (see Theorem 3.5 of [Sch] for a detailed explanation). The reflections in the faces of K generate a Coxeter group that contains W as a special subgroup. The union of all W-translates of K is a convex subspace of  $\mathbb{H}^3$  on which Wacts as a reflection group with fundamental chamber K.

### 4. Proof of the main theorem

The goal of this section is to prove that the boundary of a Coxeter group W is homeomorphic to the Menger curve when the nerve is  $K_n$  for  $n \ge 5$ . By Proposition 3.5, it suffices to show that  $\partial W$  is nonplanar when  $n \ge 5$ . Thus the following result completes the proof of Theorem 1.2.

**Proposition 4.1.** If W is any Coxeter group with nerve  $K_n$  for  $n \ge 5$ , then the complete graph  $K_5$  embeds in  $\partial W$ . In particular,  $\partial W$  is not planar.

*Proof.* Let  $W_5$  be any five generator special subgroup of W. By Proposition 3.2(1) it suffices to embed  $K_5$  into  $\partial W_5$ .

Suppose  $s_1, \ldots, s_5$  are the five generators of  $W_5$ . For each  $i \in \{1, \ldots, 5\}$ , let  $W_4^i$  be the special subgroup of  $W_5$  generated by  $\{s_1, \ldots, \hat{s_i}, \ldots, s_5\}$ . The limit set  $\Lambda W_4^i$  is homeomorphic to the Sierpiński carpet by Proposition 3.6. Similarly for each  $i \neq j$  in  $\{1, \ldots, 5\}$  let  $W_3^{i,j}$  denote the special subgroup generated by  $\{s_1, \ldots, \hat{s_i}, \ldots, \hat{s_j}, \ldots, \hat{s_j}, \ldots, s_5\}$ , whose limit set is a circle.

Since  $W_3^{i,j}$  is a subgroup of  $W_4^i$ , its limit set is a peripheral circle of the Sierpiński carpet  $\Lambda W_4^i$ , and similarly it is a peripheral circle in the carpet  $\Lambda W_4^j$ . Indeed this circle is precisely the intersection of these two Sierpiński carpets by Proposition 3.2(2). The five Sierpiński carpets  $\Lambda W_4^i$  and their circles of intersection are illustrated in Figure 2.

Choose points  $p_{i,j}$  on the circles  $\Lambda W_3^{i,j}$  such that  $p_{i,j} = p_{j,i}$  for  $i \neq j$ . Let  $E_4^i$  be a collection of 4-pointed stars for  $1 \leq i \leq 5$ . For a fixed i, we label the four edges of  $E_4^i$  as  $e_j^i$ , where  $1 \leq j \leq 5$  and  $j \neq i$ . By Proposition 2.3 for every i, there is a topological embedding  $h_i: E_4^i \to \partial W_4^i$  such that  $h_i \circ e_j^i(1) = p_{i,j} \in \partial W_3^{i,j}$ . Then  $h_i \circ e_j^i(0)$  is the center of the star in  $\Lambda W_4^i$ , and we will denote it by  $v_i$ .

The union of the five stars is an embedded complete graph  $K_5$  in  $\partial W_5$ . Indeed, we have 5 vertices  $v_1, \ldots, v_5$  and an edge between every two vertices.

216



The five Sierpiński carpets  $\Lambda W_4^1, \dots \Lambda W_4^5$  and their circles of intersection. The dotted edges between carpets indicate the pairs of peripheral circles that are identified in  $\partial W_5$ .

An edge between vertices  $v_i$  and  $v_j$  is given by concatenating the images of the edges  $e_j^i$  and  $e_i^j$  in  $\Lambda W_4^i$  and  $\Lambda W_4^j$  respectively. These edges do not intersect except at their endpoints  $v_i$ .

Acknowledgements. During the preparation of this paper, the authors had many helpful conversations with Craig Guilbault, Mike Mihalik, Wiktor Mogilski, Boris Okun, Kim Ruane, and Kevin Schreve. We are grateful for the insights obtained during these conversations. The authors also thank the referees for useful feedback that has led to improvements in the exposition.

This work was partially supported by a grant from the Simons Foundation (#318815 to G. Christopher Hruska).

# References

[Ben]	N. BENAKLI, Polyèdres hyperboliques: passage du local au global. Ph.D. thesis, Université Paris–Sud (1992).
[Bes]	<ul> <li>M. BESTVINA, Local homology properties of boundaries of groups. <i>Michigan Math. J.</i> 43 (1996), 123–139. Doi 10.1307/mmj/1029005393. Zbl 0872.57005 MR 1381603</li> </ul>
[BK]	M. BONK and B. KLEINER, Quasi-hyperbolic planes in hyperbolic groups. Proc. Amer. Math. Soc. 133 (2005), 2491–2494. Doi 10.1090/S0002-9939-05- 07564-7. Zbl 1123.20036 MR 2146190
[BKK]	M. BESTVINA, M. KAPOVICH and B. KLEINER, Van Kampen's embedding obstruction for discrete groups. <i>Invent. Math.</i> 150 (2002), 219–235. Doi 10.1007/s00222-002-0246-7. Zbl 1041.57016 MR 1933584
[BM]	M. BESTVINA and G. MESS, The boundary of negatively curved groups. J. Amer. Math. Soc. 4 (1991), 469–481. Zbl 0767.20014 MR 1096169
[Bow]	<ul> <li>B. BOWDITCH, Peripheral splittings of groups. <i>Trans. Amer. Math. Soc.</i> 353 (2001), 4057–4082. Zbl 1037.20041 MR 1837220</li> </ul>
[Cap]	PE. CAPRACE, Buildings with isolated subspaces and relatively hyperbolic Coxeter groups. <i>Innov. Incidence Geom.</i> <b>10</b> (2009), 15–31. Zbl1254.20036 MR 2665193
[Cha]	<ul> <li>C. CHAMPETIER, Propriétés statistiques des groupes de présentation finie. <i>Adv.</i> <i>Math.</i> <b>116</b> (1995), 197–262. Doi 10.1006/aima.1995.1067. Zbl 0847.20030 MR 1363765</li> </ul>
[Dav1]	M. W. DAVIS, Groups generated by reflections and aspherical manifolds not covered by Euclidean space. Ann. of Math. (2) 117 (1983), 293–324. Zbl 0531.57041
[Dav2]	— The Geometry and Topology of Coxeter Groups, vol. 32 of London Mathematical Society Monographs Series. Princeton University Press, Princeton, NJ (2008). Zbl1142.20020 MR 2360474
[Dave]	R. J. DAVERMAN, Decompositions of manifolds, vol. 124 of Pure and Applied Mathematics. Academic Press, Inc., Orlando, FL (1986). Zbl 0608.57002 MR 0872468
[DGP]	F. DAHMANI, V. GUIRARDEL and P. PRZYTYCKI, Random groups do not split. <i>Math. Ann.</i> <b>349</b> (2011), 657–673. Doi 10.1007/s00208-010-0532-4. Zbl 1278.20051
[DO]	M. W. DAVIS and B. OKUN, Vanishing theorems and conjectures for the l <sup>2</sup> - homology of right-angled Coxeter groups. <i>Geom. Topol.</i> 5 (2001), 7–74. Doi 10.2140/gt.2001.5.7. Zbl 1118.58300 MR 1812434
[Gro]	<ul> <li>M. GROMOV, Asymptotic invariants of infinite groups. In G. A. NIBLO and M. A. ROLLER (eds.), <i>Geometric group theory, Vol. 2 (Sussex, 1991)</i>. Cambridge Univ. Press, Cambridge (1993). pp. 1–295. Zb1 0841.20039 MR 1253544</li> </ul>

Nonhyperbolic Coxeter groups with Menger boundary

M. HAULMARK, Boundary classification and two-ended splittings of groups [Hau] with isolated flats. J. Topol. 11 (2018), 645-665. Doi 10.1112/topo.12062. Zbl 06963801 MR 3830879 G.C. HRUSKA and B. KLEINER, Hadamard spaces with isolated flats. Geom. [HK] Topol. 9 (2005), 1501–1538. With an appendix by the authors and Mohamad Hindawi. Doi 10.2140/gt.2005.9.1501. Zbl 1087.20034 MR 2469527 [HR1] G. C. HRUSKA and K. RUANE, Connectedness properties and splittings of groups with isolated flats. Preprint, arXiv:1705.00784 [math.GR] [Hru] G.C. HRUSKA, Nonpositively curved 2-complexes with isolated flats. Geom. Topol. 8 (2004), 205–275. Zb11063.20048 MR 2033482 [KK] M. KAPOVICH and B. KLEINER, Hyperbolic groups with low-dimensional boundary. Ann. Sci. École Norm. Sup. (4) 33 (2000), 647-669. Doi 10.1016/S0012-9593(00)01049-1. Zbl 0989.20031 MR 1834498 W. J. MOGILSKI, The fattened Davis complex and weighted  $L^2$ -(co)homology [Mog] of Coxeter groups. Algebr. Geom. Topol. 16 (2016), 2067-2105. Doi 10.2140/agt.2016.16.2067. Zbl 1387.20030 MR 3546459 R. MOORE, Concerning upper semi-continuous collections of continua. [Moo] Trans. Amer. Math. Soc. 27 (1925), 416–428. Doi 10.2307/1989234. JFM 54.0645.07 MR 1501320 [Mou] G. MOUSSONG, Hyperbolic Coxeter groups. Ph.D. thesis, Ohio State Univ. (1988). M. MIHALIK and K. RUANE, CAT(0) groups with non-locally connected [MR] boundary. J. London Math. Soc. (2) 60 (1999), 757-770. Zb1 0973.20033 MR 1753812 C. MACLACHLAN and A.W. REID, The arithmetic of hyperbolic 3-manifolds, [MRe] vol. 219 of Graduate Texts in Mathematics. Springer-Verlag, New York (2003). Doi 10.1007/978-1-4757-6720-9. Zbl 1025.57001 MR 1937957 D. OSAJDA and J. ŚWIĄTKOWSKI, On asymptotically hereditarily as-[OS]pherical groups. Proc. Lond. Math. Soc. (3) 111 (2015), 93-126. Doi 10.1112/plms/pdv021. Zbl 1360.20040 MR 3404777 [Pan] P. PANSU, Cohomologie  $L^p$ : invariance sous quasiisométries (1995). Preprint. www.math.u-psud.fr/~pansu. [Sch] T.A. SCHROEDER, Geometrization of 3-dimensional Coxeter orbifolds and Singer's conjecture. Geom. Dedicata 140 (2009), 163-174. Doi 10.1007/s10711-008-9314-5. Zbl 1210.57020 MR 2504739 J.-P. SERRE, Arbres, amalgames, SL2, vol. 46 of Astérisque. Société Math-[Ser] ématique de France, Paris (1977). Written in collaboration with Hyman Bass. Zbl 0369.20013 MR 0476875 [Sie] W. SIERPIŃSKI, Sur une courbe cantorienne qui contient une image biunivoque et continue de toute courbe donnée. C.R. Acad. Sci. Paris 162 (1916), 629-632. JFM 46.0295.02 [Swe] E. L. SWENSON, A cut point theorem for CAT(0) groups. J. Differential Geom. 53 (1999), 327–358. Zbl 1038.20029 MR 1802725

220	M. HAULMARK, G.C. HRUSKA and B. SATHAYE
[Świ]	J. Świąткowski, Hyperbolic Coxeter groups with Sierpiński carpet boundary. Bull. Lond. Math. Soc. 48 (2016), 708–716. Doi 10.1112/blms/bdw034. Zbl 1396.20043 MR 3532145
[Tra]	<ul> <li>H. C. TRAN, Relations between various boundaries of relatively hyperbolic groups. <i>Internat. J. Algebra Comput.</i> 23 (2013), 1551–1572. Doi 10.1142/S0218196713500367. Zbl 1292.20048 MR 3143594</li> </ul>
[Why]	G. WHYBURN, Topological characterization of the Sierpiński curve. <i>Fund.</i> <i>Math.</i> <b>45</b> (1958), 320–324. Doi 10.4064/fm-45-1-320-324. Zbl 0081.16904 MR 0099638
	(Reçu le 28 janvier 2019)
Matthew	HALLIMARK Department of Mathematics 1326 Stevenson Center Vanderbilt

Matthew HAULMARK, Department of Mathematics, 1326 Stevenson Center, Vanderbilt University, Nashville, TN 37240, USA

e-mail: m.haulmark@vanderbilt.edu

G. Christopher HRUSKA, Department of Mathematical Sciences, University of Wisconsin-Milwaukee, PO Box 413, Milwaukee, WI 53211, USA

e-mail: chruska@uwm.edu

Bakul SATHAYE, Department of Mathematics and Statistics, Wake Forest University, 127 Manchester Hall. PO Box 7388, Winston-Salem, NC 27109, USA

*e-mail:* sathaybv@wfu.edu

© Fondation L'Enseignement Mathématique