

Representations of the Gauge groups of electrodynamics and general relativity

Autor(en): **Girardello, L. / Wyss, W.**

Objekttyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **45 (1972)**

Heft 2

PDF erstellt am: **24.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-114376>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Representations of the Gauge Groups of Electrodynamics and General Relativity

by L. Girardello¹⁾

Institute for Theoretical Physics, University of Colorado, Boulder, 80302, USA

and W. Wyss²⁾

Department of Mathematics and Institute for Theoretical Physics,
University of Colorado, Boulder, 80302, USA

(12. XI. 71)

Abstract. A semidirect product of the gauge groups of Electrodynamics and General Relativity is determined and unitarily represented on a Hilbertspace of the type $\mathfrak{L}_2(\mathcal{S}', \mu)$.

1. Introduction

Electrodynamics and General Relativity, being rest-mass zero theories of vector and tensor fields, enjoy special symmetries governed by gauge groups. Some implications of these symmetries have been intensively studied [1]. In fact a linearized Lorentz-covariant theory of gravitation gets uniquely promoted to Einstein's General Relativity with the help of the gauge group [2]. The generators of the gauge groups of Electrodynamics and General Relativity form a nuclear Lie algebra that we show is similar to a current algebra [3]. The Gel'fand-Vilenkin formalism [4] then gives representations of the integrated group, being the semidirect product $\mathcal{S}(\mathbb{R}^4) \wedge \text{diff}_{\mathcal{S}}(\mathbb{R}^4)$, where $\mathcal{S}(\mathbb{R}^4)$ is the gauge group for Electrodynamics and $\text{diff}_{\mathcal{S}}(\mathbb{R}^4)$ the gauge group for General Relativity. The semidirect product structure corresponds to the Klein-Kaluza formalism [5]. The representation spaces are of the type $\mathfrak{L}_2(\mathcal{S}', \mu)$, with μ a cylindrical measure on \mathcal{S}' . We might be enlightened in the physics of these representations by current investigations into a Wightman formalism for Quantum Electrodynamics and for the Theory of Gravitation [6].

2. The Gauge Lie Algebra for Electrodynamics and General Relativity

Let A_μ denote the potential in Electrodynamics and $g_{\mu\nu}$ the potential in General Relativity, $\mu, \nu = 1, 2, 3, 4$. Let $\phi \in \mathcal{S}(\mathbb{R}^4)$ be real valued, then the transformations $E(\phi)$ defined by

$$E(\phi)A_\mu = A_\mu + \phi_{,\mu} \quad (2.1)$$

¹⁾ Address for 1972: Istituto di Fisica dell'Università, 43100 Parma, and Istituto Nazionale di Fisica Nucleare, Sezione di Milano (Italy).

²⁾ Supported by NSF, GP 19479.

constitute the gauge group of Electrodynamics. The Lie algebra of the gauge group in General Relativity [2] originates from infinitesimal coordinate transformations

$$T(\xi)x^\mu = x^\mu - \xi^\mu(x), \quad (2.2)$$

where ξ is a real valued vector field $\{\xi^\mu\}$ with $\xi^\mu \in \mathcal{S}(\mathbb{R}^4)$. The generators $T(\xi)$ transform A_μ and $g_{\mu\nu}$ as follows

$$T(\xi)A_\mu = A_\mu + A_\alpha \xi_{,\mu}^\alpha + \xi^\alpha A_{\mu,\alpha} \quad (2.3)$$

$$T(\xi)g_{\mu\nu} = g_{\mu\nu} + g_{\mu\beta} \xi_{,\nu}^\beta + g_{\alpha\nu} \xi_{,\mu}^\alpha + \xi^\alpha g_{\mu\nu,\alpha}. \quad (2.4)$$

The group theoretical commutators lead to the following Lie algebra \mathfrak{L}

$$[E(\phi), E(\psi)] = 0 \quad (2.5)$$

$$[E(\phi), T(\xi)] = -E(\xi \cdot \text{grad } \phi) \quad (2.6)$$

$$[T(\xi), T(\eta)] = T([\xi, \eta]), \quad (2.7)$$

where

$$[\xi, \eta]^\mu = \eta_{,\nu}^\mu \xi^\nu - \xi_{,\nu}^\mu \eta^\nu \quad (2.8)$$

This Lie algebra is similar to the charge-current algebra of a nonrelativistic field theory, which has been discussed by G. A. Goldin [3]. Hence the Gel'fand-Vilenkin formalism for nuclear Lie groups [4] will give us unitary representations of a gauge group G , whose Lie algebra is \mathfrak{L} .

3. Representations of \mathfrak{L} on the Field Algebra

Recall that the Field algebra \mathfrak{U} , as a topological vector space, is given by the topological direct sum $\mathfrak{U} = \bigoplus_{n=0}^{\infty} \mathcal{S}^n(\mathbb{R}^4)$, where $\mathcal{S}^0(\mathbb{R}^4) = \mathbb{C}$ [7]. We then get the following representation for the generators $E(\phi)$ and $T(\xi)$:

$$(\mathbb{E}(\phi)f)_n(x_1, \dots, x_n) = \sum_{k=1}^n \phi(x_k) f_n(x_1, \dots, x_n) \quad (3.1)$$

$$\begin{aligned} (\mathbb{T}(\xi)f)_n(x_1, \dots, x_n) &= \sum_{k=1}^n \left\{ \xi^\mu(x_k) \frac{\partial}{\partial x_k^\mu} f_n(x_1, \dots, x_n) \right. \\ &\quad \left. + \frac{1}{2} \frac{\partial}{\partial x_k^\mu} \xi^\mu(x_k) \cdot f_n(x_1, \dots, x_n) \right\} \end{aligned} \quad (3.2)$$

It is readily found that $\mathbb{T}(\xi)$ is antihermitian with respect to the \mathfrak{L}_2 -inner product on $\mathcal{S}^n(\mathbb{R}^4)$, and satisfies the commutation relation (2.7). $\mathbb{E}(\phi)$ is hermitian and satisfies (2.5) and (2.6).

4. The Gauge Group G

We first look at the one-parameter subgroup of G of the form

$$(e^{t\mathbb{T}(\xi)}f)_n(x_1, \dots, x_n) \equiv (V(\Phi_t(\xi, \cdot))f)_n(x_1, \dots, x_n). \quad (4.1)$$

Lemma 4.1. $(V(\Phi_t(\xi, \cdot))f)_n(x_1, \dots, x_n)$

$$= f_n(\Phi_t(\xi, x_1), \dots, \Phi_t(\xi, x_n)) \prod_{k=1}^n \sqrt{\det \frac{\partial \Phi_t(\xi, x_k)}{\partial x_k}}, \quad (4.2)$$

where $\Phi_t(\xi, \cdot)$ is the flow generated by the vector field $\xi(x)$, i.e.

$$\frac{d}{dt} \Phi_t(\xi, x) = \xi(\Phi_t(\xi, x)), \quad \Phi_0(\xi, x) = x. \quad (4.3)$$

Proof: See Appendix A.

Since ξ is an \mathcal{S} -vector field, the corresponding flow $\Phi_t(\xi, x)$ is then C_∞ in x for all t . Let $\text{diff}_{\mathcal{S}}(\mathbb{R}^4)$ stand for the C_∞ -diffeomorphisms on \mathbb{R}^4 , generated by the flows $\Phi_t(\xi, x)$. The composition of flows turns $\text{diff}_{\mathcal{S}}(\mathbb{R}^4)$ into a group, the gauge group for General Relativity. Hence on the Field algebra \mathfrak{U} we have for $\Phi \in \text{diff}_{\mathcal{S}}(\mathbb{R}^4)$

$$(V(\Phi)f)_n(x_1, \dots, x_n) = f_n(\Phi(x_1), \dots, \Phi(x_n)) \prod_{k=1}^n \sqrt{\det \frac{\partial \Phi(x_k)}{\partial x_k}}. \quad (4.4)$$

The gauge group for Electrodynamics is $\mathcal{S}(\mathbb{R}^4)$ under addition and its unitary representation on \mathfrak{U} is given by

$$(e^{i\mathbb{E}(\phi)}f)_n(x_1, \dots, x_n) \equiv (U(\phi)f)_n(x_1, \dots, x_n) \quad (4.5)$$

$$(U(\phi)f)_n(x_1, \dots, x_n) = e^{i \sum_{k=1}^n \phi(x_k)} f_n(x_1, \dots, x_n). \quad (4.6)$$

The full group is now given by the semidirect product

$$G = \mathcal{S}(\mathbb{R}^4) \wedge \text{diff}_{\mathcal{S}}(\mathbb{R}^4) \quad (4.7)$$

with the semidirect product map

$$\mathcal{S}(\mathbb{R}^4) \times \text{diff}_{\mathcal{S}}(\mathbb{R}^4) \rightarrow \mathcal{S}(\mathbb{R}^4) \quad (4.8)$$

$$(\phi, \Phi)(x) = (\phi \circ \Phi)(x) = \phi(\Phi(x)). \quad (4.9)$$

The multiplication law of G thus is

$$(\phi_1 \wedge \Phi_1)(\phi_2 \wedge \Phi_2) = (\phi_1 + \phi_2 \circ \Phi_1) \wedge \Phi_2 \circ \Phi_1. \quad (4.10)$$

For a unitary representation this reads

$$U(\phi_1)V(\Phi_1)U(\phi_2)V(\Phi_2) = U(\phi_1 + \phi_2 \circ \Phi_1)V(\Phi_2 \circ \Phi_1). \quad (4.11)$$

The gauge group G inherits a nuclear topology from $\mathcal{S}(\mathbb{R}^4)$, and \mathcal{S} is a normal subgroup of G . The operators $U(\phi)$ and $V(\Phi)$ are unitary on the Fockspace \mathfrak{H}_f which is the completion of the field algebra \mathfrak{U} in the norm

$$\|f\|^2 = \sum_{k=1}^{\infty} \|f_n\|^2 \quad (4.12)$$

$$\|f_n\|^2 = \int \overline{f_n(x_1, \dots, x_n)} f_n(x_1, \dots, x_n) dx_1 \dots dx_n \quad (4.13)$$

5. Representations of the Gauge Group G

Here we give a summary of the Gel'fand-Vilenkin method [4] as applied to G . For details see Ref. [3]. A continuous unitary representation $U(\phi)$ of the normal subgroup $\mathcal{S}(\mathbb{R}^4) \subset G$ on a Hilbertspace \mathfrak{H} with cyclic vector Ω gives us a functional $L(\phi)$, defined by

$$L(\phi) = (\Omega, U(\phi)\Omega) \quad (5.1)$$

$L(\phi)$ has the following properties

1. L is continuous on $\mathcal{S}(\mathbb{R}^4)$ (5.2)

2. $L(0) = 1$ (5.3)

3. $\sum_{k,l=1}^N \bar{c}_k c_l L(\phi_k - \phi_l) \geq 0$ for any complex numbers c_1, \dots, c_N (5.4)

and hence by Bochner's Theorem there exists a unique cylinder measure μ on \mathcal{S}' such that

$$L(\phi) = \int_{\mathcal{S}'} e^{i(T, \phi)} d\mu(T). \quad T \in \mathcal{S}'(\mathbb{R}^4) \quad (5.5)$$

A strongly continuous representation of \mathcal{S} in $\mathfrak{H} = \mathfrak{L}_2(\mathcal{S}', \mu)$ is thus given by

$$U(\phi)\Psi(T) = e^{i(T, \phi)} \Psi(T) \quad (5.6)$$

and the cyclic vector Ω is realized by the unit function on \mathcal{S}' . For $\Phi \in \text{diff}_{\mathcal{S}}(\mathbb{R}^4)$ let $\Phi^* : \mathcal{S}'(\mathbb{R}^4) \rightarrow \mathcal{S}'(\mathbb{R}^4)$ be defined by

$$(\Phi^* T, \phi) = (T, \phi \circ \Phi) \quad (5.7)$$

and define a transformed measure μ^{Φ^*} on \mathcal{S}' by

$$\mu^{\Phi^*}(T) = \mu(\Phi^* T). \quad (5.8)$$

The group $\text{diff}_{\mathcal{S}}(\mathbb{R}^4)$ is then represented on $\mathfrak{L}_2(\mathcal{S}', \mu)$ by

$$V(\Phi) \Psi(T) = \chi(\Phi, T) \Psi(\Phi^* T) \sqrt{\frac{d\mu^{\Phi^*}(T)}{d\mu(T)}} \quad (5.9)$$

where $d\mu^{\Phi^*}/d\mu$ is the Radon-Nikodym derivative and $\chi(\Phi, T)$ a complex valued function of modulus one, satisfying

$$\chi(\Phi_2, T) \chi(\Phi_1, \Phi_2 T) = \chi(\Phi_1 \circ \Phi_2, T). \quad (5.10)$$

The measure μ on \mathcal{S}' is quasiinvariant under $\text{diff}_{\mathcal{S}}(\mathbb{R}^4)$, i.e. μ and μ^{Φ^*} have the same set of measure zero. Thus the expectation functional $L(\phi)$ defines a representation of G up to a phase function.

Remarks:

1. The Fock representation of G on \mathfrak{H}_F corresponds to a Gaussian measure. In the n -particle space $\mathfrak{H}_F^{(n)} \subset \mathfrak{H}_F$, μ is concentrated on the set $T = \{T_{x_1} + \dots + T_{x_n}, x_i \neq x_k\}$, where $(T_x, \phi) = \phi(x)$.
 $d\mu(T_{x_1} + \dots + T_{x_n}) = \pi^{-2n} e^{-\|x_1\|^2 - \dots - \|x_n\|^2} dx_1 \dots dx_n$. Observe that this representation is on a Fockspace of scalar functions.
2. For the recovery of the infinitesimal generators $E(\phi)$ and $T(\xi)$, Goldin [3] gives sufficient conditions, expressed as properties of the measure μ .
3. Here we do not investigate if the Gel'fand-Vilenkin representations of G are physical. If they are, then operators commuting with the $U(\phi)$ are candidates for observables in Quantum Electrodynamics and operators commuting with $V(\Phi)$ are candidates for observables in a quantum theory of gravitation.

Acknowledgments

We would like to thank Professors S. Albeverio, A. O. Barut, R. Haag, J. R. Klauder and F. Strocchi for enlightening discussions. L.G. wishes to thank Professors A. O. Barut and W. E. Brittin for the hospitality extended to him at the Institute for Theoretical Physics.

Appendix A

We want to show that

$$(e^{tT(\xi)} f)_n(x_1, \dots, x_n) = f_n(\Phi_t(\xi, x_1), \dots, \Phi_t(\xi, x_n)) \sqrt{\prod_{k=1}^n \det \frac{\partial \Phi_t(\xi, x_k)}{\partial x_k}} \quad (\text{A.1})$$

where

$$(T(\xi)f)_n(x_1, \dots, x_n) = \sum_{k=1}^n \left\{ \xi^\mu(x_k) \frac{\partial}{\partial x_k^\mu} + \frac{1}{2} \frac{\partial \xi^\mu(x_k)}{\partial x_k^\mu} \right\} f_n(x_1, \dots, x_n). \quad (\text{A.2})$$

It suffices to give the proof for one variable x , i.e.

$$e^{tT(\xi)} f(x) = f(\Phi_t(\xi, x)) \sqrt{\det \frac{\partial \Phi_t(\xi, x)}{\partial x}} \quad (\text{A.3})$$

with

$$T(\xi)f(x) = \left\{ \xi^\mu(x) \frac{\partial}{\partial x^\mu} + \frac{1}{2} \frac{\partial \xi^\mu(x)}{\partial x^\mu} \right\} f(x). \quad (\text{A.4})$$

For fixed ξ and x let

$$F(f, t) = e^{tT(\xi)} f(x) \quad (\text{A.5})$$

$$G(f, t) = \sqrt{\det \frac{\partial \Phi_t(\xi, x)}{\partial x}} f(\Phi_t(\xi, x)). \quad (\text{A.6})$$

Then $F(f, t)$ satisfies the following differential equation

$$\frac{\partial F(f, t)}{\partial t} = F(\mathbb{T}(\xi)f, t) \quad (\text{A.7})$$

with the initial condition

$$F(f, 0) = f. \quad (\text{A.8})$$

Similarly, using the formula

$$\frac{\partial}{\partial t} \det A(t) = \det A(t) \cdot \text{Tr} \left(A(t)^{-1} \frac{\partial A(t)}{\partial t} \right), \quad (\text{A.9})$$

we get a differential equation for $G(f, t)$.

$$\begin{aligned} \frac{\partial G(f, t)}{\partial t} &= \frac{\partial f(\Phi_t(\xi, x))}{\partial \Phi_t(\xi, x)} \frac{\partial \Phi_t(\xi, x)}{\partial t} \sqrt{\det \frac{\partial \Phi_t(\xi, x)}{\partial x}} \\ &\quad + \frac{1}{2} f(\Phi_t(\xi, x)) \sqrt{\det \frac{\partial \Phi_t(\xi, x)}{\partial x}} \frac{\partial x^\mu}{\partial \Phi_t^\nu(\xi, x)} \frac{\partial}{\partial t} \frac{\partial \Phi_t^\nu(\xi, x)}{\partial x^\mu} \\ &= \sqrt{\det \frac{\partial \Phi_t(\xi, x)}{\partial x}} \left\{ \frac{\partial f(\Phi_t(\xi, x))}{\partial \Phi_t(\xi, x)} \cdot \xi(\Phi_t(\xi, x)) \right. \\ &\quad \left. + \frac{1}{2} f(\Phi_t(\xi, x)) \frac{\partial x^\mu}{\partial \Phi_t^\nu(\xi, x)} \cdot \frac{\partial}{\partial x^\mu} \xi^\nu(\Phi_t(\xi, x)) \right\} \\ &= G(\mathbb{T}(\xi)f, t). \end{aligned} \quad (\text{A.10})$$

The initial condition is

$$G(f, 0) = f. \quad (\text{A.11})$$

Hence F and G satisfy the same differential equation of first order and the same initial condition, and thus have to be equal.

REFERENCES

- [1] R. Utiyama, Phys. Rev. **101**, 1597 (1956), and e.g. the review: T. W. Kibble, *High Energy Physics and Elementary Particles* (I.A.E.A., Wien 1965).
- [2] W. Wyss, Helv. phys. Acta **38**, 469 (1965).
- [3] G. A. Goldin, J. Math. Phys. **12**, 462 (1971); J. Grodlik and D. H. Sharp, Phys. Rev. [D] **1**, 1531 (1970).
- [4] I. Gel'fand and N. Vilenkin, *Generalized Functions*, Vol. 4, Chap. IV. (Academic Press, New York 1964).
- [5] W. Pauli, *Theory of Relativity* (Pergamon Press, 1967).
- [6] F. Strocchi, Phys. Rev. **162**, 1429 (1967); Phys. Rev. [D] **2**, 2334 (1970), Phys. Rev. **166**, 1302 (1968); and L. L. Bracci and F. Strocchi, preprints, S.N.S. 71/1, and references therein.
- [7] H. J. Borchers, Nuovo Cim. **24**, 214 (1962).