

Zusammenfassungen der letzten eingegangenen Arbeiten = Résumés des derniers articles reçus

Objekttyp: **Group**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **48 (1975)**

Heft 5-6

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

HELVETICA PHYSICA ACTA
Zusammenfassungen der letzten eingegangenen Arbeiten
Résumés des derniers articles reçus

The Raman Spectrum of Ferroelectric PbTiO₃

by R. A. FREY and E. SILBERMAN

Physics Department, Fisk University, Nashville, Tennessee 37203

(1. VII. 75)

Abstract. Raman scattering in ferroelectric single domain crystal PbTiO₃ reveals 14 phonons, which can readily be assigned to $k = 0$ phonons in the crystal class C_{4v}. In addition, seven quasi-phonons can be observed. This set of phonons allows the calculation of the non-observed E(LO3) phonon, a qualitative assessment of the action of long range and short range forces in the crystal as well as the calculation of a new set of dielectric constants. The ferroelectric phase has been found to persist down to 77 K.

A Mechanical Quantum Measuring Process

by BARBARA WHITTEN-WOLFE

Western College, Miami University, Oxford, Ohio 45056 (USA)

and GÉRARD G. EMCH

Depts. of Mathematics and of Physics, University of Rochester,
Rochester, New York 14627 (USA)

(22. VIII. 75)

Abstract. A mechanistic analysis of the quantum measuring process is proposed. This takes into account both the reduction of the wave packet and the recording of the measurement on a macroscopic apparatus. An exactly solvable model is given to sustain the theory.

Strukturuntersuchungen an Alkalihyperoxiden

von M. ZIEGLER, M. ROSENFELD, und W. KÄNZIG

Laboratorium für Festkörperphysik der ETH Zürich, 8049 Zürich, Schweiz,

und P. FISCHER

Institut für Reaktortechnik der ETH Zürich, EIR, 5303 Würenlingen, Schweiz

(29. VIII. 75)

Abstract. The alkali hyperoxides undergo with varying temperature a number of phase transitions in which changes of the orientational order of the O₂⁻-molecules play an important part.

By means of x-ray analysis of NaO₂, KO₂, RbO₂ and CsO₂ single crystals (including a study of the diffuse scattering) average structures, correlations between the orientations of the O₂⁻-molecules, and relationships between the various phases and domain structures were established.

A neutron diffraction analysis of NaO₂ single crystals and polycrystalline samples indicates magnetic short range order in the marcasite phase.

Electrostriktion Dotierter Alkalihalogenide

by H. BURKARD, W. KÄNZIG und M. ROSSINELLI

Laboratorium für Festkörperphysik der ETH, 8049 Zürich, Schweiz

(18. VIII. 75)

Abstract. The electrostriction of doped alkali halide crystals was investigated. For the paraelectric systems $\text{KCl}:\text{OH}^-$, $\text{KCl}:\text{Li}^+$, $\text{NaBr}:\text{F}^-$, $\text{RbCl}:\text{Ag}^+$ and $\text{RbBr}:\text{Ag}^+$ in which the equilibrium position of the substitutional impurities is off-center the electrostriction consists of the following three contributions:

- (a) The electrostriction of the host lattice.
- (b) The deformation of the lattice as a result of the reorientation of the permanent electric-elastic dipoles.
- (c) The deformation arising from the field induced shift of the equilibrium position of the defects.

In $\text{KBr}:\text{Li}^+$ the electrostriction consists only of contributions (a) and (c), indicating that it is not a typical off-center system.

Contribution (c) does not saturate in contrast to (b) and thus is particularly important for *strong electric fields*. RbCl , for example, containing $6 \cdot 10^{17} \text{ Ag}^+$ -ions per cm^3 exhibits a temperature independent electrostrictive effect which is about 100 times larger than the effect in the pure material.

At *low field strengths* the electromechanical behaviour of $\text{KCl}:\text{OH}^-$ and $\text{KCl}:\text{Li}^+$ is dominated by contribution (b). At high hydroxyl concentration the electric and elastic interaction of the dipoles manifests itself in the electrostriction.

Classical Electrodynamics with Extended Charges

by A. RICHOZ

Institut für Theoretische Physik der Universität Zürich,
Schönberggasse 9, CH-8001 Zürich, Switzerland

(17. IX. 75)

Abstract. In the first part, the problem of the existence, uniqueness, smoothness of the solution of the coupled Maxwell–Newton equations is investigated for a charge distribution with compact support. In the second part, a detailed study of this problem for the case of a spherical shell distribution shows that the limit of Lorentz–Dirac is not physical.

A Remark on Pion Capture in Heavy Nuclei

by M. P. LOCHER and F. MYHRER

S.I.N., 5234 Villigen, Switzerland

(17. IX. 75)

Abstract. The yield of neutrons from the (π^-, xn) reactions on ^{165}Ho is calculated with a two nucleon absorption mechanism followed by (nucleon, xn) reactions using experimental input only. The comparison with the experimental yield curve supports this simple two-step model for heavy nuclei.

The $P(\phi)_2$ Green's Functions: Asymptotic Perturbation Expansion

by JONATHAN DIMOCK

Department of Mathematics, SUNY at Buffalo, Amherst, N.Y. 14226

(19. IX. 75)

Abstract. The real time Green's functions in the $P(\phi)_2$ quantum field theory are infinitely differentiable functions of the coupling constant λ up to and including $\lambda = 0$. It follows that the perturbation series are asymptotic as $\lambda \rightarrow 0^+$.