Zeitschrift:	Helvetica Physica Acta
Band:	50 (1977)
Heft:	3
Artikel:	Onsager's symmetry in higher order fluid dynamics
Autor:	Grmela, Miroslav
DOI:	https://doi.org/10.5169/seals-114866

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 05.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Onsager's symmetry in higher order fluid dynamics

By Miroslav Grmela

Centre de recherches mathématiques, Université de Montréal, Montréal, Québec H3C 3J7, Canada

(1. II. 1977)

Abstract. Example of the higher order fluid dynamics that possesses the properties of the family of dissipative dynamical systems of macroscopic physics, thus in particular the Onsager symmetry, is constructed. The non-equilibrium extension of the thermodynamic entropy depends on gradients of the fluid dynamic state variables.

I. Introduction

The abstract concept of the family of dissipative dynamical systems of macroscopic physics (hereafter called DDS) that has been developed in References [1], [2], provides a unified mathematical setting for a large class of dynamical systems phenomenologically introduced in macroscopic physics (e.g., kinetic equations of the Boltzmann and Enskog–Vlasov type, Navier–Stokes–Fourier fluid dynamics, nonequilibrium thermodynamics etc. [2]).

In this paper, we construct an example of the higher order fluid dynamics that possesses the properties of DDS. In Section II the Navier–Stokes–Fourier fluid dynamics and the concept of DDS are reviewed. The higher order fluid dynamics generalizing the Navier–Stokes–Fourier fluid dynamics and satisfying the properties of DDS is constructed in Section III. In Section IV a brief review of other approaches to higher order fluid dynamics is presented.

The state of a one component physical system is completely described in fluid mechanics (completely with respect to the fluid mechanics observations and measurements) by the triple $(E, N, \mathbf{U}) \equiv f$, where $E: \Omega \to \mathbb{R}_+$, $(\mathbf{r} \mapsto E(\mathbf{r}))$; $N: \Omega \to \mathbb{R}_+$, $(\mathbf{r} \mapsto N(\mathbf{r}))$; $\mathbf{U}: \Omega \to \mathbb{R}^3 \cdot (\mathbf{r} \mapsto \mathbf{U}(\mathbf{r}))$; the physical system is confined in $\Omega \subset \mathbb{R}^3$, we shall assume that volume of Ω equals to one. \mathbb{R}_+ denotes the positive real line. Let \mathscr{H} be the set of all admissible (again with respect to the fluid mechanic observations and measurements) states. We shall assume that \mathscr{H} is a smooth (infinite dimensional) manifold. We define in \mathscr{H} the involution $J: \mathscr{H} \to \mathscr{H}$; $(E, N, \mathbf{U}) \mapsto (E, N, -\mathbf{U})$. The subset of \mathscr{H} that consists of elements f of \mathscr{H} that are invariant with respect to J (i.e., the elements of the type (E, N, 0)) is denoted $\mathscr{H}^{(+)}$, its elements $f^{(+)}$. We can also say that \mathscr{H} is a bundle $\mathscr{H} \equiv (\mathscr{H}, \pi^{(+)}, \mathscr{H}^{(+)})$, where $\mathscr{H}^{(+)}$ is the base space, $\pi^{(+)}$ is the projection $\mathscr{H} \to \mathscr{H}^{(+)}$ and \mathscr{H} is the total space. In terms of fluid mechanics observations and measurements $N(\mathbf{r})$ is the local density of mass $E(\mathbf{r})$ is the local density of the fluid considered.

(1)

(1.1)

The time evolution of f is governed by the following equations:

$$\frac{\partial f}{\partial t} = Rf$$

in components

$$\begin{split} \frac{\partial E}{\partial t} &= -U_{\alpha} \nabla_{\alpha}(E) - \frac{1}{N} (P_{\alpha\beta} \nabla_{\alpha} U_{\beta} + \nabla_{\alpha} Q_{\alpha}) \\ \frac{\partial N}{\partial t} &= -\nabla_{\alpha} (N U_{\alpha}) \\ \frac{\partial U_{\alpha}}{\partial t} &= -U_{\beta} \nabla_{\beta} (U_{\alpha}) - \frac{1}{N} (\nabla_{\beta} P_{\alpha\beta}). \end{split}$$

The symbol $\nabla_{\alpha} \equiv \partial/\partial r_{\alpha}$, the summation convention is used. The tensor $P_{\alpha\beta}$ is symmetric $P_{\alpha\beta} = P_{\beta\alpha}$. If the quantities $P_{\alpha\beta}$ and Q_{α} are specified as functions of (E, N, \mathbf{U}) , or by using the terminology used in fluid mechanics, if the constitutive laws are specified, then (1) represents a self-contained system of the time evolution equations. The parameters introduced in the constitutive laws are the phenomenological quantities entering fluid dynamics (through these phenomenological quantities the individuality of the particular physical system is expressed) and (1) is a family of the time evolution equations parametrized by the phenomenological quantities. Boundary conditions for (E, N, \mathbf{U}) have to enter into the specification of \mathcal{H} . The class of the boundary conditions used in this paper (BC1)–(BC5) in Section II and (GBC1)–(GBC5) in Section III) reflects our interest in the thermodynamic equilibrium states and the time approach to these states.

Let \mathscr{L} be the set of all constitutive laws. We shall assume that the physical system considered is isotropic and thus we require that the functional dependence of $P_{\alpha\beta}$, Q_{α} on f is invariant with respect to the orthogonal transformations in Ω . This requirement determines the subset \mathcal{L}_0 of \mathcal{L} . (See, e.g., Reference [3] pp. 135.) Physical considerations of constitutive laws are based on the physical interpretation of P and Q ($P_{\alpha\beta}$ is the pressure tensor, Q_{α} is the heat flow). From the mathematical point of view, one can look for \mathscr{L}_{math} such that (1) will be mathematically well defined (e.g., the flow generated by (1) will exist etc.). It has been shown in Reference [1]that the Navier-Stokes-Fourier constitutive laws transform (2) into an example of DDS. According to the general theory of DDS it means in particular that (i) the study of the solutions of $R^{(+)}f = 0$ and $R^{(-)}f \mid_{R^{(+)}f=0} = 0$ gives rise to thermodynamics $(R^{(\pm)} \equiv \frac{1}{2}(R \pm JRJ))$; (ii) the flow exists, at least locally, in the vicinity of the solutions of the problem (i) that moreover correspond to the thermodynamically stable equilibrium states. In this paper we shall find a generalization of the Navier-Stokes–Fourier constitutive laws that transform (1) again into an example of DDS and the resulting system of partial differential equations is of order higher than two.

For later use, we shall write (1) also in the following form:

$$\frac{\partial f}{\partial t} = R^{(-)}(f) + R^{(+)}(f),$$

in components

$$\frac{\partial E}{\partial t} = R^{(-)}_{(E)}(f) + R^{(+)}_{(E)}(f)$$

$$\frac{\partial N}{\partial t} = R_{(N)}^{(-)}(f)$$

$$\frac{\partial U_{\alpha}}{\partial t} = R_{(U)\alpha}^{(-)}(f) + R_{(U)\alpha}^{(+)}(f),$$

$$R^{(\pm)} \equiv \frac{1}{2}(R \pm JRJ), \text{ thus } R^{(+)}(Jf) = R^{(+)}(f); R^{(-)}(Jf) = -R^{(-)}(f). \text{ If we write}$$

$$P_{\alpha\beta} = P_{\alpha\beta}^{(+)} + P_{\alpha\beta}^{(-)}$$

$$Q_{\alpha} = Q_{\alpha}^{(+)} + Q_{\alpha}^{(-)}$$

$$Q_{\alpha} = Q_{\alpha}^{(+)} + Q_{\alpha}^{(-)}$$

$$(2)$$

$$(P_{\alpha\beta}^{(\pm)}(Jf) = \pm P_{\alpha\beta}^{(\pm)}(F), \text{ similarly } Q_{\alpha}), \text{ then}$$

$$R_{(E)}^{(-)}(f) = -U_{\alpha}\nabla_{\alpha}E - \frac{1}{N}P_{\alpha\beta}^{(+)}\nabla_{\alpha}U_{\beta} - \frac{1}{N}\nabla_{\alpha}Q_{\alpha}^{(-)}$$

$$R_{(E)}^{(+)}(f) = -P_{\alpha\beta}^{(-)}\nabla_{\alpha}U_{\beta} - \nabla_{\alpha}Q^{(+)}$$

$$R_{(N)}^{(-)}(f) = -U_{\beta}\nabla_{\beta}U_{\alpha} - \frac{1}{N}\nabla_{\beta}P_{\alpha\beta}^{(+)}$$

$$R_{(U)\alpha}^{(+)}(f) = -\frac{1}{N}\nabla_{\beta}P_{\alpha\beta}^{(-)}.$$
(2)

II. Navier-Stokes-Fourier fluid dynamics

The Navier-Stokes-Fourier constitutive laws are the following:

$$P_{\alpha\beta}^{(+)} = N^* \,\delta_{\alpha\beta}$$

$$P_{\alpha\beta}^{(-)} = p^{(-)} \,\delta_{\alpha\beta} + p_{\alpha\beta}^{(-)}$$

$$Q_{\alpha}^{(+)} = -\lambda \nabla_{\alpha} E^*$$

$$Q_{\alpha}^{(-)} \equiv 0$$

$$p^{(-)} = -\eta_v \nabla_{\alpha} U_{\alpha}^*$$

$$p_{\alpha\beta}^{(-)} = -2\eta (\overline{\nabla_{\alpha} U_{\alpha}^*}).$$

If $T_{\alpha\beta}$ is a second order tensor, then $\overline{T}_{\alpha\beta} = \frac{1}{2}(T_{\alpha\beta} + T_{\beta\alpha}) - \frac{1}{3}T_{\gamma\gamma}\delta_{\alpha\beta}$. Thus $p_{\alpha\beta}^{(-)}$ is a symmetric tensor with zero trace. E^* , N^* and U^* are functions $(E, N, U)(\mathbf{r}) \mapsto E^*(\mathbf{r})$, $(E, N, U)(\mathbf{r}) \mapsto N^*(\mathbf{r})$, $(E, N, U) \mapsto U^*(\mathbf{r})$ of class C^2 . Thus, the values of E^* , N^* , U^* at $\mathbf{r} \in \Omega$ depend only on the values of E, N, U at the same $\mathbf{r} \in \Omega$. η_v , η and λ are functions $(E, N)(\mathbf{r}) \mapsto \zeta_v(\mathbf{r})$ etc. The values of η_v, η, λ at $\mathbf{r} \in \Omega$ depend also only on the values of E, N at the same $\mathbf{r} \in \Omega$. We shall assume moreover that $E^*(Jf) = E^*(f)$, $N^*(Jf) = N^*(f)$ and $U^*(Jf) = -U^*(f)$. The phenomenological quantities $Q_{\rm NSF}$ introduced by the Navier–Stokes–Fourier constitutive laws are thus $Q_{\rm NSF} = \{E^*, N^*, U^*, \eta_v, \eta, \lambda$ and the relations (4.1), (4.2), (4.3) introduced later in this Section}.

We shall review the general definition of DDS.

(DDS1) The simultaneous solutions of $R^{(+)}f = 0$ and $R^{(-)}f \mid_{R^{(+)}f=0} = 0$, denoted F, are physically interpreted as thermodynamically equilibrium states.

(4)

(DDS2) The states F found in (DDS1) are identical with the extremal points of a smooth function $\mathbb{V}: \mathscr{H} \times \Xi \to \mathbb{R}$, where Ξ is a two dimensional (in the case of onecomponent system) space, its elements are physically interpreted as thermodynamical fields. In the process of the solution of (DDS1), the components σ_E , σ_N of $\sigma \in \Xi$ appear as integration constants. The requirement that \mathbb{V} is invariant with respect to the orthogonal transformations in Ω (the systems considered are isotropic) implies $\mathbb{V}(Jf) = \mathbb{V}(f)$. The function \mathbb{V} evaluated at its extremal points (extremal with respect to $f \in \mathscr{H}$), denoted γ , is in general a multivalued function $\Xi \in \mathbb{R}$. For $\sigma \in \Xi$ for which γ is single-valued the function γ is identified with the thermodynamic potential.

(DDS3) Let F be a solution of (DDS1). Thermodynamic stability of F is equivalent to the local dynamic stability of F. This point is explained later in this Section.

One particular consequence of (DDS1), (DDS2) and (DDS3) is Onsager's symmetry of the vector field R linearized at F.

In the case of the time evolution equations (1) and (4) the equation $R^{(+)}f = 0$ takes the form

$$\nabla_{\alpha}(\eta_{v}\nabla_{\beta}U_{\beta}^{*}) + 2\nabla_{\beta}(\overline{\eta}\nabla_{\beta}U_{\alpha}^{*}) = 0$$
⁽⁵⁾

$$3\eta_{v}(\nabla_{\beta}U_{\beta}^{*})(\nabla_{\alpha}U_{\alpha}) + 2\eta(\overline{\nabla_{\beta}U_{\alpha}^{*}})(\nabla_{\alpha}U_{\beta}) + \nabla_{\alpha}(\lambda\nabla_{\alpha}E^{*}) = 0.$$
(6)

If we multiply (5) by $U_{\alpha}^{(*)}$ sum over α , integrate over Ω and integrate per parts, we have

$$\int_{\Omega} d^3 \mathbf{r} [(\nabla_{\alpha} U^*_{\alpha}) \eta_v (\nabla_{\beta} U^*_{\beta}) + 2(\overline{\nabla_{\beta} U^*_{\alpha}}) \eta(\overline{\nabla_{\beta} U^*_{\alpha}})] + s_U = 0.$$
(5.1)

We shall consider the boundary conditions that will guarantee that the surface integrals s_U equal to zero

$$s_{U} = 0. \tag{BC1}$$

We shall moreover assume that

$$\eta > 0 \tag{4.1}$$

$$\eta_v > 0$$

for all E, N considered.

As a consequence of (5.1), (BC1) and (4.1), we have

$$\nabla_{\beta} U_{\alpha}^* = 0$$

$$\nabla_{\alpha} U_{\alpha}^* = 0.$$
(7)

If we assume moreover that U^*_{α} on the boundary $\partial \Omega$ of Ω equals to zero

$$U^*_{\alpha} \big|_{\partial\Omega} = 0, \tag{BC2}$$

then (7) implies

$$U_a^* = 0. \tag{8}$$

From (8) we then have

$$U_{\alpha} = 0 \tag{8.1}$$

since $U_{\alpha} \to U_{\alpha}^*$ preserves the bundle structure $(\mathcal{H}, \pi^{(+)}, \mathcal{H}^{(+)})$. By inserting (7) into (6), one obtains

$$\nabla_{\alpha}(\lambda \nabla_{\alpha} E^*) = 0. \tag{6.1}$$

If (6.1) is multiplied by E^* , integrated over Ω and integrated per parts, one obtains

$$\int_{\Omega} d^3 \mathbf{r} (\nabla_{\alpha} E^*) \lambda (\nabla_{\alpha} E^*) + s_E = 0.$$
(6.2)

The boundary conditions that we assume will guarantee that the surface integral s_E equals to zero

$$s_E = 0. \tag{BC3}$$

Moreover, we assume that

$$\lambda > 0. \tag{4.2}$$

Then (6.2) implies

$$\nabla_{\sigma} E^* = 0 \tag{9}$$

and thus

$$E^* = \sigma_E, \tag{10}$$

where σ_E is a constant.

The equation $R^{(-)}f = 0$ restricted to the solutions of $R^{(+)}f = 0$ takes the form

$$V_{\alpha}N^* \mid_{E^* = \sigma_E} = 0 \tag{11}$$

and thus

 $N^* = \tilde{\sigma}_N,\tag{12}$

where $\tilde{\sigma}_N$ is a function of E^* . By σ_N we shall denote $\tilde{\sigma}_N$ restricted to $E^* = \sigma_E$.

Following the general theory of DDS, we look for a function $\mathbb{V}: \mathscr{H} \times \Xi \to \mathbb{R}$ (the elements of Ξ are σ_E and σ_N introduced in (10) and (12)) such that the solutions of (8), (10), (12) will be identical with extremal points of \mathbb{V} . There are, of course, many such functions. First of all we restrict ourselves to isotropic systems and thus $\mathbb{V}(Jf) = \mathbb{V}(f)$. The property (DDS3) defining the families of the dynamical systems DDS will restrict considerably the class of the functions \mathbb{V} that may be considered.

Let $\sigma_0 \equiv (\sigma_{E0}, \sigma_{N0}) \in \Xi$ be such that the equations (8), (10), (12) have a unique solution (denoted F_0). We assume that if $\sigma \in U_0$, where U_0 is a neighbourhood of σ_0 in Ξ , then the corresponding unique solutions of (8), (10), (12) form a submanifold \mathscr{F}_0 in \mathscr{H} . The equilibrium state F_0 is called thermodynamically stable if the thermodynamic potential γ is a convex function of $\sigma \in (\sigma_E, \sigma_N)$ at σ_0 . According to (DDS2) F_0 is an extremal point of the function \mathbb{V} and the thermodynamic potential γ defined on U_0 is identical with the function \mathbb{V} evaluated at F_0 . One can show [1] that the function γ is convex at σ_0 if and only if \mathbb{V} , as a function of $f \in \mathscr{H}$, is concave at σ_0 . Thus F_0 is thermodynamically stable if and only if $A_0 = -D^{(2)}\mathbb{V} \mid_{F_0}$ (the second derivative of \mathbb{V} with respect to $f \in \mathscr{H}$ evaluated at F_0) is a positive definite linear operator on $\tilde{H}_0 = T_{F_0} \mathscr{H} \ominus T_{F_0} \mathscr{F}_0$, where $T_{F_0} \mathscr{H}$ resp. $T_{F_0} \mathscr{F}_0$ is the tangent space to \mathscr{H} resp. \mathscr{F}_0 at F_0 .

Let P_0 denote the Hessian of the vector field R evaluated at F_0 . We say that F_0 is locally dynamically stable if there exists a unique solution to the initial value problem

$$\frac{\partial f}{\partial t} = P_0 f,\tag{13}$$

 $f \in \mathscr{D}(P_0) \subset H_0 \subset \widetilde{H}_0$, $f(t = 0) = f_0 \in \mathscr{D}(P_0)$, for all t > 0. If $||f_0||_0 < \infty$ then also $||f_t||_0 < \infty$; $f_t = T_t f_0$, where $\{T_t, t > 0\}$ is a strongly continuous semigroup of contraction operators. H_0 is an appropriate Hilbert space, its inner product is denoted $(.,.)_0$, its norm $|| = ||_0$.

We recall the Hille-Yoshida-Phillips [4] theorem. A necessary and sufficient condition for an operator P_0 to generate a strongly continuous semigroup of contraction operator on H_0 to itself is that P is closed and both P and its adjoint P^{\dagger} are dissipative operators. A linear operator P_0 with dense domain $\mathcal{D}(P) \subset H_0$ is said to be dissipative if

$$(P_0 f, f)_0 + (f, P_0 f)_0 \le 0$$

for all $f^{K} \mathcal{D}(P_{0})$.

We want to satisfy the necessary and sufficient condition in the Hille–Yoshida– Phillips theorem by

- (i) assuming (4.1), (4.2) already used in the solution of $R^{(+)}f = 0$ and $R^{(-)}f \mid_{R^{(+)}f=0} = 0$,
- (ii) choosing appropriately the Hilbert space H_0 .

Let \tilde{H}_0 be a fixed Hilbert space, say L_2 space, its inner product is denoted by (\cdot, \cdot) . We shall find, by using (4.1), (4.2), an operator A such that the necessary and sufficient condition in the Hille-Yoshida-Phillips theorem will be satisfied for AP_0 in \tilde{H}_0 . The boundary condition that will guarantee that AP_0 is closed will be formally denoted (BC4) (see, e.g., Reference [5]).

$$(AP_0 \text{ is closed}).$$

(BC4)

Let us suppose that such an operator A has been found. The operator A will be identified with $A_0 = -D^{(2)} \mathbb{V} |_{F_0}$. The assumption of thermodynamic stability of F_0 becomes equivalent to the assumption that A defines an inner product $(\cdot, A_{\cdot}) \equiv (\cdot, \cdot)_0$ in $\mathcal{D}(A) \subset H_0$ and subsequently in the Hilbert space $H_0 \subset \tilde{H}_0$. We shall see that the identification of A with $A_0 = -D^{(2)} \mathbb{V} |_{F_0}$ that is equivalent to (DDS3), restricts considerably the class of the functions \mathbb{V} that can be considered in the context of DDS.

Now, we shall find the operator A. Since $\mathbb{V}(Jf) = \mathbb{V}(f)$, we look for A in the form

$$A = \begin{pmatrix} a_1 & a_{12} & 0 & 0 & 0 \\ a_{12} & a_2 & 0 & 0 & 0 \\ 0 & 0 & a_3 & 0 & 0 \\ 0 & 0 & 0 & a_3 & 0 \\ 0 & 0 & 0 & 0 & a_3 \end{pmatrix},$$
 (14)

where a_1, a_2, a_3, a_{12} are real numbers (in the next Section (see (25) they will be differential operators). Thus, in this Section, H_0 and \tilde{H}_0 are identical ($\mathscr{D}(A) \equiv \tilde{H}_0$) except the difference in the definition of the inner product. The inequalities (4.1) and (4.2) enter only in $P^{(+)}(P^{(+)})$ is the Hessian of $R^{(+)}$ evaluated at F_0). Thus, we look for A such that $(AP_0)^{(+)}$ is selfadjoint (more generally, essentially selfadjoint) and dissipative and $(AP_0)^{(-)}$ is skewadjoint (more generally essentially skewadjoint). We assume the boundary conditions, denoted (BC5) that will guarantee the identity of the domains of AP and its adjoint (or a weaker requirement in the case of essential selfadjointness—see Reference [5]). By using the terminology introduced in [1], [2], we look for A such that AP_0 is an Onsager operator in \tilde{H}_0 or in other words P_0 is an Onsager operator in H_0 . [The Hilbert space H_0 is equipped with the inner product $(.,.)_A$ and with the fundamental decomposition $H_0 = H_0^{(+)} \oplus H_0^{(-)}$ that reflects locally the bundle structure of \mathscr{H} . If $\Pi^{(+)}$ resp. $\Pi^{(-)}$ are projection operators to the complete subspaces $H_0^{(+)}$ resp. $H_0^{(-)}$, then the operator $J = \Pi^{(+)} - \Pi^{(-)}$. One can define in H_0 an indefinite inner product $(., J.)_A$ and H_0 becomes a Krein space [6]. We say that P_0 is an Onsager operator in H_0 if P_0 is selfadjoint with respect to the inner product $(., .)_A$.]

From (1), (3), (4), we have (the notation $E_E^* \equiv \partial E^* / \partial E |_{F_0}$ etc. is used);

where

$$\begin{split} P_{02}^{(+)} &= \begin{pmatrix} (P_{02}^{(+)})_{11} & (P_{02}^{(+)})_{12} & (P_{02}^{(+)})_{13} \\ (P_{02}^{(+)})_{12} & (P_{02}^{(+)})_{22} & (P_{02}^{(+)})_{23} \\ (P_{02}^{(+)})_{13} & (P_{02}^{(+)})_{23} & (P_{02}^{(+)})_{33} \end{pmatrix} \\ (P_{02}^{(+)})_{ij} &= \eta_0 \, \delta_{ij} \Delta + (\eta_{v0} + \frac{1}{3}\eta_0) \nabla_i \nabla_j, \\ \Delta &= \nabla_i \nabla_i, \quad \lambda_0 &= \lambda \mid_{F_0}, \quad \eta_{v0} = \eta_v \mid_{F_0}, \quad \eta_0 = \eta \mid_{F_0}. \end{split}$$

Thus, if we make the identifications

$$a_{1} = -E_{E}^{*}, \qquad a_{12} = -E_{N}^{*}$$

$$a_{3} = -U_{U_{1}}^{*} = -U_{U_{2}}^{*} = -U_{U_{3}}^{*} = -U_{U}^{*},$$
(16)

then $(AP_0)^{(+)}$ (using (4, 1), (4, 2) and the boundary conditions (BC5)) is selfadjoint and dissipative.

From (1), (3), (4), we obtain also

$$P_{0}^{(-)} = \begin{pmatrix} 0 & 0 & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{2} & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{3} \\ 0 & 0 & -N_{0} \nabla_{1} & \cdot & -N_{0} \nabla_{2} & -N_{0} \nabla_{3} \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{1} & 0 & 0 & 0 \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{2} & 0 & 0 & 0 \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{2} & 0 & 0 & 0 \end{pmatrix}$$
(17)

where

$$N_0 = N \big|_{F_0}, N_0^* = N^* \big|_{F_0}, N_E^* = \frac{\partial N^*}{\partial E} \bigg|_{F_0}, N_N^* = \frac{\partial N^*}{\partial N} \bigg|_{F_0}$$

By multiplying (17) from the left by A, taking into account (16), one obtains the following conditions which have to be satisfied for $AP_0^{(-)}$ to be skewadjoint:

$$\frac{1}{N_0} U_U^* N_E^* = \frac{1}{N_0} E_E^* N_0^* + N_0 E_N^*$$

$$\frac{1}{N_0} U_U^* N_N^* = \frac{1}{N_0} E_N^* N_0^* - N_0 a_2$$
(18)

If we choose the function V such that the following relations

$$\frac{\partial \mathbb{V}}{\partial E} = E^* - \sigma_E$$

$$\frac{\partial \mathbb{V}}{\partial N} = -\frac{1}{N_2} (N^* E^* - \sigma_N)$$

$$\frac{\partial \mathbb{V}}{\partial U_{\alpha}} = -E^* U_{\alpha}$$
(19)

are satisfied then (18) is satisfied and also the solutions of (8), (10), (12) are identical with the extremal points of \mathbb{V} . The equations (19) imply that the functions E^* , N^* entering the Navier-Stokes-Fourier constitutive laws are not independent. The relation between them (the generalized Maxwell's relation) is implied by the symmetry of $D^{(2)}\mathbb{V}$

$$(D^{(2)}\mathbb{V} \text{ is symmetric})$$
 (4.3)

The general solution of (19) is

$$\mathbb{V}(E, N, \mathbf{U}; \sigma_E, \sigma_N) = \int_{\Omega} d^3 \mathbf{r} \left[S(E(\mathbf{r}), N(\mathbf{r}), \mathbf{U}(\mathbf{r})) - \sigma_E E - \frac{1}{N} \sigma_N \right], \tag{20}$$

400

where

$$\frac{\partial S}{\partial E} = E^*, \ \frac{\partial S}{\partial (1/N)} = E^*N^* \text{ and } \frac{\partial S}{\partial U_{\alpha}} = -E^*U_{\alpha}.$$

This function V is identical with the non-equilibrium extension of the thermodynamic potential postulated in non-equilibrium thermodynamics [7]. We have obtained this function V as the consequence of the postulated Navier–Stokes–Fourier constitutive laws and the requirement that the resulting family of the time evolution equations possesses the formal structure of DDS.

III. Higher order Navier-Stokes-Fourier fluid dynamics

We shall consider the following generalization of the Navier–Stokes–Fourier constitutive laws:

$$P_{\alpha\beta}^{(+)} = \mathcal{N}^* \delta_{\alpha\beta} + p_{\alpha\beta}^{(+)}$$

$$P_{\alpha\beta}^{(-)} = p^{(-)} \delta_{\alpha\beta} + p_{\alpha\beta}^{(-)}$$

$$Q_{\alpha}^{(+)} = -\lambda \nabla_{\alpha} \mathscr{E}^*$$

$$Q_{\alpha}^{(-)} = \mu_v \nabla_{\alpha} (\nabla_{\beta} U_{\beta}^*) + \mu \Delta U_{\alpha}^*$$

$$p_{\alpha\beta}^{(+)} = \overline{\frac{3}{2} \nabla_{\alpha} (\Lambda \nabla_{\beta} E^*)}$$

$$p^{(-)} = -\eta_v \nabla_{\alpha} U_{\alpha}^*$$

$$p_{\alpha\beta}^{(-)} = \overline{-2\eta \nabla_{\beta} U_{\alpha}^*}.$$

Let $j(E, N) \equiv (\Delta E, \Delta N, \nabla_{\alpha} E \nabla_{\alpha} E, \nabla_{\alpha} N \nabla_{\alpha} N, \nabla_{\alpha} E \nabla_{\alpha} N)$. The functions \mathscr{E}^* and \mathscr{N}^* are defined as follows: \mathscr{E}^* : $(E, N, j(E, N), \mathbf{U})(\mathbf{r}) \mapsto \mathscr{E}^*(\mathbf{r})$ and $\mathscr{E}^*(\mathbf{r}) |_{j(E, N) = 0} = E^*(\mathbf{r})$ introduced in (4). Similarly \mathscr{N}^* : $(E, N, j(E, N), \mathbf{U})(\mathbf{r}) \mapsto \mathscr{N}^*(\mathbf{r})$ and $\mathscr{N}^*(\mathbf{r}) |_{j(E, N) = 0} =$ $N^*(\mathbf{r})$ introduced in (4). We assume moreover that $\mathscr{E}^*(Jf) = \mathscr{E}^*(f)$ and $\mathscr{N}^*(Jf) =$ $\mathscr{N}^*(f)$. U* is assumed to be the same as in (4). $\Lambda, \mu_v, \mu, \lambda, \eta_v, \eta$ are real valued functions of (E, N, j(E, N)) (e.g., Λ : $(E, N, j(E, N))(\mathbf{r}) \to \Lambda(\mathbf{r})$ etc.). The phenomenological quantities Q_{GNSF} introduced by the Generalized Navier–Stokes–Fourier constitutive laws are thus $Q_{\text{GNSF}} \equiv {\mathscr{E}^*, \mathscr{N}^*, U^*, \Lambda, \mu_v, \mu, \lambda, \eta_v, \eta}$ and the relations (21.1), (21.2), (21.3) introduced later in this Section}.

If (21) is inserted into (1) then clearly the system of partial differential equations of the order higher than two is obtained. In the constitutive laws (21) the assumption, used in (4), that $P_{\alpha\beta}$ and Q_{α} at r depend only on the values of E, N, U at the same r has been relaxed. The non-locality of the dependence of $P_{\alpha\beta}$ and Q_{α} on (E, N, U) is expressed in (21) by letting the quantities entering (4) to depend also on the gradients of E and N.

The discussion of (1), (21) follows closely the discussion of (1), (4) in Section II. The equation $R^{(+)}f = 0$ remains the same as in (5)–(10). We assume again that

$$\eta > 0, \eta_v > 0 \tag{21.1}$$

$$\lambda > 0 \tag{21.2}$$

for all (E, N, j(E, N)) considered and the boundary conditions (GBC1)-(GBC3) corresponding to (BC1)-(BC3) in Section II. The equation (10) reads now

$$\mathscr{E}^* = \sigma_E$$

(22)

(21)

The equation $R^{(-)}f = 0$ restricted to the solutions of $R^{(+)}f = 0$ takes the form

$$\mathcal{N}^* \mid_{\mathcal{B}^* = \sigma_F} = 0 \tag{23}$$

and thus

$$\mathscr{N}^* = \widetilde{\sigma}_N \tag{24}$$

where $\tilde{\sigma}_N$ is a function of \mathscr{E}^* . By σ_N we shall denote again $\tilde{\sigma}_N$ restricted to $\mathscr{E}^* = \sigma_E$. The problem is now to find the function $\mathscr{V}: (E, N, \mathbf{U}, j(E, N); \sigma_E, \sigma_N) \rightarrow \mathscr{V}(E, N, \mathbf{U}, j(E, N); \sigma_E, \sigma_N)$ such that (i) the solutions of (8), (22), (24) will be identical with the extremal points of \mathscr{V} , (ii) thermodynamic stability of a thermodynamically equilibrium state will be equivalent to its local dynamic stability, (iii) if $j(E, N) \equiv 0$ then \mathscr{V} reduces to \mathbb{V} that has been introduced in (20). We shall indeed find an example of such a function \mathscr{V} . It will become evident however that the requirement (DDS3) – the equivalence of thermodynamic and local dynamic stability – is not strong enough to single out the function \mathscr{V} . A systematic study of the higher order fluid dynamic requires a generalization of (DDS3). The local dynamic stability has to be replaced by a type of global dynamic stability (the requirement that \mathscr{V} is a Liapunov function).

Let F_0 be a solution of (8), (22), (24) (thus also an extremal point of the function \mathscr{V}) and let moreover F_0 does not depend on **r**. Similarly as in Section II, we shall find the operator A such that AP is an Onsager operator. The operator A will be then identified with $-D^{(2)}\mathscr{V}|_{F_0}$. Since F_0 is independent of **r** and \mathscr{V} can depend only on $(E, N, \mathbf{U}, j(E, N))$ the operator A has to have the form

$$A = A^{(0)} + A^{(1)}\Delta, \tag{25}$$

where $A^{(0)}$ is a 5 \times 5 matrix of real numbers identical with (14) and

	b_1	b_{12}	0	0	0		
	b_{12}	b_2	0	0	0	и и ₁	
$A^{(1)} =$	0	0	b^3	0	0	(26)
	0	0	0	b_3	0		
	0	0	0	0	b_3		

where b_1, b_2, b_{12}, b_3 are real numbers. Since the operator A is a differential operator, i.e., A is unbounded operator in \tilde{H}_0 , $(\mathcal{D}(A) \neq H_0)$ the Hilbert space $H_0 \neq \tilde{H}_0$. Construction of $H_0 \subset \tilde{H}_0$ in which the given operator A defined in $\mathcal{D}(A) \subset \tilde{H}_0$ is bounded and can thus be used, if moreover A is positive definite, to define an inner product in H_0 is explained for example in Reference [8].

From (1), (3), (21), we have

where $P_{02}^{(+)}$ is the same as in (15), $\lambda_0 = \lambda |_{F_0}$, E_E^* , E_N^* , a_{12} , a_2 are the same as in (15) and \mathscr{E}_{2E}^* is the part of the $\partial \mathscr{E}^* / \partial E |_{F_0}$ that is proportional to Δ . Thus if a_1 , a_2 , a_{12} are the same as in (16),

$$b_{1} = -\mathscr{E}_{2E}^{*}$$

$$b_{12} = -\mathscr{E}_{2E}^{*},$$

$$b_{3} = 0.$$
(28)

and appropriate boundary conditions (GBC4), (GBC5) that correspond to (BC4), (BC5) are assumed, then $(AP_0)^{(+)}$ is selfadjoint and dissipative.

From (1), (3), (21), we obtain also

$$P_{0}^{(-)} = \begin{pmatrix} 0 & 0 & 0 & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{2} & -\frac{1}{N_{0}} N_{0}^{*} \nabla_{3} \\ 0 & 0 & 0 & -N_{0} \nabla_{1} & -N_{0} \nabla_{2} & -N_{0} \nabla_{3} \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{1} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{1} & 0 & 0 & 0 \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{2} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{2} & 0 & 0 & 0 \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{3} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{3} & 0 & 0 & 0 \\ -\frac{1}{N_{0}} N_{E}^{*} \nabla_{3} & -\frac{1}{N_{0}} N_{N}^{*} \nabla_{3} & 0 & 0 & 0 \\ \end{pmatrix} \begin{pmatrix} 0 & 0 & p_{1} \nabla_{1} & p_{2} \nabla_{2} & p_{3} \nabla_{3} \\ 0 & 0 & 0 & 0 & 0 \\ p_{2} \nabla_{1} & p_{3} \nabla_{1} & 0 & 0 & 0 \\ p_{2} \nabla_{2} & p_{3} \nabla_{2} & 0 & 0 & 0 \\ p_{2} \nabla_{2} & p_{3} \nabla_{3} & 0 & 0 & 0 \end{pmatrix} \Delta,$$
(29)

where

$$p_{1} = -\frac{1}{N_{0}} (\mu_{v_{0}} + \mu_{0}), \qquad p_{2} = -\frac{1}{N_{0}} (\mathcal{N}_{2E}^{*} + \Lambda E_{E}^{*})$$

$$p_{3} = -\frac{1}{N_{0}} (\mathcal{N}_{2N}^{*} + \Lambda E_{N}^{*}).$$
(30)

The first term on the right hand side of (29) is identical with (17) since, according to our assumption, F_0 is independent of **r** and thus $j(E, N) |_{F_0} \equiv 0$. By multiplying (27) from the left by A (see (25), (26)), taking into account the boundary conditions

(GBC4), (GBC5), and considering only the derivatives up to and including the fourth order, one obtains the following conditions that have to be satisfied for $(AP_0)^{(-)}$ to be skewadjoint:

$$a_{1}p_{1} + b_{1}\left(-\frac{1}{N_{0}}N_{0}^{*}\right) + b_{12}(-N_{0}) = a_{3}p_{2}$$

$$a_{12}p_{1} + b_{12}\left(-\frac{1}{N_{0}}N_{0}^{*}\right) + b_{2}(-N_{0}) = a_{3}p_{3}.$$
(31)

By inserting (28), (16) and (30) into (31), we have

$$\frac{1}{N_0} E_E^*(\mu_{v_0} + \mu_0) + \frac{1}{N_0} \mathscr{E}_{2E}^* N_0^* + N_0 \mathscr{E}_{2N}^* = +\frac{1}{N_0} E_0^*(\mathscr{N}_{2E}^* + \Lambda E_E^*)$$

$$\frac{1}{N_0} E_N^*(\mu_{v_0} + \mu_0) + \frac{1}{N_0} \mathscr{E}_{2N}^* N_0^* + N_0 b_2 = -\frac{1}{N_0} E_0^*(\mathscr{N}_{2N}^* + \Lambda E_N^*).$$
(32)

If the function \mathscr{V} satisfies the following relations:

$$\frac{\delta_v \mathscr{V}}{\delta_v E} = \mathscr{E}^* - \sigma_E$$

$$\frac{\delta_v \mathscr{V}}{\delta_v N} = -\frac{1}{N^2} (\mathscr{E}^* \mathscr{N}^* + \nabla_\alpha (\Lambda \nabla_\alpha E^*) + (\mu_v + \mu) \Delta E^* - \sigma_N)$$

$$\frac{\delta_v \mathscr{V}}{\delta_v U_\alpha} = -\mathscr{E}^* U_\alpha,$$
(33)

then (32) is satisfied, the extremal points of \mathscr{V} are identical with the solutions of (8), (22), (24) and \mathscr{V} reduces to \mathbb{V} introduced in (20) if $j(E, N) \equiv 0$. The differentiation

$$\frac{\delta_v \mathscr{V}}{\delta_v \phi} \equiv \frac{\delta \mathscr{V}}{\delta \phi} - \nabla_\alpha \left(\frac{\delta \mathscr{V}}{\delta \nabla_\alpha \phi} \right) + \Delta \left(\frac{\delta \mathscr{V}}{\delta \Delta \phi} \right)$$

is the variational differentiation, $\phi \equiv E$ or $\phi \equiv N$. The quantities \mathscr{E}^* , \mathscr{N}^* , μ_v , μ , Λ are not independent. The relation among them is implied by the generalized Maxwell's relations

$$(D^{(2)}\mathscr{V} \text{ is symmetric}).$$
 (21.3)

A function \mathscr{V} satisfying (33) is

$$\mathscr{V}(E, N, \mathbf{U}, j(E, N); \sigma_E, \sigma_N) = \int_{\Omega} d^3 \mathbf{r} \left[\mathscr{S}(E(\mathbf{r}), N(\mathbf{r}), \mathbf{U}(\mathbf{r}), j(E, N)(\mathbf{r})) - \sigma_E E - \frac{1}{N} \sigma_N \right], \quad (34)$$

$$\begin{split} & \mathcal{S}^{*} \mid_{j(E,N) \equiv 0} = \mathcal{S}^{*} \text{ Introduced in (20),} \\ & \frac{\delta_{v} \mathscr{S}}{\delta_{v} E} = \mathscr{E}^{*}, \\ & \frac{\delta_{v} \mathscr{S}}{\delta_{v} \left(\frac{1}{N}\right)} = \mathscr{E}^{*} \mathscr{N}^{*} + \nabla_{\alpha} (\Lambda \nabla_{\alpha} E^{*}) + (\mu_{v} + \mu) \Delta E^{*}, \\ & \frac{\delta_{v} \mathscr{S}}{\delta_{v} U_{\alpha}} = -\mathscr{E}^{*} U_{\alpha}. \end{split}$$

The thermodynamic potential $\gamma_{\mathscr{V}}$ constructed from \mathscr{V} is in general different from the thermodynamic potential $\gamma_{\mathscr{V}}$ constructed from \mathbb{V} . In particular, a thermodynamic equilibrium state F_0 can be thermodynamically stable with respect to $\gamma_{\mathscr{V}}$ but thermodynamically unstable with respect to $\gamma_{\mathscr{V}}$. There is a possibility of relating these different notions of thermodynamic stability to the appearance of the metastable states known from experience with fluids. This problem will be hopefully discussed in detail in a subsequent paper.

We have found that if the constitutive laws (21) are postulated and the nonequilibrium extension of the thermodynamic potential is given by (34) then the linearized infinitesimal generator of the time evolution possesses the Onsager symmetry. The non-equilibrium thermodynamic potential \mathscr{V} is not however determined uniquely by studying the equivalence of the thermodynamic and local dynamic stability. We cannot decide for example if in (34) should be the term $\nabla_{\alpha}(\Delta \nabla_{\alpha} E^*)$ or the term $\Lambda \Delta E^*$. A systematic study of higher order fluid dynamics will require the study of a type of global dynamic stability where the function \mathscr{V} will play the role of a Liapunov function.

IV. Discussion

An example of the higher order fluid dynamics possessing the structure of the family of dissipative dynamical systems of macroscopic physics [1], [2] has been constructed.

The non-equilibrium extension of the thermodynamic potential cannot be obtained in higher order fluid dynamics by using the method used in non-equilibrium thermodynamics [7] (i.e., S(E, N) is extended to $\int_{\Omega} d^3 \mathbf{r} S(E(\mathbf{r}), N(\mathbf{r}))$, where E, N are the thermodynamical conjugates of the thermodynamic fields σ_E and σ_N , $E(\mathbf{r})$ and $N(\mathbf{r})$ are the fluid mechanics state variables, called the assumption of the local thermodynamic equilibrium). The situation in higher order fluid mechanics is similar, as for example in the Boltzmann kinetic theory. The non-equilibrium extension of the thermodynamic potential, so called Boltzmann's H-function, cannot be also obtained by using the method used in non-equilibrium thermodynamics. We have seen that the method associated with the theory of DDS applies in the Navier–Stokes–Fourier fluid dynamics (Section II) kinetic theory [1] and higher order fluid dynamics (Section III). The higher order fluid dynamics indicates however the necessity of generalizing the point (DDS3) in the definition of the formal structure of DDS. A type of global–Liapunov–dynamic stability has to replace the local, Hille–Yoshida–Phillips–dynamic stability.

Several different approaches to higher order fluid dynamics have been developed in fluid mechanics. We mention three of them.

In the point of view advocated by Coleman [3] the inequality $\partial S/\partial t + J \ge 0$ is added to the system of equations (1). J is a known function of $P_{\alpha\beta}$, Q_{α} ; S is another, in addition to $P_{\alpha\beta}$ and Q_{α} , unknown function of fluid mechanics state variables. The higher order fluid dynamics in the framework of Coleman's point of view has been discussed recently by de Sobrino [9].

Reduction of the Boltzmann kinetic theory to fluid dynamics inspired the Burnett [10] [11] and the Waldmann [12] [13] approach to higher order fluid dynamics. The Burnett approach has been inspired by the iterations of the Chapman– Enskog method, the Waldmann approach by Grad's thirteen moment method of solution of Boltzmann's equation. The linearized Burnett equations together with the non-equilibrium thermodynamic potential used in non-equilibrium thermodynamics do not possess the Onsager symmetry [11]. Waldmann has suggested the generalization of the entropy production and the entropy flux that are appropriate for the constitutive laws he used (e.g., the entropy production is evidently positive definite) but the generalization of the non-equilibrium entropy itself, that has to depend on gradients, has not been discussed.

REFERENCES

- [1] M. GRMELA, Helv. Phys. Acta 47, 667 (1974); J. Math. Phys. 16, 2441 (1975).
- [2] M. GRMELA and I. ISCOE, Ann. Inst. H. Poincaré, Sect. A. (submitted).
- [3] C. TRUESDELL, Rational Thermodynamics (McGraw-Hill, New York 1969).
- [4] R. S. PHILLIPS, Trans. Amer. Math. Soc. 90, 193 (1959).
- [5] G. HELLWIG, Differential Operators of Mathematical Physics (Addison-Wesley, Reading, Mass. 1967).
- [6] J. BOGNÁR, Indefinite Inner Product Spaces (Springer-Verlag, Heidelburg, New York 1974).
- [7] S. R. DE GROOT and P. MAZUR, Non-Equilibrium Thermodynamics (North-Holland, Amsterdam 1962).
- [8] J.-P. AUBIN, Applied Functional Analysis, Wiley-Interscience (to appear).
- [9] L. DE SOBRINO, Can. J. Phys. 54, 105 (1976).
- [10] D. BURNETT, Proc. London Math. Soc. 40, 382 (1935).
 S. CHAPMAN and T. G. COWLING, The Mathematical Theory of Non-Uniform Gases (Cambridge U.P., Cambridge, England 1953).
- J. A. MCLENNAN, Phys. Rev. A8, 1479 (1973).
- [11] J. A. MCLENNAN, Phys. Rev. A10, 1272 (1974).
- [12] L. WALDMANN, Z. Naturforsch. 22a, 1269 (1967).
- [13] L. WALDMANN and H. VESTNER, Physica 80A, 523 (1975).