

Equation of hydrostatic equilibrium and temperature dependent gravitational constant

Autor(en): **Massa, Corrado**

Objekttyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **62 (1989)**

Heft 4

PDF erstellt am: **24.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116040>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Equation of hydrostatic equilibrium and temperature dependent gravitational constant

By Corrado Massa

Via Fratelli Manfredi 55 42100–Reggio Emilia Italy

(21. XII. 1988 revised, 7.II.1989)

Abstract. The Oppenheimer–Volkoff (shortly OV) equation of hydrostatic equilibrium describes the balance between gravitational force and pressure gradient in a self-gravitating perfect fluid. In the following, a generalized form of the OV equation is obtained by the assumption that the gravitational constant G is temperature-dependent according to the law $G = G_0(1 - bT^2)^{-1}$ where b is a positive constant. Such a law is required by a variety of gauge theories used in the unification program. A direct consequence of the generalized OV equation is the existence of an upper bound $T < (2/b)^{1/2}$ on the temperature of any self-gravitating radiation field.

A variety of gauge theories used in the unification program imply a temperature-dependent gravitational constant G given by:

$$G = G_0(1 - bT^2)^{-1} \quad (1)$$

where T is the temperature, G_0 is the zero temperature value of G , very close to the currently observed value $6.67 \cdot 10^{-8} \text{ cm}^{-3} \text{ g}^{-1} \text{ s}^{-2}$, and b is a positive constant whose numerical value depends on the details of the theory concerned [1–3]. The simplest way to fit the law (1) on the framework of Einstein's gravitational theory is to write the gravitational field equations [1, 4]

$$R^{ik} - (\frac{1}{2})g^{ik}R = 8\pi GT^{ik} \quad (2)$$

where $G = G(T)$ according to equation (1). R^{ik} is the Ricci tensor, g_{ik} is the fundamental tensor, $R = g^{ik}R_{ik}$ is the spacetime curvature scalar, and T^{ik} is the energy tensor. In this paper, latin indices run from 0 to 3, greek indices run from 1 to 3; commas denote partial derivative and semicolons denote covariant derivative with respect to the spacetime coordinates x^i ; dots denote differentiation with respect to x^0 , and apices denote differentiation with respect to x^1 .

By equation (2) and by the Bianchi identity $R_{i;k}^k = (\frac{1}{2})R_{,i}$ we have $(GT^{ik})_{;k} = 0$, namely

$$\dot{G}T^{00} + Gt^0 = 0 \quad (3)$$

$$G'T^{11} + Gt^1 = 0 \quad (3a)$$

where $t^i = T_{;k}^{ik}$.

Consider now a self-gravitating mass described by the energy tensor of a perfect fluid:

$$T^{ik} = (\rho + P)U^i U^k - Pg^{ik} \quad (4)$$

where ρ is the rest frame mass density and P the scalar pressure; U^i is the 4-velocity of the local rest frame of the fluid.

Consider a static, spherically symmetric field; thus:

$$U^\beta = 0, \quad ds^2 = e^\nu(c dt)^2 - e^\lambda dr^2 - d\Omega^2 \quad (5)$$

where ds^2 is the spacetime line-element, ν and λ are functions of r only, and $d\Omega^2$ means $(r d\vartheta)^2 + (r \sin \vartheta d\varphi)^2$ where $(ct, r, \vartheta, \varphi) = x^i$ are the Schwarzschild coordinates (r is the radial coordinate).

The normalization of 4-velocity $U^i U_i = 1$ determines $U^0 = e^{-\nu/2}$, and, by (4)(5), $T^{00} = \rho e^{-\nu}$, $T_0^0 = \rho$, $T^{11} = Pe^{-\lambda}$. For $\dot{G} = 0$, equation (3) gives $t^0 = 0$ (neither creation nor destruction of matter) and we have to consider equation (3b) only. Equation (3b) with $T^{00} = \rho e^{-\nu}$ and $T^{11} = Pe^{-\lambda}$ gives:

$$(G'/G) + P' + (\frac{1}{2})(\rho + P)\nu' = 0 \quad (6)$$

The (00) component of the field equation (2) in the proper reference frames of fluid elements is;

$$(1/r^2)(d/dr)[r(1 - e^{-\lambda})] = 8\pi G\rho,$$

which integrated gives;

$$\left(\frac{1}{4\pi}\right)m(r) = \int_0^r G\rho r^2 dr \quad (7)$$

where $2m(r) = r(1 - e^{-\lambda})$.

The (11) component is

$$-(1/r^2) + (1/r^2)e^{-\lambda} + (1/r)e^{-\lambda}\nu' = 8\pi GP,$$

that gives:

$$(\frac{1}{2})\nu' = (m + 4\pi PG r^3)(r^2 - 2mr)^{-1} \quad (8)$$

Equations (1) (8) lead to;

$$P' = -(\rho + P)(m + 4\pi PG r^3)(r^2 - 2mr)^{-1} - PG'/G \quad (9)$$

with $m = m(r)$ given by equation (7). Equation (9), with $G = G_0 = \text{constant}$, reduces to the usual OV equation for a perfect fluid [5-7]. Consider now equation (1), which gives $G'/G = (T'/T)2bT^2(1 - bT^2)^{-1}$. If the fluid is a radiation field with the equation of state $\rho = 3P = aT^4$ (a = the radiation constant) then equation (9) reads:

$$AT'/T + [Bm + (4\pi/3)aT^4 r^3 G_0](BC)^{-1} = 0, \quad (10)$$

where;

$$A = 1 + (\frac{1}{2})bT^2 B^{-1}, \quad (10a)$$

$$B = 1 - bT^2, \quad (10b)$$

$$C = r^2 - 2mr. \quad (10c)$$

A direct consequence of equation (10) is the existence of an upper bound on the temperature T of the fluid. To see this fact, assume $T > 1/\sqrt{b}$ at some point r_0 inside the fluid, namely $B < 0$. For equation (1), the equation (7) gives $m(r_0) < 0$, viz. (for equation (10c)) $C > 0$, and equation (10) leads to

$$AT' > 0 \quad (11)$$

T' is certainly negative. Indeed, if $T' > 0$, then $A > 0$, and (for equations (10,a,b) and for $B < 0$) $T > (2/b)^{1/2}$; all values between $(1/b)^{1/2}$ and $(2/b)^{1/2}$ would be forbidden, and T could not be continuous.

Put $T' < 0$ into equation (11) and obtain $A < 0$, viz. $T < (2/b)^{1/2}$ that is an upper bound on T .

A more severe bound, namely $T < (1/b)^{1/2}$, can be obtained via the quite different procedure described in Reference (1). In the special case of the Hawking black hole temperature, the bound $T < (1/b)^{1/2}$ can be derived by the very simple procedure of Reference (2). My thanks to an anonymous referee of this Review which brought Reference (3) to my attention.

REFERENCES

- [1] C. MASSA, Helv. Phys. Acta (1989) to be published, and references cited therein.
- [2] P. C. W. DAVIES, Phys. Lett. *101B* 399 (1981).
- [3] P. C. W. DAVIES, "Temperature-Dependent G and Black Hole Thermodynamics", in the Proceedings of the 2nd Moscow Quantum Gravity Seminar (ed. Markov & West, Plenum 1984).
- [4] A. K. RAYCHAUDHURI, and B. BAGCHI, Phys. Lett. *124B*, 168 (1983).
- [5] J. R. OPPENHEIMER, and G. VOLKOFF, Phys. Rev. *55*, 374 (1939).
- [6] C. W. MISNER, K. S. THORNE, and J. A. WHEELER, Gravitation (W. H. Freeman and Company, San Francisco 1973) Chapter 23, Section 5, p. 605.
- [7] R. D. SORKIN, R. M. WALD, and Z. Z. JIU, Gen. Rel. Grav. *13*, 1127 (1981).