

# **Speeding-up simulations of interacting systems near critical points by the multigrid Monte Carlo method**

Autor(en): **Stoll, E.P.**

Objekttyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **62 (1989)**

Heft 6-7

PDF erstellt am: **24.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116171>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

SPEEDING-UP SIMULATIONS OF INTERACTING SYSTEMS NEAR CRITICAL POINTS  
BY THE MULTIGRID MONTE CARLO METHOD

E.P. Stoll, IBM Research Division, Zurich Research Laboratory, CH-8803 RÜSCHLIKON, Switzerland

**Abstract:** With multigrid coarse-to-fine transformations, loss of computer performance due to very slow long-wavelength relaxations can be eliminated. For systems with short-range interactions, effective interactions can however be modified according to renormalization-group considerations. With coarse-to-fine grid transformations alone, deviations indicate whether fine-to-coarse transformations are also necessary.

### 1. Introduction

Computer simulations have become very effective for many problems which cannot be treated by analytic methods. However, in complex systems, very slow relaxations at long wavelength restrict the application of vector and parallel computers. For the numerical solution of partial differential equations, the multigrid method [1] has demonstrated how it is possible to speed up slow long-wavelength relaxations by many orders of magnitude. A condition for the solution of such systems by this method is their selfsimilarity, i.e. the couplings between grid points remain unchanged in the mapping procedure from fine-to-coarse grids. For equilibrium systems with short-range interactions, renormalization-group theory [2] shows that effective interactions have to be recalculated at each fine-to-coarse mapping except at fixed points corresponding to critical points.

### 2. Multigrid theory

Multigrid procedures [1] usually start with fine grids, iterate, transform to coarser grids, iterate again and return to the fine grid. This algorithm can be extended to several levels. For a short-range interaction, the interaction has to be renormalized as demonstrated for the Ising model at  $T > T_c$  in [3]. The mapping from fine-to-coarse grids specifies the "melting" of blockspins on the fine grid. In our simulations, large blockspins with renormalized interactions were used at the outset and were allowed to melt only during successive iterations. The entropy was kept constant in each melting step.

### 3. Results

Figure 1 shows thermodynamic quantities for the two-dimensional Ising model normalized to exact values [4-6] computed with the multigrid Monte Carlo method. Deviations from

1 close to the critical point indicate that if the information of additional multigrid levels is used, blockspin melting at constant entropy is by itself unsatisfactory. Therefore, fine-to-coarse mappings are also needed, in which renormalized interactions can be estimated as demonstrated in [3].




Figure 1: Plot of specific heat  $c_V$ , energy  $E$ , susceptibility  $\chi$  and order parameter  $m$  normalized to their exact theoretical values [4-6] for two-dimensional Ising models with  $1024 \times 1024$  and  $4096 \times 4096$  spins.

#### 4. Conclusions

Renormalization-group multigrid Monte Carlo calculations are very useful to improve the performance of numerically intensive calculations. Interactions have to be renormalized if the fine-to-coarse decimation step cannot be treated via a mapping to a percolation problem [7]. Fine-to-coarse mappings have to be combined with coarse-to-fine mappings for adequate blockspin melting.

#### 5. References

- [1] W.L. Briggs, *A Multigrid Tutorial* (Philadelphia: Society for Industrial and Applied Mathematics, 1987)
- [2] K.G. Wilson, *Rev. Mod. Phys.* **47**, 773 (1975)
- [3] H.H. Hahn and T.S.J. Streit, *Physica A* **154**, 108 (1988)
- [4] C.N. Yang, *Phys. Rev.* **85**, 808 (1952)
- [5] J.W. Essam and M.E. Fisher, *J. Chem. Phys.* **38**, 802 (1963)
- [6] M.F. Sykes, D.S. Gaunt, P.D. Roberts and J.A. Wyles, *J. Phys. A: Math. Gen.* **5**, 624 (1972)
- [7] D. Kandel, E. Domany, D. Ron, A. Brandt and E. Loh Jr., *Phys. Rev. Lett.* **60**, 1591 (1988)