

Erratum on the paper Helv. Phys. Acta 65 (1992) 748 : semiclassical expansions of the thermodynamic limit for a Schrödinger equation. II, The double well case

Autor(en): **Helffer, Bernard**

Objekttyp: **Corrections**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **67 (1994)**

Heft 1

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Erratum on the Paper

Helv.Phys.Acta 65 (1992) 748:

**Semiclassical expansions of the thermodynamic limit
for a Schrödinger equation**

II. The double well case

By Bernard Helffer

DMI-ENS
45 rue d'Ulm
F-75230 Paris Cédex

and Johannes Sjöstrand

Département de Mathématiques
Université Paris-Sud
F-91405 Orsay
(2. II. 1994)

In our paper there is a gap in the proof of (3.7) p.761. It is indeed not always possible to construct κ with the property that $\kappa(x^\Omega) = x^{\Omega_+}$ is the center of the ℓ^∞ -ball containing one well and satisfying (3.7). We overlooked the rôle of $\Sigma_k(x_k - x_{k+1})^2$ in the comparison of V and $V \circ \kappa$. The text starting from p.760 line -8 to p.761 line -9 has to be modified as follows.

Let $\kappa : \Omega \rightarrow \Omega_+ = (I_+)^m$ be the translation with $\kappa(x^\Omega) = \tilde{x}^{\Omega_+}$. Here $\tilde{x}_j^{\Omega_+} = 0$, if I_j contains t_0 and in the other case $\tilde{x}_j^{\Omega_+} = \pm \rho \delta_0$ for a suitable ρ independent of Ω and δ satisfying: $0 < \rho < 1$ where we take the sign + if I_j is on the right of t_0 and the sign - if I_j

is on the left of t_0 .

The choice of ρ :

In order to precise the choice of ρ , we observe that the problems occur for the boxes which are in the vicinity of one well and we consider a simplified function by taking the quadratic approximation of v at t_0 . This permits us to approximate the function $(s, t) \rightarrow W(t, s) = (1/16)(t - s)^2 + v(t_0 + (t + s)/2)$ by

$$(s, t) \rightarrow W_0(t, s) = C(t - s)^2 + D(t + s)^2$$

with $C > 0$ and $D > 0$. We consider the evolution of this potential restricted to an interval $I(x, \delta)$ centered at the point $(t = x, s = 0)$ and of size $2\delta_0$.

We observe that after dilation we can reduce our study to the case $\delta_0 = 1$, and we consider consequently the function:

$$]0, +\infty[\ni x \rightarrow W_0(x, s) \text{ for } s \in [-1, 1].$$

We have

$$(\partial_t W_0)(x, s) = 2C(x - s) + 2D(x + s) = (2C + 2D)x + (2D - 2C)s.$$

We first observe that if $C = D$, the function is a monotone increasing function of $x \geq 0$ and we could have taken $\rho = 0$ in this case but this is not the case here (in other papers we do a change of coordinates in order to be in this situation but it is not convenient here); we recall that $C = 1/16$, $D = v''(t_0)/8$ in our case.

Let us assume in order to fix the ideas that $C > D$. Then the derivative as a function of s is minimal at $s = 1$ and $(\partial_t W_0)(x, s)$ is positive for $x \geq (C - D)/(C + D)$.

Our ρ is chosen such that $|(C - D)|/(C + D) < \rho < 1$.

Using (3.4) and our choice of ρ , we see that, choosing ϵ_0 so small such that $\epsilon_0 \ll \delta_0$ with $\delta_0 \leq 1$, we get for $x \in \Omega$:

$$(3.7) \quad V(x) - V(\kappa(x)) \geq C\delta_0^2\beta(\Omega)$$

where $\beta(\Omega)$ is the number of intervals I_j which do not contain t_0 (or $-t_0$). Notice that $\beta(\Omega)$ is unchanged by the first " κ ".

Actually in this discussion, we have different cases to consider depending on the vanishing $\tilde{x}_j^{\Omega+}\tilde{x}_{j+1}^{\Omega+} = 0$ or not. We have only discussed above the most difficult case when this product is 0 with $|\tilde{x}_j^{\Omega+}| + |\tilde{x}_{j+1}^{\Omega+}| \neq 0$.

We now compose our two maps, we notice that $\alpha_0(\Omega) \leq \beta(\Omega)$ and we then get a new map $x \rightarrow \kappa(x)$, being the composition of reflexions in 0 in some of the coordinates and of a translation, such that

$$\kappa : \Omega \rightarrow \tilde{\Omega}_+ \subset 2\Omega_+,$$

(where Ω is the original box)

$$\kappa(x^\Omega) = \tilde{x}^{\Omega+}$$

and

$$(3.8) \quad V(x) - V(\kappa(x)) \geq (1/C)(\alpha_+(\Omega) + \beta(\Omega)), x \in \Omega.$$

C is here a strictly positive constant, independent of Ω , once we have fixed ϵ_0 and δ_0 conveniently as explained before.

Let P_Ω denote the Dirichlet realization of $-h^2\Delta + V$ in Ω and let μ_+ denote the lowest eigenvalue of $P_{\tilde{\Omega}_+}$. Let μ_0 denote the lowest eigenvalue of $-h^2\Delta + V$ on \mathbb{R}^n . By the minimax principle, we have

$$(3.9) \quad \mu_0 \leq \mu_+$$

and recall from Section 1 that, under the assumption $m = \mathcal{O}(h^{-N_0})$, we have a good knowledge of the asymptotics for μ_+ deduced from WKB constructions.

Formula (3.8) shows that:

$$(3.10) \quad P_\Omega - \mu_+ \geq (1/C)(\alpha_+(\Omega) + \beta(\Omega)).$$

Using κ we will also get, **under the assumption** $m = \mathcal{O}(h^{-N_0})$,

$$(3.11) \quad P_\Omega - \sum_1^m (x_j - x_j^\Omega)^{2M} - \mu_+^M(h) \geq (1/C)(\alpha_+(\Omega) + \beta(\Omega)), \text{ if } \Omega \neq \Omega_\pm,$$

with a new constant $C > 0$, where

$$(3.12) \quad \mu_+^M(h) - \mu_+(h) = \mathcal{O}(h^{N(M)}),$$

with $N(M) \rightarrow \infty$ as $M \rightarrow \infty$ and $h > 0$ sufficiently small. We observe indeed that:

$$\sum_1^m (\tilde{x}_j - \tilde{x}_j^\Omega)^{2M} = \sum' (\tilde{x}_j)^{2M} + \sum'' (\tilde{x}_j - \tilde{x}_j^\Omega)^{2M},$$

where \sum' corresponds to the sum over the j such that $t_0 \in I_j$. We now obtain the majoration:

$$\sum_1^m (\tilde{x}_j - \tilde{x}_j^\Omega)^{2M} \leq \sum_1^m \tilde{x}_j^{2M} + \beta(\Omega) \epsilon_0^{2M}; \quad \forall \tilde{x} \in \kappa(\Omega)$$

and then get (3.11) as in [Sj]1 (Section 6), [Sj]2 or [He-Sj].