

Parasitism of *Agriotes obscurus* linneaus (Coleoptera : Elatreidae) by *Hexamermis* sp. (Nematoda : Mermithidae)

Autor(en): **Doane, J.F. / Klingler, Jakob / Welch, H.E.**

Objekttyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft =
Bulletin de la Société Entomologique Suisse = Journal of the
Swiss Entomological Society**

Band (Jahr): **45 (1972)**

Heft 4

PDF erstellt am: **24.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-401692>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

PARASITISM OF AGRIOTES OBSCURUS LINNEAUS (COLEOPTERA: ELATERIDAE) BY HEXAMERMIS SP. (NEMATODA: MERMITHIDAE)

J. F. DOANE¹ and JAKOB KLINGLER

Swiss Federal Research Station, Wädenswil, Switzerland

H. E. WELCH

Department of Zoology, University of Manitoba, Winnipeg, Manitoba

Records of nematode parasitism of elaterids are few. VAN ZWALUWENBURG (1928) reported that an elaterid pupa, probably *Monocrepidius* sp., was parasitized by an unidentified mermithid. Several genera of nematodes also have been found associated with dead elaterid larvae (KÖRNER 1954; VEREMCHUK 1962, 1969), but such associations are probably not the result of primary parasitism. Primary parasitism of *Agriotes lineatus* by the nematode *Neoaplectana carpocapsae* was reported by POINAR & VEREMCHUK (1970). The different types of insect-nematode relationships were reviewed by WELCH 1963, 1965.

Nematodes, later identified as belonging to the mermithid genus *Hexameris*, were found parasitizing larvae of a wireworm, *Agriotes obscurus*² (Linneaus). The wireworms, collected near Wädenswil³, Switzerland, during the autumn of 1970, were being reared for other studies when the nematodes emerged.

A. obscurus larvae collected 15 September were taken to the laboratory, held for 3 days at 6°C, and then 10 large larvae (about 20 mm) and 20 medium-sized larvae (about 13 mm) were placed in small plastic boxes filled with sieved soil and supplied with live wheat kernels for food. The larvae were examined two or three times a month and at these times provided with fresh soil and food. During the entire period of examination the larvae were held at approximately 22°C.




Fig. 1. *Agriotes obscurus* (Linneaus) larva (actual size 13 mm) and a mermithid parasite (*Hexameris* sp.) that emerged from this larva.

Between 13 October and 19 December, 10 nematodes ranging in size from about 15 to 35 mm emerged from six of the medium-sized wireworms, an average of 1.7 per infected host. None emerged from wireworms longer than 14 mm. The numbers of nematodes per wireworm in the sample of 30 larvae were as follows: 0 nematodes in 22 larvae; 1 in 3; 2 in 2; 3 in 1. The per cent infection of the population sample of 30 wireworms was therefore 20 per cent, or if only the medium-sized larvae are considered, 30 per cent. Parasite emergence, which killed the host, appeared to take place through the mouth, anus or possibly the spiracles because no break in the integument could usually be found. However, in one case emergence was apparently through the intersegmental membrane between the second and third abdominal segments on the dorsum. After emergence the nematodes were heat relaxed in water on a cavity slide over a low flame, fixed in 4 per cent formalin and held for examination.

The feeding activity of the six parasitized wireworms did not seem greatly affected as compared to that of healthy larvae except perhaps within the 2-week period before parasite emergence. Even so, three of the parasitized larvae fed within 2 weeks of nematode emergence. During the last few days before parasite emergence parasitized wireworms became sluggish or quiescent, appeared thicker than normal, and resembled larvae about to moult. The apparent small effect of the nematode parasites on the feeding activity of the wireworms is consistent with the observation that mermithids do little damage to their hosts except shortly before parasite emergence (WELCH 1965).

Since parasites emerged from medium-sized wireworms 12–14 mm long apparently either a particular larval stage is attacked or the nematode infection tends to affect the growth rates of larvae so that they are all about equal size when the nematodes emerge.

Acknowledgments

The authors wish to thank Mr. F. L. Dolezsar, Canada Department of Agriculture Research Station, Saskatoon, for the photography.

¹Present address: Canada Agriculture Research Station, Saskatoon, Sask., Canada.

² Larvae of *A. obscurus* cannot be definitely separated from those of *Agriotes lineatus* (Linneaus). However, because adults collected from this location were all *A. obscurus* the larval population probably contained no *A. lineatus* larvae.

³ Coordinates of this location are: 693.375/299.300. (Kilometer-grid of Swiss national map).

REFERENCES

- KÖRNER, H. 1954. Die Nematodenfauna des vergehenden Holzes und ihre Beziehungen zu den Insekten. *Zool. Jahrb. Abt. 82*: 245–353.
- POINAR, JR., G. O. & VEREMCHUK, G. V., 1970. (A new strain of nematodes pathogenic to insects and the geographical distribution of *Neoaplectana carpocapsae* Weiser [Rhabditida, Steinernematidae]). *Zool. ZH.* 49, 966–969, (In Russian).
- WELCH, H. E. 1961. Nematodes as agents of insect control. *Proc. ent. Soc. Ontario* 92: 11–19.
- 1963. Nematode infections, pp. 363–392. In E. A. Steinhaus (ed.), *Insect pathology*. Academic Press, New York.
- 1965. Entomophilic nematodes. *A. Rev. Ent.* 10: 275–302.
- VAN ZWALUWENBURG, R. H. 1928. The interrelationships of insects and wireworms. *Bull. Hawaiian Sugar Planters Assoc. Expt. Sta., ent. Ser.*, 20, 68 pp.
- VEREMCHUK, G. V. 1962. (Entomohelminthological family Steinernematidae Chitwood and Chitwood 1937) *Tezisy Dokl. nauch. Konf. vses. Obshch. Gel'minto. Akad. Nauk S. S. R.*, 10–14 Dekabrya 1962, Chast 1: 24–25, (In Russian).
- VEREMCHUK, G. V. 1969. (New species of *Neoaplectana* [Rhabditida: Steinernematidae] nematodes pathogenic to Elateridae) *Mater. nauch. Konf. vses. Obshch. Gel'minto.*, 1969, Part 1, pp 44–50 (In Russian). *Helm. Abst.* 1971, 40 (3) No. 1779.