

# **Two new species of Kempnyia from southern Brazil (Plecoptera : Perlidae)**

Autor(en): **Froehlich, C.G.**

Objekttyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society**

Band (Jahr): **69 (1996)**

Heft 1

PDF erstellt am: **23.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-402621>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## Two new species of *Kempnyia* from southern Brazil (Plecoptera: Perlidae)

C.G. FROEHLICH

Departamento de Biologia, FFCLRP, Universidade de São Paulo, BR-14040-901 Ribeirão Preto, SP, Brazil.

Two new species of *Kempnyia*, *K. auberti* and *K. petersorum* are described. Both were collected in southern São Paulo State and in Paraná State and are related to *K. neotropica* J. & B., a widespread species in southeastern and southern Brazil.

Keywords: Plecoptera, *Kempnyia*, new species, Brazil

### INTRODUCTION

Two new species of *Kempnyia* Klapalek, both collected in southern São Paulo State and in Paraná State, are described. The types are deposited in the Museum of Zoology, University of São Paulo (MZSP); the paratypes in the National Museum of Natural History, Smithsonian Institution, Washington, DC, USA (NMNH).

### DESCRIPTIONS


#### *Kempnyia auberti* sp.n.

(Figs. 1-8)

*Material:* Holotype male: SP, Ribeirão Grande, Fazenda Intervales, 10.XI.1993, C.G. FROEHLICH and H. PAPROCKI (MZSP). Paratype male: PR, Rio Marumbi, 15-16.II.1969, W.L. and J.G. PETERS (NMNH).

*Measurements:* Holotype: forewing length, 22.0 mm; paratype: forewing length, 18.4 mm.


*Male:* Head (Fig. 1) brown on both sides of the ochraceous M-line, at lappets, and backwards to postfrontal line. Sides of head ochraceous brown to ochraceous. Scape of antennae ochraceous on dorsal and ventral sides, brownish on anterior and posterior faces. Pedicel brown, darker basally. Flagellum medium brown. Palpi light brown. Pronotum (Fig. 1) brown, rugosities a shade lighter, broad median ochraceous stripe. Meso- and metanotum brown to ochraceous. Dorsal abdomen pale yellowish. Femora yellowish, changing to brown apically. Tibiae brownish, darker at both ends. Tarsi ochraceous, brown apically. Wing membrane brown, lighter in centre of many cells. C, Sc, R, M in part, A<sub>2</sub> and A<sub>3</sub> pale, rest of veins brownish. Cerci medium brown, brown hairs on paler cuticle. Ventral side pale, basisterna in part brown.



Figs. 1-8: *Kempnyia auberti* sp.n., male holotype. Fig. 1, head and pronotum; Fig. 2, sternum 9; Fig. 3, tergum 10; Fig. 4, detail of tergum 10 (peg-like setae); Fig. 5, detail of paraproct; Figs. 6-8, penial armature in lateral, ventral and dorsal views.

Sternum 9 (Fig. 2) light in colour, brownish at sides and darker at bases of the subgenital plate rim. A median pale brown stripe backwards from hammer. Subgenital plate relatively short and broad, hammer with an elongate triangular shape. Field of special (peg-like) bristles on tergum 10 (Fig. 3) moderate in size, most bristles carrot-shaped (Fig. 4). Paraprocts digitiform, with minute sensillae apically (Fig. 5). Penial armature (Figs. 6-8) elongate, hooks moderately twisted, a pair of spiny pads on ventral membranous penis wall, basally from armature.

*Female and nymph:* Unknown.



Figs. 9-15: *Kempnyia petersorum* sp.n., male holotype. Fig. 9, head and pronotum; Fig. 10, sternum 9; Fig. 11, tergum 10; Fig. 12, detail of tergum 10 (peg-like setae); Figs. 13-15, penial armature in lateral, dorsal and ventral views.

**Discussion:** The penis of *K. auberti* is very similar to that of *K. neotropica* (ZWICK, 1972) both in the shape of the armature and the presence of the ventral spiny pads. *K. auberti*, however, is larger and lighter in colour. *K. neotropica* is a dark species, as several of the names given to it indicate: *obscura* PICTET, *aterrima* KЛАРАЛЕК, and *fusca* NAVAS. In the type locality both species are sympatric.

**Etymology:** The name honours the late Prof. J. AUBERT.

#### *Kempnyia petersorum* sp.n.

(Figs. 9-15)

**Material:** Holotype, male: SP, Ribeirão Grande, Fazenda Intervales, 7.II.1989, E. TRAJANO (MZSP). Paratypes, 4 males: PR, Rio Marumbi, 15-16.II.1969, W.L. and J.G. PETERS (NMNH).

*Measurements:* Holotype, forewing length, 12.4 mm; paratypes: forewing length, 11.1-11.7 mm.

*Male:* Head (Fig. 9) with broad longitudinal brown band; sides of head brownish-ochraceous but parietalia brown posteriorly. Scape brownish-ochraceous, rest of antennae medium to light brown. Palpi light brown. In both antennae and palpi, brown hairs on pale cuticle. Pronotum (Fig. 9) brown, rugosities a shade lighter. Meso- and metanotum ochraceous, some lines brownish. Abdominal terga light brown, darker towards terminalia. Wing membrane brownish, veins pale but with row of brown microtrichia along them. Legs brown, bases of femora yellowish, tarsi light brown. Ventral side yellowish, basisterna in part brown. Abdominal sterna light brown, darker towards terminalia. Cerci brownish; from about the 10th cercomere on, bicoloured.

Sternum 9 (Fig. 10) relatively long, light-coloured. Subgenital plate rounded, hammer long-oval; anteriorly to hammer, a pale brown longitudinal stripe. Field of special (peg-like) bristles on tergum 10 (Fig. 11) moderate in size, most bristles elliptical (Fig. 12). Paraprocts digitiform, with minute sensillae apically. Penial armature (Figs. 13-15) elongate, hooks moderately twisted, a pair of spiny pads on ventral membranous penis wall, basally from armature.

*Female and nymph:* Unknown.

*Discussion:* The penis of *K. petersorum* is also similar to that of *K. neotropicica* both in the shape of the armature and in the presence of spiny pads. It differs from *K. neotropicica* and *K. auberti* in the colour pattern, the brown band along the head being very characteristic. Also the contrast of the brown pronotum with the light-coloured meso- and metanotum is striking. It is sympatric with both *K. neotropicica* and *K. auberti*. If the spiny penial pads prove to be a synapomorphy, they form a natural group.

*Etymology:* Dedicated to William L. and Janice G. PETERS, the collectors of the paratypes.

#### ACKNOWLEDGEMENTS

To the CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for a Senior Fellowship (Grant No. 201124/94.7). To the authorities of the Fundação Florestal do Estado de São Paulo for permission and facilities to work in Fazenda Intervales. To Dr. Oliver S. FLINT, Jr., for making possible my visit to the Natural History Museum of the Smithsonian Institution, Washington DC, USA. To Dr. M. SARTORI for the invitation to participate in the *Festschrift* honouring Prof. J. AUBERT.

#### REFERENCE

- ZWICK, P. 1972. Die Plecopteren Pictets und Burmeisters, mit Angaben über weitere Arten (Insecta). *Rev. Suisse Zool.* 78 (4): 1123-1194.