

**Zeitschrift:** Acta Tropica  
**Herausgeber:** Schweizerisches Tropeninstitut (Basel)  
**Band:** 27 (1970)  
**Heft:** 2

**Artikel:** Miscellanea : Terrestrial snail a new host for trombiculid larvae  
**Autor:** Vercammen-Grandjean, P.H. / Benoit, P.L.G. / Mol, J.J. van  
**DOI:** <https://doi.org/10.5169/seals-311639>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 01.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Terrestrial Snail a New Host for Trombiculid Larvae

P. H. VERCAMMEN-GRANDJEAN, P. L. G. BENOIT \* and J. J. VAN MOL \*\*

The G. W. Hooper Foundation, University of California,  
San Francisco, California 94122, U.S.A.

To our knowledge, larval chiggers (Acarina: Trombiculidae) have never been found parasitizing molluscs. In fact, very few groups of acarines do parasitize those invertebrates, and, even so, most of them only live on or in molluscs as commensals and not at the expense of the host. In this way members of the families Pontarachnidae, Unionicolidae, and Halacaridae inhabit the mantle and the gills of certain lamellibranches (marine and fresh-water) such as oysters, clams, and mussels, and the gills of certain aquatic gastropods like *Ampullaria* in South




Fig. 1. A *Granularion* sp. (Urocyclidae: Mollusca) with – as distinct intradermal white patches – *Schoengastia* (*Endotrombicula*) sp. (Trombiculidae: Acarina); collected in Sierra Leone (Mt. Loma), Africa, 1964.

America. A group of Ereyneidae parasitize the lungs of terrestrial gastropods, like *Helix pomatia* and various slugs, in Europe, America and Australia (genus *Riccardoella*). Finally, only one Halacaridae (*Halixodes chitonis*) really lives at the expense of its host, a polyplacophore, or chiton, from New Zealand. Restriction of parasitism to vertebrates was regarded as characteristic of the Trombiculidae until a few species were discovered parasitizing arthropods.

Several specimens of a terrestrial snail, *Granularion* n. sp. (Urocyclidae) collected in 1964 from Mont Loma, Sierra Leone, by an expedition of the "Institut Fondamental d'Afrique Noire (I. F. A. N.)" have now been found freely parasitized by an intradermal mite, *Schoengastia* (*Endotrombicula*) sp. (see photograph). A description is in preparation.

\* Département des Invertébrés, Musée d'Afrique Centrale, Tervuren, Belgium.

\*\* Chaire de Zoologie Systématique, Université Libre de Bruxelles, Belgium.