

Zeitschrift: Archives des sciences [1948-1980]
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 33 (1980)
Heft: 1-3

Artikel: High grade rodingites from the Central Alps : metamorphism and geochemistry
Autor: Trommsdorff, Volkmar / Evans, Bernard W.
DOI: <https://doi.org/10.5169/seals-739487>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 15.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

HIGH GRADE RODINGITES FROM THE CENTRAL ALPS: METAMORPHISM AND GEOCHEMISTRY

BY

Volkmar TROMMSDORFF¹ and Bernard W. EVANS²

High grade rodingites occur in at least two different suites in the Central Alps (Fig. 1). *The first suite* occurs in the Cima Lunga-Adula nappe (especially well developed in the area of Cima di Gagnone) and comprises mafic rocks consisting predominantly of garnet and clinopyroxene and varying from eclogite (CaO 12, Na₂O 2.8 weight percent) to metarodingite (CaO up to 25, Na₂O, O weight percent, Trommsdorff *et al.* 1975, Evans *et al.* 1979). Garnets vary correspondingly from pyralspite 84 ugrandite 16 to ugrandite 80 pyralspite 20 and clinopyroxenes from omphacite (jadeite 45) to diopside and fassite (Cats 23). Trace element abundances appear to have been little affected by the process of rodingitization and suggest sea floor basalt tholeiitic affinities of the suite.

Associated ultramafic bodies vary from chlorite-enstatite peridotite (with in part 0% CaO) enclosing rodingites, to garnet peridotite (CaO up to 4%) enclosing eclogite and are believed to at least in part represent progressively metamorphosed products of serpentinite.

For garnet peridotite and eclogites several independent geothermometers and —barometers suggest conditions of ~25kbar and 800° C (see also Heinrich (1978)). The same conditions hold for the rodingites based upon a new calibration for low Na-pyroxene-calcic garnet pairs by Ellis and Green (1979). Within the eclogites, metarodingites and garnet peridotite exchange equilibria of the high pressure assemblages were little affected by the later, main Alpine amphibolite facies metamorphism of the Central Alps.

However, in contact reaction zones between ultramafics and enclosed mafic lenses these early assemblages were destroyed by the main Alpine event resulting in partial amphibolitisation of eclogites and metarodingites.

¹ Inst. für Kristallographie und Petrographie, ETH-Zentrum, CH-8092 Zürich.

² Dept. of Geological Sciences, University of Washington, Seattle, Wash. 98195, U.S.A.




FIG. 1. — Outline tectonic map of the Central Alps near Bellinzona.

Table 1 gives a summary of events recognized for the mafic-ultramafic suite of Cima di Gagnone.

TABLE 1. — *Sequence of events in Cima-Lunga mafic-ultramafic rocks.*

Time	Stage	Event	Rock types	
			Lherzolithic ultramafics	Tholeiitic basalts
			↓	↓
	Oceanic	serpentinization/ rodingitization	serpentinization	in part rodingitization
			↓	↓
	Deep burial	High-pressure metamorphism (25kb/800 °C)	chlorite peridotite	garnet peridotite
			↓	↓
	Alpine orogeny	Amphibolite facies meta- morphism	in part chlorite peridotite ± spinel	in part amphibolite (blackwall, amphiboliti- zation)

In the Bellinzona zone (Fig. 1), at Alpe Albion east of Bellinzona a *second ultramafic-rodingite suite* has been recognized. This suite shows no signs of an early high-pressure metamorphic event. The Alpine metamorphic event in this area was of sillimanite grade and the ultramafic rocks contain a spinel-tremolite-chlorite-enstatite-olivine assemblage. Metarodingite dykes are mostly flattened parallel to a strong Alpine foliation and are scarcely crosscutting. They resulted from original coarse grained dykes that locally show relic gabbro textures. Their CaO content varies from 20 to over 33 weight percent. The original inhomogeneity of these rocks lead to domains in the rodingites that are dominated by various minerals i.e. pyroxene domains with a highly fassaitic clino-pyroxene, epidote domains, garnet domains, vesuvianite domains and iron titanium oxide domains. In the complex mineral assemblages of the latter sphene, perovskite and melanite are found, in the vesuvianite domains xanthophyllite is not uncommon. Blackwall reaction zones between the ultramafics and metarodingite contain much chlorite, orthopyroxene, olivine, spinel, and calcic amphibole. Hoegbomite occurs in local association with spinel. Against the metarodingite a 5 mm wall of black hornblende is found.

This postkinematic hornblende frequently encloses clinopyroxene relics indicating at least growth of some blackwall material on expense of the metarodingite during Alpine metamorphism.

On the other hand shearing of blackwall within folded metarodingite boudins indicates formation of blackwall material in a prekinematic stage.

The petrogenetic history of the Bellinzona zone-ultramafics can be derived from these observations and involves at least three stages:

- 1) Metasomatic rodingitization of coarse grained gabbro dykes accompanying serpentinization of enclosing peridotite.
- 2) Severe deformation mylonitization and folding involving probably more than one event the last being Alpine in age.
- 3) Alpine metamorphism, sillimanite grade, in part postkinematic.

The subdivision of the Central Alpine ultramafics into two zones, one that has experienced high pressure metamorphism and one that doesn't is in agreement with observations by Fumasoli (1974). In both zones, however, an early stage of serpentinization accompanied by rodingitization is recognized.

ACKNOWLEDGEMENTS

The authors acknowledge the support of Schweizerischer Nationalfonds No. 2.115-0.78 and U.S. National Science Foundation (Grant EAR-78-03350).

REFERENCES

- ELLIS, D. J. and D. H. GREEN (1979). An experimental study of the effect of Ca upon garnet-clinopyroxene Fe-Mg exchange equilibria. *Contrib. Mineral. Petrol.*, 71, 13-22.
- EVANS, B. W., V. TROMMSDORFF and W. RICHTER (1979). Petrology of an eclogite-metarodingite suite at Cima di Gagnone, Ticino, Switzerland. *Amer. Mineralogist*, 64, 15-31.
- FUMASOLI, M. W. (1974). Geologie des Gebietes nördlich und südlich der Iorio-Tonale-Linie im Westen von Gravedona (Como Italia). Thesis, University Zürich.
- HEINRICH, C. (1978). Metamorphose und Strukturen der Cima-Lunga Serie. I. Cima di Gagnone-Val Motto. Diplomarbeit, ETH Zürich.
- TROMMSDORFF, V., B. W. EVANS und W. RICHTER (1975). Eklogit/Rodingit Übergänge in Ultramafiten der Cima-Lunga-Serie. *Schweiz. mineral. petrogr. Mitt.*, 55/3.