

**Zeitschrift:** Archives des sciences et compte rendu des séances de la Société  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 44 (1991)  
**Heft:** 3: Archives des Sciences

**Artikel:** New and interesting mites from the Geneva Museum LXXII : some  
anoetid and tarsonemid mites from the Cape Verde Islands (Acari)  
**Autor:** Mahunka, Sandor / Mahunka-Papp, Luise  
**DOI:** <https://doi.org/10.5169/seals-740203>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 02.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Archs Sci. Genève	Vol. 44	Fasc. 3	pp. 283-287	Décembre 1991
-------------------	---------	---------	-------------	---------------

NEW AND INTERESTING MITES FROM THE  
GENEVA MUSEUM LXXII.  
SOME ANOETID AND TARSONEMID MITES  
FROM THE CAPE VERDE ISLANDS (ACARI)

BY

**Sandor MAHUNKA\*** and **Luise MAHUNKA-PAPP\***

ABSTRACT

**New and interesting mites from the Geneva Museum LXXII. Some Anoeetid and Tarsonemid mites from the Cape Verde Islands (Acari).** – Three Anoeetid and six Tarsonemid species are recorded originating from soil samples from the Cape Verde Islands. One species is new to science and necessitates the erection of a new genus: *Stercoranoetus harteni* gen. n., sp. n. A second species of this genus is *S. woolleyi* (Mahunka & Eraky, 1987) comb. n., transferred from *Histiostoma* Kramer, 1876.

RÉSUMÉ

Dans des échantillons de sol provenant des îles du Cap Vert, les auteurs ont trouvé trois espèces d'Anoeetidés et six de Tarsonemidés (Acariens). Une espèce est nouvelle et nécessite la création d'un genre nouveau: *Stercoranoetus harteni* gen. n., sp. n.; une seconde espèce *Stercoranoetus woolleyi* (Mahunka & Eraky, 1987) comb. n., a été transférée du genre *Histiostoma* Kramer, 1876.

INTRODUCTION

The mite fauna of the Cape Verde Islands has been poorly explored and no data have been published – as far as we know – either for *Anoetidae* or for *Tarsonemina*. Studying soil samples collected by Mr. A. van Harten we have found some mites belonging to these taxa.

Owing to their geographical situation the fauna of the Cape Verde Islands is very interesting from a zoogeographical point of view (MAHUNKA 1987, 1991). The following list contains species found until now only in the Palaearctic or Holarctic Region (*Brennandania csibiae*, *Heterodispus pubescens*), and species which live in both temperate and tropical zone (*Histiostoma phyllophorus*, *Pseudopygmephorus tarsalis*). It is also very interesting, that the species most related to the new genus live in the tropics and in Europe.

We are very grateful to Dr. B. Hauser and Dr. C. Lienhard for allowing us to study this very interesting material deposited in the Muséum d'Histoire naturelle, Geneva.

LIST OF LOCALITIES

- 655: Santiago: São Jorge dos Orgãos, in litter, IX. 1983.  
1532: Santiago: São Jorge dos Orgãos, in litter, I. 1986.

---

\*Zoological Department, Hungarian Natural History Museum, Baross utca 13, H-1088 Budapest, Hungary.

- 1787: Santiago: São Jorge dos Orgãos, in litter, I. 1986.  
 1804: Santiago: Santa Cruz, in litter, 28.III.1988.  
 1809: Santiago: Ribeira de Charco, in litter, 6.V.1988.  
 1837: Santiago: Achada Fazenda, in litter, 16.V.1988.  
 1882: Santiago: Serra de Malagueta, in litter, 29.V.1988.

### LIST OF SPECIES

ACARIDA Latreille, 1802

**Anoetidae** Oudemans, 1904

*Stercoranoetus harteni* gen. n., sp.n.

Localities: 1804, 1837, 1882.

*Histiostoma phyllophorus* (Oudemans, 1905)

A wide-spread species, extracted mostly from manure.

Locality: 1804: 2 specimens.

*Glyphanoetus phyllotrichus* (Berlese, 1881)

This species was described from dung, until now many authors published it also from manure and from animal excrement.

Locality: 655: 1 specimen.

TARSONEMINA Canestrini Fanzago, 1877

**Pygmephoridae** Cross, 1965

*Pseudopygmephorus quadratus* Ewing, 1909

A cosmopolitan species present wherever organic material – above all manure or dung – is accumulated.

Locality: 1809: 1 specimen.

*Pseudopygmephorus tarsalis* (Hirst, 1921)

A cosmopolitan (?) species. It is frequent in animal manure.

Localities: 1532: 2 specimens; 1809: 1 specimen; 1882: 1 specimen.

**Microdispidae** Cross, 1965

*Brennandania csibiae* Mahunka, 1981

So far it has been known only from Hungary and Greece (MAHUNKA, 1981).

Locality: 655: 1 specimen.

*Brennandania silvestris* (Jacot, 1936)

A common Holarctic species.

Locality: 655: 2 specimens.

**Scutacaridae** Oudemans, 1916

*Heterodispus pubescens* Mahunka, 1969

It was described from Hungary and known from here only.

Localities: 655: 4 specimens; 1882: 10 specimens.

*Scutacarus longitarsus* (Berlese, 1905)

Common in the Palaearctic Region, several localities also known in the tropical regions.

Locality: 655: 2 specimens.

## DESCRIPTION

***Stercoranoetus* gen. n.**

**D i a g n o s i s :** Family *Anoetidae*. Humeral part of hysterosoma anteriorly protruding, covering the posterolateral part of the propodosoma. Dorsal surface well ornamented. Infracapitulum normal, palps very short. Sejugal apodemes straight, posterior sternal apodeme fused with them. Legs 1 and 2 stout, joints short but wide. Tibia and genu of leg 1 with spiniform appendages, tibia and genu of leg 2 with longitudinal lateral edges. All legs with claws.

**T y p e s p e c i e s :** *Stercoranoetus harteni* sp. n.

**R e m a r k s :** The new genus is well characterized by the humeral apophysis and by the shape of legs 1 and 2. On this basis it belongs to the alliance of *Pteranoetus* Mahunka, 1978, however, the latter has a modified gnathosoma and completely different apodemes. A very similar configuration exists in an other recently described *Anoetidae* species (*Histiostoma woolleyi* Mahunka & Eraky, 1987), which has to be transferred to the new genus: *Stercoranoetus woolleyi* (Mahunka & Eraky, 1987) comb. n.

***Stercoranoetus harteni* sp. n.**

**M e a s u r e m e n t s .** – Length: 155-165  $\mu\text{m}$ , width: 118-125  $\mu\text{m}$ .

**D o r s a l s i d e** (Fig. 1): Propodosoma very large, wide, trilobate anteriorly. Beside the sharp rostral apex two rounded lateral ones present; the rostral apex not longer than the others. Surface ornamented by polygonate pattern of arcuate lines. Both pairs of prodorsal setae straight, strong and erect; the inner pair originating only slightly anteriorly to the outer pair. Dorsosejugal region striate. Hysterosoma also very wide, its humeral part protruding forwards and covering the lateral part of propodosoma. Lateral part of hysterosoma also ornamented by scratched lines, like propodosoma, this pattern well observable also near posterior margin. Eleven pairs of notogastral setae present, five pairs of them very long, slightly dilated. The other setae simple, short, but well observable.


**V e n t r a l s i d e** (Fig. 4): Infracapitulum long, slightly dilated medially, palps very short. Anterior sternal apodeme ending free, *ap.* 2 fused with the straight sejugal apodemes. These latter with a small lateral arch. Posterior sternal apodeme also fused with the sejugal apodemes. Epimeral suckers well developed,  $S_1$  originating above apodeme 2,  $S_2$  far from apodeme 4. Adhering plate normal, ending far from posterior margin of the body.

**L e g s :** Legs 1 and 2 thick (Figs 2-3) all joints short and wide. Tibia and genu of leg 1 with a sharp spur on their outer corner (Fig. 7), tibia and genu of leg 2 with a sharp,

arched lateral edge. Tarsi of legs 3 (Fig. 5) and 4 (Fig. 6) each with a short, phylliform terminal seta.

**Material examined:** Holotype: 1804; 7 paratypes: from the same sample; 2 paratypes: 1837; 3 paratypes: 1882. Holotype and 8 paratype: MHNG<sup>1</sup>; 4 paratypes (1331-PA-1988): HNHM<sup>2</sup>.

**Remarks:** *Stercoranoetus harteni* differs from *S. wooleyi* by the length of notogastral setae and by the form of the propodosoma.




FIGS 1-3.

*Stercoranoetus harteni* gen. n., sp. n. - 1: dorsal side, 2: leg 1, 3: leg 2.

<sup>1</sup> MHNG = deposited in the Muséum d'Histoire naturelle, Genève.

<sup>2</sup> HNHM = deposited in the Hungarian Natural History Museum, Budapest, with the identification number of the specimens in the Collection of Arachnoidea.



FIGS 4-7.

*Stercoranoetus harteni* gen. n., sp. n. — 4: ventral side, 5: leg 3, 6: leg 4, 7: tibia and genu of leg 1 from ventral view.

## REFERENCES

- MAHUNKA, S. 1972. Tetüatkák – Tarsonemina. *Fauna Hung.* 110: 215 pp.
- 1981. The Pygmephoroid Fauna of the Hortobágy National Park (Acari: Tarsonemida). In: MAHUNKA, S. (ed.): *The Fauna of the Hortobágy National Park, Akadémiai Kiadó, Budapest*, vol. 1: 343-370.
- 1987. Neue und interessante Milben aus dem Genfer Museum LVIII. Some primitive Oribatids from the Cape Verde Island (Acari: Oribatida). *Revue suisse Zool.* 94: 109-116.
- 1991. New and interesting mites from the Geneva Museum. LXX. Oribatids from the Cape Verde Islands II (Acari: Oribatida). *Revue suisse Zool.* 98: 567-580.
- MAHUNKA, S. & S. A. ERAKY 1987. Mites extracted from manure and silo (Acari: Acaridae, Anoetidae and Tarsonemina), I. *Fol. ent. hung.* 48: 129-140.

