

**Zeitschrift:** Botanica Helvetica  
**Herausgeber:** Schweizerische Botanische Gesellschaft  
**Band:** 100 (1990)  
**Heft:** 1

**Artikel:** Durio bukitrayaensis Kosterm. (Bombacaceae), a new species from Borneo  
**Autor:** Kostermans, A.J.G.H.  
**DOI:** <https://doi.org/10.5169/seals-69709>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## *Durio bukitrayaensis* Kosterm. (Bombacaceae), a new species from Borneo

A. J. G. H. Kostermans

Herbarium Bogoriense, Bogor, Indonesia

Manuscript accepted January 8, 1990

### Abstract

Kostermans A. J. G. H. 1990. *Durio bukitrayaensis* Kosterm. (Bombacaceae), a new species from Borneo. Bot. Helv. 100: 29–31.

*Durio bukitrayaensis* is described as a new species from Borneo. It is related to *D. dulcis*, *D. grandiflorus*, and *D. testudinarum*.

### *Durio bukitrayaensis* Kosterm., spec. nov.

Arbor mediocris, foliis alternantibus chartaceis oblongis (pro genere longis), longe cuspidatis basi rotundatis, supra subnitidis, sub lente minutissime reticulatis glabris, nervo mediano conspicue impresso, nervis filiformibus impressis, subtus nitidis adpresse lepidotis, squamulis non fimbriatis, nervo mediano prominente, nervis subpatentibus, epicalycis partibus ovato-acutis lepidotis, calycis partibus lepidotis squamulis majusculis, fructibus ellipsoideis, parvis, flavoviridibus spinis subcylindricis parvis acutis squamulis minutis fimbriatis obtectis.

*Typus*: Mogeia 4128 (BO).

Tree, 15 m tall, dbh. 15 cm. Bark dark brown, scurfy. Branchlets tightly crusty-scaly. Leaves scattered, chartaceous, oblong, 6 × 15 – 7 × 22 cm – 10 × 32 cm, acumen very slender, sharp, up to 4 cm long, base rounded; above green, under the lens minutely reticulate, nerves scabrous, no scales; below silvery brown or brownish-glaucous (fresh), glossy, covered with very tightly packed, flat, non-fimbriate scales of 0.5–0.75 mm diam., so tightly packed that their shape is hardly discernible, midrib strongly prominent, nerves rather patent, slender, prominent, c. 17 pairs with in between shorter ones. Petiole rather slender, densely lepidote, 10–15 mm long.

Flowers on the very base of the trunk, cauliflorous or on the thicker branches. Buds shortly ovoid, or conical. Epicalyx of broadly ovoid, acute, 2.5 cm long lobes, densely and tightly covered by non-fimbriate scales. Calyx lepidote, scales 1–1.2 mm diam.

Fruit ellipsoid, yellowish green, c. 7 cm long, 6 cm diam., rounded at both ends; spines rather slender, 1.5 cm long, densely, minutely fimbriate-scaly, at base c. 1 mm diam., stiff, sharp.

Distribution/Ecology. Lowland, 350 m, in wet, evergreen, tropical rainforest in the Bukit Raya Mts., Central Kalimantan.




Fig. 1. *Durio bukitrayaensis* Kosterm.

*Note:* The non-fimbriate, very tightly packed scales of the under surface of the leaves resemble those of *D. dulcis* and *D. grandiflorus* but the scales are much larger. Moreover, the leaves of *D. dulcis* and *D. grandiflorus* are smaller and differently shaped, the patent nerves are also different. Like in *D. testudinarum*, the flowers may occur on the very base of the bole, but the leaves are different, characteristic is the long draw-out slender acumen, and the scale are much smaller than those of *D. testudinarum*. The fruits are small, the spines look like those of *D. grandiflorus*.

Central Kalimantan, Bukit Raya in Upper Katingan (Mendawi) R. area, Upper Samba R., 60 km NNW of Tumbang Samba, 112°5 E, 0°60 S, alt. 350 m, Dec., fr. Mogeia 4128 (BO); Upper Katingan R., NNW of Tumbau Samba, 112°30 E, 1°15 S, Dec., buds, Mogeia 4417 (BO).