

Zeitschrift: Botanica Helvetica
Herausgeber: Schweizerische Botanische Gesellschaft
Band: 109 (1999)
Heft: 1

Artikel: Plant communities with *Pinus sylvestris* L. and *P. nigra* Arnold subsp. *salzmanii* (Dunal) Franco of the Spanish Sistema Central : a phytosociological approximation
Autor: Galán de Mera, Antonio / Hagen, M. Angeles / Vicente Orellana, José Alfredo
DOI: <https://doi.org/10.5169/seals-73285>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 01.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Plant communities with *Pinus sylvestris* L.
and *P. nigra* Arnold subsp. *salzmannii* (Dunal)
Franco of the Spanish Sistema Central:
a phytosociological approximation

Antonio Galán de Mera, M^a Angeles Hagen and José Alfredo Vicente Orellana

Department of Biology, Laboratory of Botany, San Pablo-CEU University, P.O. Box 67,
28660 Boadilla del Monte (Madrid), Spain

Manuscript accepted March 16, 1998

Abstract

Galán de Mera A., Hagen M. A. and Vicente Orellana J. A. 1999. Plant communities with *Pinus sylvestris* L. and *P. nigra* Arnold subsp. *salzmannii* (Dunal) Franco of the Spanish Sistema Central: a phytosociological approximation. Bot. Helv. 109: 21–54.

A phytosociological study of forests with *Pinus sylvestris* and *P. nigra* subsp. *salzmannii* of the Spanish Sistema Central has been made, establishing a comparison with 683 coniferous communities. As a result of the application of the concepts of Kopecký & Hejný (1974), Foucault (1981), Dierschke (1993) and Kopecký et al. (1995), on the relevés made following Braun-Blanquet (1964), the Vaccinio-Piceetea class and the Pinetalia *sylvestris* order are recognized inside the Iberian Peninsula. Moreover, the plant communities with *Echinopartum barnadesii* and *Senecio carpetanus* are interpreted as geographical races. Other aspects of the juniper communities of Junipero *nanae*-Cytisetum *oromediterranei* (subassociations, variants and relic forms) are also races. The presence of *P. nigra* subsp. *salzmannii* inside the forests of *Quercus pyrenaica* is considered as a thermic relic form, and the communities with *P. sylvestris* as altitudinal forms.

Key words: Phytosociology, multivariate analysis, Sistema Central, Iberian Peninsula, pine forests.

Introduction

In Europe the natural communities with *Pinus subsectio sylvestris* (Little & Critchfield 1969) have a boreal origin (Jan du Chene 1976, Bauerochse & Katenhusen 1997, Millar 1993). Their expansion was favoured by the glacial periods (Costa Tenorio et al. 1988, Cristina Peñalba 1994, Wilmanns 1997). Presently, *Pinus sylvestris* L. is a widely distributed species in eurosiberian Europe, with some radiations in the Mediterranean region (Hultén & Fries 1986). On the other hand, *P. nigra* Arnold, is distributed in the mountains near the Mediterranean sea (Wendelberger 1963, Blanco Castro et al. 1996), diversifying in several subspecies taxa.

Some contributions to the ecology and phytosociology of *P. sylvestris* and *P. nigra* subsp. *salzmannii* (Dunal) Franco in the Iberian Peninsula have been noted: Rivas Goday 1955, Rivas Goday & Borja Carbonell 1961, Rivas-Martínez 1963, 1964, Esteve Chueca 1973–74, Vigo 1979, Losa Quintana et al. 1986, Rivas-Martínez & Cantó 1987, Rivas-Martínez et al. 1987, 1991, Gamisans & Gruber 1988, Valle et al. 1988, Regato & Escudero 1989, Elena Rosselló & Sánchez Palomares 1991, Fernández-González 1991, Gamisans et al. 1991, Ninot 1996, Rojo y Alboreca & Montero González 1996. They have been considered for the construction of the current syntaxonomic scheme in this article.

P. sylvestris is one of the most important forest species in Spain and is widely distributed in the Sistema Central, while *P. nigra* subsp. *salzmannii* is only a relic form in this territory. As Costa Tenorio et al. (1990) comment, the interpretations of the vegetation where the former is included are much too strict if the classic aspect of the phytosociological method is considered (Rivas-Martínez 1987). Some authors point out the absence of *P. sylvestris* (Sánchez Mata 1989) in areas visited by other authors (Mancebo et al. 1993) who allege its existence, as in the case of the Gredos Mountains. The communities with *P. nigra* subsp. *salzmannii* have been known in the Sistema Central for a long time, including in the Guadarrama Mountains (Gómez Manzaneque 1988, Regato et al. 1992). However, there has been no phytosociological interpretation (Rivas-Martínez 1975, Regato et al. 1995).

The aim of the present study is precisely to show a new phytosociological approximation which explains the ecology of these communities with *P. sylvestris* and *P. nigra* subsp. *salzmannii* in the Sistema Central, in the context of the coniferous European forests.

Phytogeography of the area studied

The Sistema Central are the siliceous Paleozoic mountains which go through the Iberian Peninsula (Fig. 1), from W-SW to E-NE from the Estrela Mountains (Portugal) to the Ayllón and Las Cabras mountains (Guadalajara-Segovia-Soria, Spain). The highest peaks are: Estrela (1891 m), La Ceja (2425 m), Peña de Francia (1723 m), Calvitero (2401 m), Pico de Almanzor (2592 m), Cabeza de Hierro (2383 m), Pico de Peñalara (2489 m), Ocejón (2058 m) and Pico del Lobo (2273 m).

The Sistema Central belongs to the Iberomarroqui-Atlantica superprovince (Pérez Latorre et al. 1996, Deil & Galán de Mera 1998). This encompasses the areas of the Iberian Peninsula and northern Africa with Atlantic-Mediterranean climatic regime (Gausson et al. 1958), according to the distribution of Atlantic elements in northern Africa (Dahlgren & Lassen 1972). This superprovince is a migratory space of Atlantic eurosiberian species to the southern Iberian Peninsula and northern Africa. From the 8 provinces in which this wide Iberic-Northafrican territory is divided (Carpetano-Iberico-Leonesa, Luso-Extremadurese, Tingitano-Onubo-Algarviense, Betica, Rifeña, Atlasica, Atlantica and Sud-Occidental), our area studied is encompassed in the Carpetano-Iberico-Leonesa province (Sistema Central, León, Orense and Soria mountains). This province is divided into various sectors (Rivas-Martínez et al. 1990): A – Guadarramico sector (Guadarrama Mountains), B – Bejarano-Gredense sector (Bejar and Gredos mountains), C – Salmantino sector (Peña de Francia and plains of Salamanca) and D – Estrellense sector (Estrela Mountain). The localities where relevés have been made are included in the Guadarramico and Bejarano-Gredense sectors (Fig. 1).

Following the bioclimatic classification of Rivas-Martínez et al. (1991), the data obtained from Müller (1982), and the National Institute of Meteorology (Spain), and the discriminatory indexes, Summer and Winter Humidity of Galán de Mera et al. (1995), the Sistema Cen-




Fig. 1. Map of the Spanish Sistema Central, phytogeography and localities studied (following several references and relevés made by the authors). Symbols: □ Junipero nanae-Cytisetum oromediterranei geographical race with *Senecio carpetanus pinetosum sylvestris*, ▲ Junipero nanae-Cytisetum oromediterranei geographic race with *Echinopartum barnadesii pinetosum sylvestris* altitudinal form with *Pinus nigra* subsp. *salzmannii*, ● *Pinus sylvestris* DC, ◻ *Luzulo forsteri-Quercetum pyrenaicae* relic form with *Pinus nigra* subsp. *salzmannii*, ■ *Genisto falcatae-Quercetum pyrenaicae* relic form with *Pinus nigra* subsp. *salzmannii*. The numbers indicate the relevés of Table 3. A – Guadarramico sector, B – Bejarano-Gredense sector, C – Salmantino sector.

tral is included in the Mediterranean Region (Table 1), which ranges from subhumid to hyperhumid (Table 2).

The diagonal position which the Sistema Central presents in the Iberian Peninsula implies different perception of temperature and rainfall in the mountains, depending on the localities. The Gredos Mountains and the northern slopes are more influenced by the Atlantic disturbances. Moreover, the Ayllón Mountains receive the rainfall of the Mediterranean summer-autumn low pressures as a consequence of their distance from the Azores anticyclone (Capel Molina 1981). This causes the southern slopes of the Guadarrama Mountains to be the driest and also the most continental, because of the higher contrast between temperatures. Therefore, this is the area of wide distribution of *P. sylvestris*. However, *P. nigra* subsp. *salzmannii* is only found in the Gredos Mountains and some western sites of the Guadarrama Mountains because of its high thermic exigences. The bioclimatic belts are displaced in the Sistema Central because of the thermic differences between the slopes. Thus, for example, on the southern slope of the Guadarrama Mountains, the Oromediterranean belt extends from 1500 to 2300 m, while on the northern slope it descends to 1300 m.

Table 1. Values of the summer (HE) and winter (HI) humidity indexes in European meteorological stations and of the Sistema Central. Eurosiberian Region $PI > PE/HE > 1$; Mediterranean region $PI > PE/HE < 1$; $HI = \sum_{D-M} (P + HR/ETP)$; $HE = \sum_{J-A-S} (P + HR/ETP)$; P: rainfall in mm, HR: relative humidity of the air in %, ETP: potential evaporation in mm.

METEOROLOGICAL STATION	ALTITUDE (m)	P(D,Jan,F,M)	P(J,J,A,S)	HR(D,Jan,F,M)	HR(J,J,A,S)	ETP(D,Jan,F,M)	ETP(J,J,A,S)	HI	HE	CLIMATE TYPE AND PLANT FORMATIONS
Madrid (Spain), 40°25'N/3°41'W	667	165	85	298	208	66	479	7,0	0,6	Mediterranean, Sclerophyllous vegetation
Lyon (France), 45°43'N/4°57'E	200	208	323	324	279	57	453	9,3	1,3	Temperate climate, Deciduous forests
Genève (Switzerland), 46°12'N/6°09'E	405	233	346	319	282	37	417	14,9	1,5	Temperate climate, Deciduous forests
Zürich (Switzerland), 47°23'N/8°34'E	569	275	518	312	276	25	399	23,5	1,9	Temperate climate, Deciduous forests
Stuttgart (Germany), 48°42'N/9°12'E	401	161	311	324	301	39	422	12,4	1,4	Temperate climate, Deciduous forests
Nürnberg (Germany), 49°30'N/11°06'E	310	159	282	328	292	23	412	21,2	1,4	Temperate climate, Deciduous forests
Praha (Czech Republic), 50°05'N/14°25'E	197	96	251	335	288	19	430	22,7	1,3	Temperate climate, Deciduous forests
Dresden (Germany), 51°07'N/13°41'E	246	148	297	318	292	22	407	21,2	1,4	Temperate climate, Deciduous forests

Table 2. Climatic values of some meteorological stations near the localities studied with *Pinus sylvestris* and *P. nigra* subsp. *salzmannii*. Symbols: A = altitude of the meteorological station (m), It = thermicity index (T + M + m) 10, T = annual mean temperature (°C), M = maximal mean temperatures of the coldest month (°C), m = minimal mean temperature of the coldest month (°C), P = annual mean rainfall (mm), H = number of days with sure frost, N = number of days with snow covering. * Only pluviometric stations.

METEOROLOGICAL STATION	A	It	T	M	m	P	H	N	BIOClimATIC DIAGNOSIS
Cerezo de Arriba "Gran Plato" (Segovia)	1880	83,1	7,0	3,5	-2,2	1218,4	36	30-31	Oromediterranean humid
Puerto de Navacerrada (Madrid)	1890	48,7	6,2	1,9	-3,3	1335,4	45	115-116	Oromediterranean humid
Cercedilla "Fuenfria" (Madrid) *	1350	—	—	—	—	1121,3	—	6-7	Humid
Guadarrama (Madrid)	981	107,8	6,7	5,7	-1,7	655,3	15	4-5	Supramediterranean subhumid
Valle de los Caidos (Madrid) *	1300	—	—	—	—	980,5	—	16-17	Subhumid
Navalperal de Pinares (Ávila) *	1287	—	—	—	—	704,7	—	7-8	Subhumid
San Bartolomé de Pinares (Ávila) *	1150	—	—	—	—	416,9	—	39	Dry
Serranillos (Ávila) *	1235	—	—	—	—	1366,6	—	8	Humid
Puerto del Pico (Ávila) *	1395	—	—	—	—	946,8	—	11-12	Subhumid
Embalse La Jarosa (Madrid)	1060	165	10,7	7,1	-1,3	938,6	9-10	3-4	Supramediterranean subhumid
Navarredonda de Gredos (Ávila)	1525	119	8,5	5,9	-2,5	828,4	37	22-23	Supramediterranean subhumid

Materials and methods

Treatment of the data.

The present study is based on 683 phytosociological relevés made following the method of Braun-Blanquet (1964). These relevés are located in the mountains of western Europe and northern Africa, to establish a comparison with those of the Sistema Central. From these relevés, 131 correspond to the area studied, and are represented in Table 3. Table 5 is a synthetic table of the communities of the *Vaccinio-Piceetea* class in western Europe and northern Africa, with the different species of *Pinus* of the mountain pine groves of the Iberian Peninsula (*P. sylvestris*, *P. nigra* subsp. *salzmannii* and *P. uncinata*). The references of Table 3 are in Table 4, and those of Table 5 in Table 6.

We have made statistical analyses of Tables 3 and 5, with the SYN-TAX program (Podani 1994). Dendrograms to observe the degree of similarity between relevés were obtained applying the Jaccard index (1929) (Figs. 2 and 3). After this, the tables were put in order considering the relationship between relevés, to study the ecological and phytosociological behaviour.

Phytosociological treatment

The conceptual treatment of the phytosociological association and its subdivision made by the Iberian phytosociologists have limited ecological facts to a rigid syntaxonomic scheme and forced the placing of some plant communities in specific associations. The same occurs with the subassociation, which, since its definition by Braun-Blanquet (1964), has been interpreted in several ways. It has been used to design altitudinal, geological, ecological and edaphic variations. These induce us to consider the studies of Kopecký & Hejný (1974), Foucault (1981), Dierschke (1993) and Kopecký et al. (1995), which explain the basal community (BC) and derived community (DC) concepts. A basal community is a plant community sited in anthropogenic places and colonized by plants of the highest syntaxonomic unities. The derived community means the invasion of one association by plants different from the characteristic ones, and whose number decreases considerably because of a derived change in human activity. The concept of relic form (as opposed to normal historic form), given by Schuhwerk (1990), is used to distinguish small regions with critical plants which lend a historic and ancestral aspect to the association, for example, the result of a low exposure to past glacial periods (Ojeda et al. 1995). This concept was used in Spain to explain some aspects of the Aragonese relic vegetation (Montserrat 1975), without, however, a phytosociological viewpoint.

The interpretation of the variability of the association by Matuszkiewicz (1981) leads us to define the meaning of subassociation more accurately. This author considers vegetation as a relative continuum and concludes that the association has 3 variants (Fig. 5): 1 – horizontal-referring to the geographical races, 2 – vertical-referring to the altitudinal forms, and 3 – local-referring to the edaphic differences, considering this as the concept of subassociation.

Results and discussion

Vaccinio-Piceetea and Pino-Juniperetea

The *Vaccinio-Piceetea* class encompasses the coniferous forests and the heaths with boreal origin, characteristic of continental areas (Julve 1993, Ellenberg 1996). According to Rivas-Martínez et al. (1991), they were developed in northeastern Europe and the southern Alps after the Tardiglacial, as a wooded tundra. Braun-Blanquet et al. (1952) and Rivas Goday & Borja Carbonell (1961) consider that this phytosociological class exists inside the Iberian Peninsula. Rivas-Martínez (1963) maintains that the pine groves of the Sistema Central do not belong to the *Vaccinio-Piceetea*, and have to be included in the *Cytision oromediterranei* alliance because of the importance of the *Genistea* in our mountains and the impoverishment in some characteristic plants of the class. Later Rivas-Martínez (1964) created the *Pino-Juniperetea* class to isolate the Mediterranean pine groves from the centereuropean ones.

Table 3. (continued)

Relevé number	1	11	1 1 1	11111111111 1	11	1	11111111111
3444461888883333	4444	99999900111112	22222223333311515161669990000101011	011	56111415652777745678222233222567557756767888		11111111111
15168727146823697836752534202145634867890431724569208134350516185689702136370912850293749194532717830909014689012305445784669280798							
Arctostaphylos uva-ursi	2	2443333315243443	2	1133241+32			
Halimium umbellatum viscosum	3111	222332231.2	2	++21211			
Pteridium aquilinum	222312	23	4	2	1224		++
Gallium rotundifolium	11212311			12121			2+11
Cistus laurifolius		122121	31113				
Quercus-Fagetea CL	221	1.2	+				1111
Genista florida	+	1	+	+			
Genista cinerea	1	+	+	+	1		
Veronica officinalis	1	+			11		
Viola riviniana	11+1						
Poa nemoralis	11+1111						
Clinopodium vulgare arundanum	+	1	+				
Viola odorata	+	1	+				
Populus tremula	+	2111					
Gallium rivulare	22		++				
Rubus idaeus	++						
Sanicula europaea	111						2.2
Holcus mollis	12.2						
Melica uniflora		121					
Rosa corymbifera	1						1+
Fragaria vesca	2	2					
Campanula rapunculoides		1	1				
Helleborus niger	+	1					
Aconitum napellus castellanum		1+					
Athyrium filix-femina		11					
Betula pendula		++					
Rosa tomentosa	+	+					
Sorbus aucuparia							
Crataegus monogyna		+					
Hepatica nobilis		1					
Geum sylvaticum							
Rosa micrantha							
Oriqanum vulgare							
Paris quadrifolia		+					
Conopodium filifolium							
Rubus ulmifolius		1					
Brachypodium sylvaticum							
Rosa canina							
Mosses and lichens							
Hypnum cupressiforme	1						
Moss	1		21.1				
Dicranum pellucidum	1						
Pseudoevernia furfuracea	11						
Bartramia halleriana		+					
Polytrichum sp.		11					
Polytrichum piliferum							
Cladonia furcata							
Cladonia fimbriata							
Companions							
Erica arborea	122.2	+.1+	432.3331+.1.11333.1	11.1+	++22+232	+.4	2112

Table 3. (continued)

Relevé number	1	11	111	1111	11111	111111	1111111	11111111	111111111	1111111111	11111111111
Avenula sulcata s. str.											
Arenaria montana	34444618888883333	4444	99999990011112	22222223333311515161669990000101011	011	5611414565277774	567822222332222567557756767888				
Rumex angiocarpus	15168727146523697836752534202145634867890431724569208134350516185689702136370912850293749194532717830909014689012305445784669280798										
Festuca indigesta aragonensis											
Agrostis castellana											
Arrhenatherum album											
Thymus bracteatus											
Nardus stricta											
Adenocarpus hispanicus											
Conopodium pyrenaicum											
Hieracium pilosella											
Lavandula pedunculata											
Jasione laevis carpetana											
Carduus carpetanus											
Solidago virgaurea											
Agrostis capillaris											
Rumex acetosella											
Leucanthemopsis pallida alpina											
Lactuca viminea											
Cerastium gracile											
Thymus praecox											
Pinus pinaster											
Hypochaeris radicata											
Cytisus scoparius											
Agrostis tenuis											
Arrhenatherum elatius s.l.											
Hieracium vahlii											
Cruciata glabra											
Festuca iberica											
Agrostis delicatula											
Hieracium castellanum											
Sedum forsterianum											
Jasione montana echinata											
Helianthemum apenninum											
Teucrium scorodonia											
Further do occur:											
Erica australis 45:2, 46:4, 48:4, 62:1; Festuca ampla 35:2, 59:1, 58:1, 61:2; Leontodon crispus bourgeanus 50:4, 76:4, 101:4, 115:4; Sedum brevifolium 13:1, 123:4, 127:4, 130:4; Gagea quadarramica 11:4, 124:4, 125:1, 129:4; Crocus carpetanus 3:4, 71:4, 77:4, 78:4; Leontodon carpetanus 128:4, 129:1, 130:1, 131:4, 131:4; Armeria caespitosa 128:4, 129:4, 130:4; Omalotheca sylvatica 82:1, 83:1, 123:1; Cladonia pyxidata 122:1, 123:1, 131:1, 131:1; Lactuca chondrillifolia 57:4, 58:1, 120:4; Dianthus laricifolius 103:1, 105:1, 109:4; Leucanthemopsis pallida s. str. 94:4, 95:4, 96:1; Gentiana lutea 10:2, 78:4, 102:4; Carex muricata 82:4, 83:1, 86:4; Centaurea alba 18:4, 19:4, 55:4; Thymus zygis 21:4, 22:1, 27:4; Urtica dioica 23:4, 43:1, 44:1; Rosa sp. 23:4, 24:4, 53:4; Jurinea humilis 61:1, 103:1, 106:1; Ranunculus ollisiponensis s. str. 120:4, 123:4; Senecio sp. 43:4, 47:1; Oxalis corniculata 43:1, 44:1; Senecio jacobaea 58:4, 59:1; Thymus mastichina 51:1, 55:2; Picris hieracioides 45:4, 46:4; Dactylis glomerata 26:1, 85:4; Festuca rothmaleri 81:2, 84:1; Festuca agr. rubra 82:4, 83:1; Acinos alpinus micranthum 109:4, 110:4; Ranunculus ollisiponensis alpinus 4:4, sect. Histrices 82:2, 83:1; Geranium robertianum 85:4, 86:4; Narcissus rupicola 9:4, 12:4; Sedum album micranthum 109:4, 110:4; Ranunculus ollisiponensis alpinus 4:4, spicant 44:1; Lapsana communis 62:1; Epilobium lanceolatum 81:4; Conopodium majus burgaei 122:1; Quercus rotundifolia 2:1; Galeopsis ladanum 82:4; Mucizonia sedoides 49:1; Blechnum sect. Histrices 82:2, 83:1; Geranium robertianum 85:4, 86:4; Potentilla micrantha 85:4; Quercus rotundifolia 2:1; Quercus setigiflorus 55:2; Juncus hirsutum 51:4; Avenula sulcata 52:4; Silene saxifraga 49:4; Taraxacum sroeterianum 12:4; Saxifraga willkommia 116:4; Holcus setigiflorus 44:4; Viola sp.? 47:4; Sedum hirsutum 51:4; Avenula sulcata albinervis 58:1; Pentaglottis sempervirens 85:1; Andryala integrifolia 25:4; Geranium purpureum 23:1; Thlaspi stenopterum 5:4; Juncus conglomeratus 44:4; Artemisia glutinosa 45:4; Spergula morisonii 51:4; Pinus pinaster (young) 25:4; Jasione crispa centralis 10:4; Festuca costei 5:3; Scutellaria minor 44:1; Calluna vulgaris 45:1.											

Table 4. References of the relevés of Table 3.

- 1, 2. Relevés of the authors. Valle de Enmedio, Malagón Mountain, Ávila
- 3-8. Relevés of the authors. Cueva Valiente, Malagón Mountain, Ávila
- 9-12. Relevés of the authors. Peñalara, Guadarrama Mountains, Madrid
- 13, 14. Relevés of the authors. Puerto de Navacerrada, Guadarrama Mountains, Madrid
15. Relevés of the authors. Bola del Mundo, Guadarrama Mountains, Madrid
- 16-20. Relevés of the authors. Calle de los Álamos, Guadarrama Mountains, Madrid
- 21-27. Relevés of the authors. Cabeza de la Parra, Gredos Mountains, Ávila
- 28-34. Relevés of the authors. El Arenal, Gredos Mountains, Ávila
- 35-41. Relevés of the authors. Los Leones, Guadarrama Mountains, Madrid-Segovia
- 42-49. Relevés of the authors. La Pinilla, Ayllón Mountain, Segovia
50. Rivas-Martínez et al. (1987). Peñalara, Guadarrama Mountains, Madrid
51. Rivas-Martínez et al. (1987). Cerro del Telégrafo, Guadarrama Mountains, Madrid
52. Rivas-Martínez et al. (1987). From Valdesquí to Valdemartín, Guadarrama Mountains, Madrid
53. Rivas-Martínez et al. (1987). Pico del Nevero, Guadarrama Mountains, Madrid
- 54, 97. Rivas-Martínez et al. (1987). Puerto de Navafría, Guadarrama Mountains, Madrid
55. Rivas-Martínez et al. (1987). Lomo del Noruego, from Cotos to Valdesquí, Guadarrama Mountains, Madrid
56. Rivas-Martínez et al. (1987). From puerto de la Morcuera to Las Najarras, Guadarrama Mountains, Madrid
- 57, 58. Rivas-Martínez et al. (1987). Puerto de Navafría, Guadarrama Mountains, Madrid
- 59, 60, 62. Rivas-Martínez et al. (1987). Circo de Hoyo Cerrado, Peñalara, Guadarrama Mountains, Madrid
61. Rivas-Martínez et al. (1987). El Nevero, Guadarrama Mountains, Madrid
- 63, 65, 66, 98. Fernández-González (1991). Puerto de Cotos, Guadarrama Mountains, Madrid
- 64, 73-76. Fernández-González (1991). Puerto de Navafría, Guadarrama Mountains, Madrid
- 67, 118. Fernández-González (1991). El Nevero, Guadarrama Mountains, Madrid
68. Fernández-González (1991). Northern slope of Mondalindo, Riofrío, Guadarrama Mountains, Madrid
- 69, 70, 72. Fernández-González (1991). Puerto de la Morcuera, Guadarrama Mountains, Madrid
- 71, 77. Fernández-González (1991). Lomo del Noruego, from Cotos to Valdesquí, Guadarrama Mountains, Madrid
78. Fernández-González (1991). From Cotos to Valdesquí, Guadarrama Mountains, Madrid
- 79, 80, 117. Fernández-González (1991). Circo de Hoyo Cerrado, Guadarrama Mountains, Madrid
- 81, 84. Fernández-González (1991). Cabeza Mediana, Guadarrama Mountains, Madrid
- 82, 83. Fernández-González (1991). Southern slopes of Los Pájaros, Guadarrama Mountains, Madrid
85. Fernández-González (1991). From Lozoya to Navafría, Guadarrama Mountains, Madrid
86. Fernández-González (1991). Arroyo de las Guarramillas, Cotos, Guadarrama Mountains, Madrid
87. Rivas-Martínez and Cantó (1987). Tablada, Malagón Mountain, Madrid
- 88, 90. Rivas-Martínez and Cantó (1987). Cabeza de Lijar, Malagón Mountain, Madrid
89. Rivas-Martínez and Cantó (1987). Abantos, Malagón Mountain, Madrid
91. Rivas-Martínez and Cantó (1987). Pinares Llanos, Malagón Mountain, Madrid

Table 4. (continued)

- 92, 94. Rivas-Martínez and Cantó (1987). Collado de la Mina, Guadarrama Mountains, Madrid
93. Rivas-Martínez and Cantó (1987). La Salamanca, Guadarrama Mountains, Madrid
95. Rivas-Martínez and Cantó (1987). Collado del Hornillo, Malagón Mountain, Ávila
96. Rivas-Martínez and Cantó (1987). Cueva Valiente, Malagón Mountain, Ávila
99. Rivas-Martínez et al. (1987). Somosierra, Guadarrama Mountains, Madrid
101. Rivas-Martínez et al. (1987). Hoyo de Pepe Hernando, Peñalara, Guadarrama Mountains, Madrid
102. Rivas-Martínez et al. (1987). Dos Hermanas, Peñalara, Guadarrama Mountains, Madrid
- 103, 113. Rivas-Martínez et al. (1987). La Salamanca, Guadarrama Mountains, Ávila
- 104, 105, 107-110. Rivas-Martínez et al. (1987). Cueva Valiente, Malagón Mountain, Ávila
106. Rivas-Martínez et al. (1987). Collado de la Mina, Guadarrama Mountains, Madrid
- 100, 111, 112. Rivas-Martínez et al. (1987). Cabeza de Lijar, Malagón Mountain, Madrid
114. Rivas-Martínez et al. (1987). From puerto de Canencia to Collado Cerrado, Guadarrama Mountains, Madrid
115. Fernández-González (1991). Peñalara, Guadarrama Mountains, Madrid
116. Fernández-González (1991). Dos Hermanas, Guadarrama Mountains, Madrid
119. Fernández-González (1991). From puerto de Canencia to Collado Cerrado, Cotos, Guadarrama Mountains, Madrid
120. Rivas-Martínez (1963). Collado Albo, Guadarrama Mountains, Madrid
121. Rivas-Martínez (1963). Barranco del río Moro, Guadarrama Mountains, Segovia
123. Rivas-Martínez (1963). Puerto de la Fuenfría, Guadarrama Mountains, Madrid
124. Rivas-Martínez (1963). Puerto de Navafría, Guadarrama Mountains, Madrid
125. Rivas-Martínez (1963). El Ventorrillo, Guadarrama Mountains, Madrid
126. Rivas-Martínez (1963). Puerto de Cotos, Guadarrama Mountains, Madrid
127. Rivas-Martínez (1963). Dos Hermanas, Peñalara, Guadarrama Mountains, Madrid
128. Rivas-Martínez (1963). Western slope of Siete Picos, Guadarrama Mountains, Madrid
129. Rivas-Martínez (1963). Eastern slope of Siete Picos, Guadarrama Mountains, Madrid
130. Rivas-Martínez (1963). Northern slope of Cerro del Telégrafo, Guadarrama Mountains, Madrid
131. Rivas-Martínez (1963). Northern slope of Siete Picos, Guadarrama Mountains, Madrid




Fig. 2. Similarity dendrogram of the communities of *Pinus sylvestris* and *P. nigra* subsp. *salzmannii* of the Sistema Central.




Fig. 3. Similarity dendrogram of the communities of Vaccinio-Piceetea in western Europe and of Querceto-Fagetea with *Pinus sylvestris* and *P. nigra* subsp. *salzmannii*.

In Table 4 we can observe that there are several eurosiberian plants, frequent in the Vaccinio-Piceetea, which extend to the Iberian Peninsula (Font Quer 1954, Ceballos 1966): *Arc-tostaphylos uva-ursi*, *Juniperus communis*, *J. sabina*, *Pinus sylvestris*, *P. uncinata*, *Pyrola chlorantha*, *P. minor*, *Vaccinium myrtillus*. Though the Genisteae are highly diversified in the Mediterranean Region (Cristofolini 1997), and show a very different aspect from the Mediterranean communities, the distribution of *J. communis* and *P. sylvestris* is so evident in the boreal world (Hultén & Fries 1986) that it is not necessary to create another phytosociological class in the Iberian Peninsula. In the mountains of the Anatolian Peninsula (Turkey), there is a similar occurrence with the Abietion bornmuellerianae alliance (Rehder et al. 1994).

Group *a* from Table 4 consists of characteristic elements of the *Querceto-Fagetea*, whose presence is higher in the Iberian and Pyrenean pine groves, and so we include these in the Pinetalia *sylvestris* order (Folch i Guillén 1986, Oberdorfer 1990, Bolòs et al. 1993). This order encompasses the mountain pine groves which extend from the center of Europe to the Mediterranean Basin. On the other hand, in group *b*, the frequency of boreal elements is higher, defining the *Piceetalia* order (*Clematis alpina*, *Erica herbacea*, *Homogyne alpina*, *Larix decidua*, *Linnaea borealis*, *Listera cordata*, *Picea abies*, *Pinus cembra*, *Trientalis europaea*...). The Vaccinio-Piceetea class is divided into 2 orders: Pinetalia *sylvestris*-pine groves which go from Central Europe to the Iberian Peninsula, contacting with elements of Querceto-Fagetea class, and *Piceetalia*-subalpine and alpine coniferous forests which go from boreal Europe to the Pyrenees.




Fig. 4. Photo of a basal community (BC) developed in a track of a chair lift, a derived community (DC), because pine groves are favoured by crops, and natural pine groves of *Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris* (JC) [La Pinilla, Segovia].

The communities with Pinus sylvestris and Pinus nigra subsp. salzmannii of the Sistema Central – Luzula lactea-Pinus sylvestris Basal Community (BC)

In Figure 4 there is a photo of a *Pinus sylvestris* climax pine grove in the center of Spain. A great part of the forest was destroyed to install a chair lift of a ski resort. Sometime later, this strip was invaded by *Luzula lactea*, while in nearby unaltered zones, there are the elements of *Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris*. Thus, the community with *Pinus sylvestris* and *Luzula lactea* is a basal community (BC).

Junipero nanae-Cytisetum oromediterranei

It is a siliceous community dominated by *Cytisus balansae* subsp. *europaeus*, whose origin is the adaptation of some Genisteae to the high mountain during the Alpine orogeny. The characteristic floristic composition includes *Cytisus balansae* subsp. *europaeus*, *Deschampsia flexuosa* subsp. *iberica*, *Juniperus communis* and *Luzula lactea*. From a phytotopographical point of view, it is sited on central and lateral moraines and ice fields of the Sistema Central, between 1500 and 2500 m, on the southern slope, and 1300 to 2500 m on the northern slope. Pine groves appear in the deepest and most humid soils.

From a chorological point of view, it is a Carpetano-Iberico-Leonesa association with different geographical races (Schuhwerk 1990) based on microendemisms such as: *Senecio carpetanus* (= *Senecioni carpetani-Cytisetum oromediterranei*, Guadarrama Mountains), *Echinopartum barnadesii* (= *Cytiso oromediterranei-Echinopartum barnadesii*, Gredos Moun-




Fig. 5. Variability of the associations with *Pinus sylvestris* and *P. nigra* subsp. *salzmannii* in Navacerada-Madrid (1) and Pico pass-Ávila (2) following the association concept of Kopecký and Hejný (1974), Matuszkiewicz (1981) and Schuhwerk (1990): a) *Junipero nanae-Cytisetum oromediterranei*, b) *Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris*, c) *Pinus sylvestris* DC, d) *Luzulo forsteri-Quercetum pyrenaicae*; a') *Junipero nanae-Cytisetum oromediterranei* geographic race with *Echinopartum barnadesii*, b') *Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris* altitudinal form with *Pinus nigra* subsp. *salzmannii*, e) *Genisto falcatae-Quercetum pyrenaicae*, f) *Genisto falcatae-Quercetum pyrenaicae* relic form with *P. nigra* subsp. *salzmannii*.

tains), *Echinopartum barnadesii* subsp. *dorsisericeum* (= *Echinoparto pulviniformis-Cytisetum oromediterranei*, Gredos Mountains and Peña de Francia), *Teucrium salviastrum* (= *Lycopodio clavati-Juniperetum nanae*, Estrela Mountains).

Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris

This encompasses the pine groves of the Guadarrama and the Gredos mountains, sited on more developed soil than the typical association, with a similar floristic court, though there are some important plants such as *Adenocarpus hispanicus*, *Linaria nivea*, *Luzula lactea* or *Vaccinium myrtillus*.

We can deduce from Table 3 that the altitudinal limit of *Pinus sylvestris* is quite variable, depending on the relief. For example, we can find it below 1900 m in the Ayllón Mountains (Segovia), while near Peñalara (Madrid), it can be found below 2200–2300 m, depending on the highest continentality of the locality. (Fig. 5).

The typical association, *Junipero-Cytisetum oromediterranei*, is a substitution stage of the natural pine groves, and it is a perennial community where *P. sylvestris* cannot grow.




Fig. 6. *Junipero nanae*-*Cytisetum oromediterranei* geographic race with *Echinospartum barnadesii* pinetosum *sylvestris* and altitudinal form with *Pinus nigra* subsp. *salzmannii* (El Arenal, Sierra de Gredos, Avila).

The variability of these communities is high if we also consider the development of pines in the belt of the oak grove. In Table 3 we point out the variant with *Linaria nivea*. This plant, together with *Digitalis purpurea*, indicates soils mainly altered by fire. The pine groves were cleared a long time ago to obtain pasturages for cattle (Gil García et al. 1996). These pine groves belong to the *Koelerio-Corynephoretea* class and define a subserial variant with *Koeleria crassipes* and *Corynephorus canescens*. In some relevés of Table 3, the presence of *Vaccinium myrtillus*, which is not very frequent in the Guadarrama Mountains, is important. This plant is more abundant in the Ayllón Mountains, in the pine groves above 1620 m (Pinilla ski resort, Segovia); on the other hand, in Madrid it appears above 2000 m (Telégrafo hill). This Ericaceae can always be found exposed to the cold northern winds or protected in the glacial cirques, defining a relic form of cooler times.

Though Font Quer (1954) cited *Pinus sylvestris* in the Gredos Mountains, it seems that some authors do not find it in the Sistema Central (Rivas-Martínez et al. 1987, Sánchez Mata 1989). Our relevés have been made near the Pico peak (Gredos, Avila), over 1300 m. Most of these localities are on the southern slope. Here *Pinus nigra* subsp. *salzmannii* is found frequently, giving rise to an altitudinal form of more thermic exposures. (Fig. 6). The ecology and phytogeography of these localities have already been defined by Regato Pajares et al. (1992), Mancebo et al. (1993), Regato et al. (1995).

Table 5. Synthetic Table of the communities of Vaccinio-Piceetea in western Europe and of Quercu-Fagetea with *Pinus sylvestris* and *P. nigra* subsp. *salzmannii* (the Arabic numbers indicate the presence: r < 6%, + = 6-10%, 1 = 11-20%, 2 = 21-40%, 3 = 41-60%, 4 = 61-80%, 5 = > 81%). Abbreviations: AL = alliance, O = order, CL = class.

Community number	3	527	93	411499	3	344432779999	616666666661441117777772441633233116733425525299888888258888882552555	37470796658001378261587899678012393601264572123475623415455801928990346012138455678949012342453678					
Relieves per column	1	1	1	1121	1121111111	1	111	1111	123	1	11		
463181-77325862786725127212-----												15565-6	
Species number	111	11132123443521121111	3664545454964445436225356657684358223323245253343212353765551123354	2153216426020593781602737718775792547047231077928954831645781844858618368876906832258734391868038233039									
Vaccinio-Piceetea CL													
Pinus sylvestris	443115133444444444234555123235544523332	3	431414531551424	1	3					1	212	22555555	
Juniperus communis	4431512	2	413243234	45551151455445123454532	34455255	44434423134	21	144	42			25	253
Vaccinium myrtillus													
Pinus uncinata													
Arctostaphylos uva-ursi													
Pinus x thaetica													
Vaccinium vitis-idaea													
Pinetalia sylvestris O, Cytision oromediterranei AL	443	2314241432	34531										
Cytisus balaneae europaeus													
Luzula lactea													
Deschampsia flexuosa iberica													
Juniperus sabina													
Juniperus thurifera													
Genista lobelii longipes													
Echinopatum barnadesii													
Piceetalia O													
Orthilia secunda													
Festuca gautieri													
Pyrola chlorantha													
Homogyne alpina													
Rhododendron ferrugineum													
Cotoneaster integerrimus													
Pyrola minor													
Larix decidua													
Luzula nivea													
Monotropa hypopitys													
Moneses uniflora													
Pinus cembra													
Vaccinium uliginosum microphyllum													
Goodyera repens													
Polygala chamaebuxus													
Loniceria nigra													
Picea abies													
Erica herbacea													
Luzula sieberii													
Listera cordata													
Empetrum nigrum hermaphroditum													
Loiseleuria procumbens													
Huperzia selago													
Blechnum spicant													
Chamaecytisus supinus													
Arctostaphylos alpinus													
Daphne striata													
Coronilla vaginialis													
Bupthalamum salicifolium													
Clematis alpina													




Tab. 5. (continued)

Community number				
3	527	93	411499	3
37470796658001378261587899678012393601264572123475623415455801928990346012138445678949012342453678				
Linnaea borealis				154
Lycopodium annotinum				153
Luzula luzulina				131
Aquilegia atrata				313
Trientalis europaea				145
Lonicera alpigena				2
Lonicera coerulea				11
Daphne cneorum				112
Galium pinetorum				123
Rhododendron hirsutum				15
Pyrola rotundifolia				34
Vaccinium uliginosum pubescens				25
Crepis alpestris				2
Monotropa hypophega				1
Corallorhiza trifida				1
Carex ericetorum				21
Vaccinium uliginosum s.str.				13
Rhododendron intermedium				1
Epipactis atrorubens				2
Thesium rostratum				1
Chamaecytisus ratibonensis				4
Mosses and lichens				
Hylacomium splendens				14
Pleurozium schreberi				11
Dicranum scoparium				14
Rhytidiadelphus triqueter				13
Hypnum cupressiforme				22
Cetraria islandica				55
Tortella tortuosa				23
Polytrichum formosum				33
Leucobryum glaucum				1
Dicranum bergeri				1111
Ptilium crista-castrensis				5555
Mnium spinosum				1213
Cladonia furcata				41222
Plagiochila asplenioioides				241
Scleropodium purum				3222
Peltigera gr. aphchosa				33
Cladonia pyxidata				154
Barbilephozia lycopodioides				222
Pseudoscleropodium purum				121
Ctenidium molluscum				1
Peltigera canina				4
Rhytidiadelphus loreus				23
Eurynchium striatum				11
Ditrichum flexicaule				33
Cladonia foliosa				21
Homalothecium sericium				32
Hylacomium proliferum				32
Polytrichum piliferum				1
Polytrichum juniperum				1
Polytrichum sp.				1
Cetraria nivalis				4
Cladonia fimbriata				1

b

Tab. 5. (continued)

Community number	
3	527 93 411499 3 344432779999 61666666666441117777772441633323116733425525299888888255525555
37470796658001378261587899678012393601264572123475623415456801928990346012138455678949012342453678	
.....2.....
.....1.....
.....2.....1.....
.....3.....1.....
.....4.....
.....5.....
.....4.....4544.44455432.....
.....1.....31.....122.4...314.....1.5
.....3.....1.....245142.14.342553.....155
.....1.....2113.4321355221.....4453211.2224.....
.....1.....45455132.4314133115354.1.2.1.....43.....13415..34344..33344...11..
.....13.....11.233.....243243.31.....1.13..314.25.....
.....251425.12112.....2.1.....111...1.1.....1.22.....1.2
.....245.....122422.3.21.....32..333.1.....224141.22.13.....1232
.....244.3.....111.2.12..21.112.....232434..2.....11.15..55..11.....
.....1.....2311.553.1.....1.12111151.....43.113.2.3.....
.....154.45.1224..4.1.....43.113.2.3.....1.2.31..12.....1
.....3.....2355.152.....11.....1.1.....12.....42..21..2545
.....2.3.....111.....11.....1.1.....34..2.2.....
.....3.....123.32..111.1.....1.1.....1122.21.11.....33
.....222.4.....111.13.21.....1225555.....
.....1.4.....4233.....11.....111.....1124..44112.....
.....1.1.....4233.....11.....1.2.1.....1225555.....
.....3.1.....1.1.1.....32.122.4..2.....52.5
.....12.3.55.....332123.....131.....
.....53.....111242.2.....1.4..343.....
.....53.....4534.233.....1.1111.1..4..1.1.....
.....54.24.4523..4.1.....1.....21.....
.....451.22233.1.....21.....
.....3.....12.....1121.121.....24..2.....
.....11.....1124..2.....11.....1.....
.....1114243.1.....311422.....11.2.2.....1.2.1.....
.....11.....311422.....123.4334.1.3.1.....
.....34.....11..24.....3.....11..12311..3.1.....
.....11.....12122.4.1.....52.....12311..3.1.....
.....11.1.1.112.3.....1.....21..11.12..1.....
.....1.....1.....
Cetraria cucullata	
Bartramia halleriana	
Cladonia rangiferina	
Pseudoevernia furfuracea	
Dicranum sp.	
Mosses	
Mnium undulatum	
Dicranum pellucidum	
Rhytidium rugosum	
<u>Pinus nigra s.l. communities</u>	
<u>Pinus nigra salzmannii</u>	
Pinus nigra s.str.	
Quercus-Fagetea CL	
Hepatica nobilis	
Amelanchier ovalis	
Deschampsia flexuosa s.str.	
Fragaria vesca	
Buxus sempervirens	
Sorbus aucuparia	
Viola reichenbachiana	
Sorbus aria	
Lonicera xylosteum	
Rosa pendulina	
Corylus avellana	
Crataegus monogyna	
Abies alba	
Viburnum lantana	
Helleborus foetidus	
Fagus sylvatica	
Brachypodium sylvaticum	
Melampyrum pratense s.str.	
Poa nemoralis	
Carex ornithopoda	
Rubus idaeus	
Prenanthes purpurea	
Melampyrum sylvaticum	
Geum sylvaticum	
Quercus petraea	
Hedera helix	
Quercus faginea	
Daphne laureola s.str.	
Ligustrum vulgare	
Thalictrum tuberosum	
Fraxinus excelsior	
Galium rotundifolium	
Prunus mahaleb	
Arenaria montana	
Berberis vulgaris s.str.	
Betula pendula	
Prunus spinosa	
Berberis hispanica	
Primula veris columinae	
Acer granatense	
Genista cinerascens	
Rosa canina	



Tab. 5. (continued)

Community number	
3	Acer opalus
527	Viola riviniana
93	Cytisus sessilifolius
411499	Sorbus domestica
3	Cornus sanguinea
344432779999	Viola wilkommii
6166666666144111777777441633323116733425525299888888258888882552555	Quercus pubescens
37470796658001378261587899678012393601264572123475623415455801928990346012138455678949012342453678	Dryopteris filix-mas
	Rubus saxatilis
	Quercus robur
	Carex digitata
	Sorbus chamaemespilus
	Mycelis muralis
	Paeonia officinalis microcarpa
	Carex alba
	Aquilegia vulgaris
	Genista florida
	Adenocarpus hispanicus
	Lonicera etrusca
	Populus tremula
	Sanicula europaea
	Euphorbia amygdaloides
	Laserpitium latifolium
	Melampyrum pratense ssp. alpestre
	Helleborus viridis occidentalis
	Dryopteris dilatata
	Polygonatum odoratum
	Veronica urticifolia
	Athyrium filix-femina
	Paris quadrifolia
	Luzula sylvatica
	Rhamnus alpinus
	Daphne mezereum
	Stellaria holostea
	Vicia sepium
	Lathyrus montanus
	Festuca heterophylla
	Campanula persicifolia
	Rosa sp.
	Rubus ulmifolius
	Dryopteris carthusiana
	Frangula alnus
	Epilobium montanum
	Cephalanthera rubra
	Geranium sanguineum
	Sorbus torminalis
	Betula pubescens
	Acer monspessulanum
	Hypericum montanum
	Potentilla micrantha
	Origanum vulgare
	Quercus pyrenaica
	Viola alba dehnhardtii
	Rosa pouzini
	Ilex aquifolium

Tab. 5. (continued)

Community number	3	527	93	411499	3	344432779999	6166666661441117777724416332311673342552523988888825888882552555	37470796658001378261587899678012393601264572123475623415455801928990346012138455678949012342455678
<i>Acer campestre</i>							3	1.1
<i>Acer platanoides</i>							1	1
<i>Melica uniflora</i>							1.1	1.1
<i>Prunus avium</i>							13	1
<i>Teucrium scorodonia</i>							21.1	3
<i>Viola odorata</i>							1.2	1.22
<i>Lilium martagon</i>								1
<i>Poa chaixii</i>							23	1
<i>Convallaria majalis</i>							3	11
<i>Clinopodium vulgare</i> s.str.							24	4
<i>Berberis vulgaris</i> serot								333
<i>Rosa pimpinellifolia</i>							24	1
<i>Polygonatum verticillatum</i>								11
<i>Carpinus betulus</i>							24	2
<i>Acer pseudoplatanus</i>							3.5	3
<i>Campanula rapunculoides</i>							1	1
<i>Taxus baccata</i>							1	1
<i>Sorbus mougeotii</i>								1
<i>Geranium sylvaticum</i>								11
<i>Rosa sicula</i>								1
<i>Daphne laureola latifolia</i>								11
<i>Crataegus laciniata</i>								1
<i>Rosa corymbifera</i>								1
<i>Rosa spinosissima</i> ssp. <i>myriacantha</i>							2	21
<i>Viola canina</i>								3
<i>Rhamnus catharticus</i>							3	3
<i>Pyrus communis</i>							41	
<i>Primula veris</i> s.str.							32	
<i>Lathyrus niger</i>							25	
<i>Trifolium medium</i>							52	
<i>Buglossoides purpureoaeerulea</i>							22	
<i>Eupleurum falcatum</i>							11	
<i>Galium sylvaticum</i>							21	
<i>Viola hirta</i>							34	
<i>Potentilla alba</i>							13	
<i>Rubus</i> sp.							1	1
<i>Tilia cordata</i>								22
<i>Sambucus racemosa</i>								12
<i>Galium odoratum</i>								32
<i>Petasites albus</i>								11
<i>Lamiasstrum galeobdolon</i>								11
<i>Neottia nidus-avis</i>								41
<i>Luzula luzuloides</i>								34
<i>Quercus x cerricoides</i>							4	12
<i>Tilia platyphyllos</i>							4	1
<i>Clematis recta</i>								12
<i>Clematis vitalba</i>							3	1
<i>Dictamnus albus</i>								12
<i>Euphorbia dulcis</i>							3	1
<i>Thesium bavarum</i>							3	1
<i>Abies maroccana</i>								1
<i>Aquilegia pyrenaica</i>								1
<i>Cedrus atlantica</i>								1
<i>Viola mirabilis</i>							3	1
<i>Genista cephalantha demnatisensis</i>								1

Tab. 5. (continued)

Community number					
3	527	93	411499	3	344432779999
37470796658001	37826158789967801	239360126457212347	56234415455801928990346012138455678949012342453678	61666666661441117777724416332311673342552529988888258888882552555	
Peucedanum cervaria5.1.
Melittis melissophyllum1.
Fragaria viridis1.
Rosa tomentosa5.
Crataegus sp.5.
Campanula trachelium1.
Pulmonaria angustifolia1.
Rosa arvensis2.
Salix angustifolia1.
Helleborus niger1.
Rubus chamaemorus1.
Primula vulgaris3.
Anemone nemorosa1.
Genista falcata2.
Euonymus verrucosus1.
Paeonia officinalis s.str.1.
Scilla lilio-hyacinthus1.
Cytisus striatus1.
Carex sylvatica1.
Cotoneaster granatensis1.
Cotinus coggygria4.
Luzula forsteri s.str.4.
Cotoneaster nebrodensis2.
Loniceera periclymenum1.
Rosa stylosa4.
Rubus sect. Histrices2.
Cyclamen purpurascens1.
Geum heterocarpum2.
Cephalanthera damasonium1.
Rosa micrantha1.
Coriaria myrtifolia1.
Cephalanthera longifolia2.
Calluno-Ulicetea CL4.
Veronica officinalis1.
Calluna vulgaris1.
Genista pilosa5.
Erica australis2.
Erica multiflora5.
Erica vagans1.
Quercetea ilicis CL2.
Erica arborea2.
Quercus rotundifolia1212432.
Juniperus oxycedrus11.3
Juniperus phoenicea3
Rubia peregrina453534
Cytisus scoparius12122
Bupleurum rigidum1212133
Rhamnus alaternus121254
Quercus coccifera1
Daphne gnidium2
Paeonia broteroi34
Rhamnus myrtifolius1
Loniceera splendida24
Aristolochia pistolochia1
Pistacia terebinthus1

Tab. 5. (continued)

Community number				
	3	527	93	411499
	37470796658001	378261587899678012	39360126457212347	5623415455801928990346012138455678949012342453678
<i>Luzula pilosa</i>323.4.555.....
<i>Maianthemum bifolium</i>42533..2..44.....
<i>Knautia dipsacifolia</i>24.....
<i>Hieracium sylvaticum</i>51.....2.....
<i>Melica nutans</i>31.....1.....
<i>Carex flacca</i>1.5.3.1.....21.....
<i>Phyteuma spicatum</i>1.....1.1.11.....34.....
<i>Festuca indigesta s.str.</i>4.....
<i>Calamagrostis villosa</i>343244.....2.....
<i>Euphorbia cyparissias</i>2321325.....
<i>Prunella grandiflora pyrenaica</i>12..242.....
<i>Leucanthemum vulgare</i>12.....42.....
<i>Catananche caerulea</i>135.4111.....
<i>Brachypodium pinnatum s.str.</i>11.3.....1..2.13.....
<i>Conopodium pyrenaicum</i>11.....
<i>Leontodon crispus bourgaeanus</i>1.2.131.22.....
<i>Agrostis castellana</i>122.....1.311.....
<i>Linaria nivea</i>2.21.....121.....1.....
<i>Festuca indigesta aragonensis</i>1.....11.1.12.....
<i>Carduus carpetanus</i>2.....1.1211.....
<i>Koeleria crassipes</i>2.3.1.....1.1.....2.....
<i>Leucanthemopsis pallida alpina</i>1.....1211.3.....
<i>Coronilla minima</i>1.....43.3.....24.....
<i>Teucrium montanum</i>3412.1.....1.....3.....
<i>Genista hispanica</i>24.....
<i>Galium ludicum</i>4444.....1.2.....
<i>Avenula bromoides</i>4.....
<i>Euphorbia nicaeensis</i>12.122.....
<i>Hellianthemum cinereum</i>1.....142.....
<i>Festuca eskia</i>11.....
<i>Calamagrostis arundinacea</i>31322.....1.....
<i>Lathyrus filiformis</i>2..32.....
<i>Leuzea conifera</i>32.23.....
<i>Biscutella valentina</i>33.....
<i>Thymus bracteatus</i>4.45.....3.33.....
<i>Thymus serpyllum</i>4234.....
Further do occur:				
<i>Gymnadenia odoratissima</i>	91:2,	92:3,	93:4,	94:3,
<i>Campanula cochlearifolia</i>	92:1,	93:5,	94:2,	95:3,
<i>Dactylis glomerata</i>	6:1,	11:1,	78:4,	33:3,
<i>Hippocrepis comosa</i>	90:4,	91:3,	92:4,	93:4,
<i>Hellianthemum nummularium s.str.</i>	27:1,	23:1,	56:1,	57:3,
<i>Geranium robertianum</i>	11:1,	74:1,	71:1,	85:1,
<i>Hypochoeris radicata</i>	3:1,	6:2,	96:1,	98:1,
<i>Hellianthemum croceum</i>	15:3,	68:4,	30:3,	31:3,
<i>Phyteuma orbiculare</i>	90:4,	92:2,	93:1,	94:2,
<i>Scorzonera humilis</i>	80:2,	81:1,	82:2,	84:4,
<i>Brachypodium retusum</i>	61:3,	65:2,	67:3,	69:5,
<i>Rumex acetosella</i>	43:4,	80:2,	81:2,	83:2,
<i>Arenaria armerina</i>	30:3,	33:1,	34:1,	46:2,
<i>Festuca hystrix</i>	65:3,	67:3,	68:1,	19:2,
<i>Hellianthemum apenninum</i>	6:1,	2:1,	36:2,	19:1,
<i>Vincetoxicum hirundinaria s.str.</i>	77:4,	78:1,	16:1,	19:1,
<i>Scabiosa turoleensis</i>	65:4,	15:1,	68:2,	33:2,
<i>Cerastium boissierianum</i>	15:1,	30:3,	33:1,	34:2,
<i>Lathyrus linifolius</i>	41:1,	42:1,	13:1,	14:1,
<i>Brachypodium pinnatum rupestre</i>	12:1,	41:2,	13:1,	14:1,
<i>Avenula mirandana</i>	9:4,	16:3,		
<i>Carduus defloratus</i>	90:14,	91:2,	92:2,	93:5,
<i>Festuca ovina</i>	78:4,	81:5,	82:5,	83:5,
<i>Brachypodium phoenicoides</i>	62:3,	66:2,	64:3,	65:4,
<i>Prunus prostrata</i>	44:2,	45:1,	68:1,	30:2,
<i>Potentilla tabernaemontani</i>	62:2,	66:4,		
<i>Arrhenatherum album</i>	4:2,	6:1,		
<i>Hieracium vahlii myriadenum</i>	3:1,	7:1,	40:1,	
<i>Pinus pinaster</i>	6:4,	96:1,	5:2,	33:1,
<i>Peucedanum oreoselinum</i>	77:3,	78:4,	91:3,	
<i>Anthericum ramosum</i>	77:3,	78:3,	44:2,	45:1,
<i>Agrostis tenuis</i>	31:2,	33:3,	34:2,	Rosmarinus officinalis
<i>Festuca paniculata</i>	23:1,	56:2,	57:2,	58:1,
<i>Sedum forsterianum</i>	11:1,	5:1,		
<i>Jaione montana echinata</i>	6:1,	96:1,	1:1,	5:1,
<i>Echinopsium boissieri</i>	15:2,	68:3,	30:3,	31:3,
<i>Thymus orospedanus</i>	15:3,	68:4,	30:4,	31:2,
<i>Vicia pyrenaica</i>	9:4,	16:3,		

Tab. 5. (continued)

70:5; Lavandula angustifolia pyrenaica 9:2, 16:3, 32:1; Lactuca viminea 96:1, 10:1, 11:3; Thesium alpinum 90:3, 93:2, 94:1; Valeriana tripteris 90:4, 85:4; Luzula multiflora 81:3, 82:5, 83:3; Stachys officinalis 78:3, 16:1, 41:1; Hieracium gr. peleteranum 75:2, 76:3, 74:1; Avena pratensis s.l. 76:1, 73:1, 74:1; Pulsatilla alpina s.str. 13:1, 14:2, 73:2; Conopodium majus s.str. 75:2, 76:1, 72:1; Erysimum grandiflorum 66:3, 65:2, 67:2; Medicago suffruticosa 62:1, 66:3, 65:2; Vicia tenuifolia 61:3, 64:1, 65:2; Vicia incana 12:1, 13:1; Hieracium sp. 26:1, 13:1; Lathyrus pratensis 91:3, 13:1; 43:1, 48:1; Jasione laevis carpatica 8:1, 10:1, 43:3; Jurinea humilis 97:1, 38:1, 47:2; Sanguisorba minor s.str. 32:3, 69:3, 33:2; Phlomis lychnitidis 31:1, 33:2, 34:2; Teucrium webbianum 31:3, 33:1, 34:2; Polygala boissieri 30:3, 29:2, 32:2; Astragalus granatensis 66:4, 68:2, 29:2; Saxifraga chneufolia 28:2, 85:2, 86:4; Globularia cordifolia 26:1, 90:4, 93:1; Centaurea scabiosa 26:1, 93:4, 94:2; Saponaria ocyroides 26:1, 93:5, 45:1; Phyteuma hemisphaericum 54:2, 23:1, 57:1; Ononis pusilla 26:1, 19:1, 33:1; Silene saxifraga 98:1, 18:1, 19:1; Poa ligulata 45:1, 18:1, 19:4; Helianthemum canum s.str. 67:3, 18:1, 19:4; Fumana ericoides 18:4, 19:3, 69:2; Acinus alpinus meridionalis 11:1, 30:2, 31:1; Teucrium pyrenaicum s.str. 91:1, 23:1, 58:2; Clinopodium vulgare arundanum 61:1, 11:2, 98:1; Agrostis delicatula 61:1, 96:1, 10:1; Gentiana lutea 3:1, 37:1, 10:1; Thymus praecox 9:2, 2:4, 5:1; Cistus laurifolius 36:3, 1:1, 5:4; Helictotrichon filifolium 15:3, 68:5, 30:3; Lonchocera pyrenaica 14:1, 23:1, 56:1; Leontodon pyrenaicus 14:1, 54:3, 55:5; Echinospartum horridum 12:1, 41:1, 25:3; Vicia incana 12:1, 13:1; Hieracium sp. 26:1, 13:1; Hieracium sp. 26:1, 13:1; Sedum brevifolium 4:1, 43:1; Carduus carlinifolius 13:1, 14:2; Fumana paradoxa 15:1, 31:1; Urtica dioica 6:1, 98:1; Galium rivulare 8:1, 40:1; Digitalis Orobanche rapum-genistae 4:1, 43:1; Festuca elegans 96:4, 5:1; Centaurea alba 96:1, 5:1; Lathyrus vernus 77:4, 16:1; Epipactis helleborine 17:1, 25:1; Purpurea s.str. 10:1, 43:2; Hieracium castellatum 40:1, 10:1; Holcus mollis 11:2, 81:2; Lathyrus vernus 77:4, 16:1; Sanguisorba rupicola 18:3, 19:2; Globularia repens 18:4, 19:4; Jasonia glutinosa 18:4, 19:2; Rhannus pulmillus 18:3, 23:1; Draba dedeana 18:4, 19:3; Sanguisorba rupicola 18:3, 19:2; Sedum dasycarpum 18:1, 19:1; Anthyllis vulneraria 18:1, 19:4; Hypericum richeri bursari 20:1, 52:1; Salix pyrenaica 20:1, 22:1; Polystichum lonchitis 72:1, 20:1; sp. 19:1, 28:1; Polystichum aculeatum 72:1, 20:1; Thymelaea nivalis 25:1, 23:1; Senecio adonidifolius 23:1, 56:1; Sideritis hyssopifolia 23:1; Geum montanum 20:1, 28:1; Silene rupestris 23:1, 56:1; Thymelaea nivalis 25:1, 23:1; Saxifraga paniculata 23:1, 56:1; Galium verum 26:1, 92:5; Asperula 58:2; Helictotrichon sedenense 23:1, 56:1; Hieracium cerinthoides 23:1, 56:1; Saxifraga paniculata 23:1, 56:1; Galium verum 26:1, 92:5; Asperula aristata 27:1, 26:1; Festuca rubra 28:1, 57:2; Teucrium leonis 30:3, 31:1; Viola suavis 30:1, 32:2; Scabiosa andryvalifolia 30:3, 31:1; Alyssum sarpyllifolium 30:2, 34:1; Genista pseudopilososa 30:1, 34:1; Piptatherum paradoxum 69:2, 33:1; Teucrium polium 33:2, 46:1; Cistus albidus 69:3, 33:1; Silene mallifera 45:1, 33:1; Crepis alba 45:1, 33:1; Galium frutescens 33:1, 34:1; Carlina corymbosa 34:3, 46:1; Santolina chamaecyparissus 65:3, 34:1; Bupleurum frutescens 69:5, 34:1; Avena lodunensis 35:1, 36:4; Dianthus laricifolius 38:1, 47:2; Sedum album micranthum 38:1, 47:2; Lactuca chondrillaefolia 8:1, 43:1; Gnaphallium sylvaticum 11:1, 43:1; Saponaria glutinosa 44:3, 45:2; Thymus zapateri 44:3, 45:3; Myosotis sylvatica 44:3, 45:2; Galium idubedae 44:2, 45:1; Arabis turrita 44:3, 45:1; Siler nestleri 44:1, 45:1; Epipactis microphylla 44:1, 45:1; Bupleurum ranunculoides 45:2; Ononis genista 44:2, 45:1; Allium sphaerocephalum 44:1, 45:1; Geum urbanum 44:1, 45:1; Epipactis microphylla 44:1, 45:1; Juncus trifidus 54:5, 44:1, 45:1; Elymus caninus 16:1, 45:1; Gentiana bursi 72:1, 52:2; Primula integrifolia 54:2, 55:2; Carex curvula 54:3, 55:3; Juncus trifidus 54:5, 55:2; Carex sp. 56:1, 57:1; Sempervivum montanum 23:1, 56:1; Meum athamanticum 56:1, 57:1; Globularia nudicaulis 56:1, 58:4; Genista heterochrous 61:4, pauciflora 75:3, 76:1; Carex pilulifera 80:3, 82:3; Danthonia decumbens 80:2, 82:5; Lycopodium clavatum 81:3, 84:2; Aster bellidiflorum 85:3, 86:1; Euphrasia salisburgensis 93:3, 94:2; Biscutella laevigata 93:4, 94:3; Leucanthemum maximum 93:4, 94:3; Mucedonia sedoides 98:1; Teucrium pyrenaicum sp. quarensis 12:2; Viola cornuta 98:1; Potentilla neumanniana 12:1; Picris hieracioides 98:1; Pulmonaria longifolia 12:1; Artemisia glutinosa 98:1; Hippoeris radicata 98:1; Hieracium gr. praecox 13:1; Viscum album ssp. austriacum 13:1; Holcus setigellus 98:1; Dianthus hyssopifolius 14:2; Juncus conglomeratus 98:1; Hieracium lactucella 14:1; Scutellaria minor 98:1; Viola rupestris 14:1; Aconitum napellus castellanum 98:1; Helianthemum canum pourtaii 15:1; Oxalis corniculata 98:1; Centaurea jaenensis 15:1; Senecio sp. 98:1; Lapsana communis 98:1; Pimpinella saxifraga 16:1; Juncus inflexus 2:1; Conopodium filifolium 97:1; Ranunculus ollisiponensis alpinus 2:1; Thlaspi stenopterum 2:1; Senecio jacobaea 97:1; Festuca costei 2:1; Avenula sulcata albinervis 97:1; Narcissus rupicola 3:2; Jasione crispa centralis 3:1; Taraxacum sroeterianum 3:1; Thymus zygis 6:2; Geranium purpureum 6:1; Festuca ampla 97:2; Andryala integrifolia 6:1; Dorycnium pentaphyllum 9:1; Arenaria queruloidea 96:1; Cirsium cf. tuberosum 9:1; Sedum hirsutum 96:1; Thymelaea pubescens 9:1; Spermogula morisonii 96:1; Carlina vulgaris 9:1; Thymus mastichina 96:1; Festuca summlusitana 96:4; Epilobium lanceolatum 11:1; Corynephorus canescens 96:2; Festuca rothmaleri 11:1; Festuca agr. rubra 11:1; Carex muricata lamprocarpa 11:2; Laserpitium krapffii gaudinii 93:5; Hieracium bupleuroides 93:3; Galeopsis ladaranum 11:1; Leontodon hispidus 93:4; Pentaglottis sempervirens 11:1; Trifolium montanum 92:4; Scabiosa columbaria 92:5; Mercurialis perennis 16:1; Gymnadenia conopsea 92:5; Dianthus monspessulanus 17:1; Salix caprea 92:3; Carex caryophylla 17:1; Molinia caerulea arundinacea 91:1; Vincetoxicum hiruandina intermedium 17:1; Prunella grandiflora s.str. 91:3; Cardamine pratensis 17:1; Potentilla caulescens 18:4; Vicia cracca gerardi 91:2; Erinus alpinus 18:2; Antirrhinum pulverulentum 18:2; Campanula cespitosa 90:2; Asplenium fontanum 18:2; Asplenium ruta-muraria 18:2; Galium austriacum 90:4; Armeria trachyphylla 19:2; Scabiosa lucida 90:4; Sedum album s.str. 19:2; Carduus crassifolius ssp.glaucus 90:5; Asphodelus albus 19:1; Ajuga reptans 85:2; Armeria obtusiflora 19:1; Dactylorhiza maculata 85:2; Salvia lavandulifolia 19:1; Chaerophyllum hirsutum 85:1; Dipcadi serotinum 19:2; Adenostyles glabra 86:1; Armeria aggregata erinacea 19:1; Satureja obovata 19:1; Molinia caerulea s.str. 84:4; Dianthus brachyanthus 19:1; Luzula nutans 20:1; Soldanella alpina 20:1; Genista tinctoria 82:4; Viscum album s.str. 81:3; Trifolium alpinum 20:1; Veronica chamaedrys 78:1; Jasione laevis s.str. 20:1; Serratula tinctoria 78:4; Festuca violacea 20:1; Ranunculus polyanthemophyllus 78:2; Carex sempervirens 20:1; Salix reticulata 22:1; Trifolium alpestre 78:4; Thymelaea dioica 23:1; Gentiana occidentalis 23:1; Inula conyza 77:2; Potentilla alchimilloides 23:1; Arabis brassica 77:1; Helianthemum oleandicum italicum 23:1; Coronilla coronata 77:3; Valeriana globularifolia 23:1; Trifolium rubens 77:4; Potentilla montana 24:1; Galium saxatile 24:1; Eryngium bourgati 25:1; Euphorbia angulata 73:1; Seseli nanum 25:1; Athyrium distentifolium 72:1; Viscum album ssp. microphyllum 26:2; Pulmonaria affinis 72:1; Odontites cf. viscosa 26:2; Plantanthea bifolia 26:1; Astragalus monspessulanus 26:1; Viola alpestris 26:1; Sedum sediforme 69:3; Stipa calamagrostis 27:1; Stipa parviflora 69:3; Silene otites 27:1; Cistus salvifolius 69:3; Pinus halepensis 69:3; Phleum alpinum 28:1; Ulex parviflorus 28:1; Agrostis schraderiana 28:1; Alchemilla vulgaris 28:1; Veratrum album 28:1; Thymelaea tinctoria 69:3; Helianthemum nummularium pyrenaicum 25:1; Satureja innota 69:5; Odontites granadensis 29:1; Melica ciliata 67:3; Prunus ramburii 29:2; Limodorum abortivum 66:2; Vella spinosa 29:3; Satureja intricata 29:1; Astragalus danicus 66:3; Daphne oleoides 29:2; Filipendula vulgaris 66:3; Galium maritimum 62:2; Daphne hispanica 30:1; Euphorbia characias 61:2; Thymus claudestinus 30:4; Thymus gadorensis 30:1; Festuca

Tab. 5. (continued)

nevadensis 30:1; Thalictrum foetidum valentinum 30:1; Festuca airoides 54:3; Satureja gracilis 30:3; Minuartia sedoides 54:3; Erysimum favargerii 30:1; Luzula lutea 54:4; Vincetoxicum nigrum 31:4; Jasione crispa s.str. 54:2; Cerastium gibraltarium 31:3; Gentiana acaulis 54:2; Globularia spinosa 31:2; Saxifraga spathularis 53:2; Teucrium similiatum 31:2; Senecio pyrenaicus s.str. 51:1; Santolina rosmarinifolia 46:1; Silene vulgaris 32:4; Crambe filiformis 46:1; Arenaria valentina 32:2; Teucrium chamaedrys gracile 46:1; Carlina acanthifolia 32:2; Origanum grosii 46:1; Anarrhinum laxiflorum 33:2; Psilostemon riphaeum 33:1; Salvia phlomisoides 33:1; Agropyron panormitanum 46:1; Linum suffruticosum s.str. 33:1; Erysimum medio-hispanicum 33:1; Nepeta tuberosa reticulata 46:1; Calamintha sylvatica s.str. 33:1; Sanguisorba minor muricata 46:2; Echium flavum 33:1; Scabiosa tomentosa 46:1; Thlaspi perfoliatum 33:2; Erica terminalis 46:1; Teucrium capitatum 34:2; Stipa tenacissima 46:2; Eryngium campestre 34:3; Asperula hirsuta 46:1; Argemone zanonii 34:2; Thymelaea tartonraira 46:2; Hippocrepis bourgaei 34:2; Ulex baeticus 46:2; Thymelaea sanamunda 34:1; Scorzonera pygmaea 46:1; Helianthemum asperum 34:1; Linum suffruticosum salsoloides 45:1; Astragalus incanus 34:1; Rhannus infectoria 45:1; Leucanthemopsis pallida s.str. 36:2; Carduncellus monspeliensis 45:1; Saxifraga wilkommiana 40:1; Aconitum lycoctonum 45:1; Arrhenatherum elatius s.str. 40:1; Teucrium chamaedrys pinnatifidum 45:1; Thalictrum minus 45:1; Hieracium argyrococomum 43:2; Lithospermum officinale 45:1; Cerastium ramosissimum 43:4; Asperula laevigata 44:2; Pseudarrhenatherum longifolium 43:2; Conopodium majus burgaei 43:1; Leontodon carpetanus 43:1; Ranunculus ollissiponensis s.str. 43:1; Agrostis alpina 43:1.

Table 6. References of the relevés of Table 5.

- 1-5. Relevés of the authors. 'Junipero nanae-Cytisetum oromediterranei geographical race with *Senecio carpetanus pinetosum sylvestris*'. Guadarrama Mountains, Sistema Central, Spain
- 6, 96, 97. Relevés of the authors. 'Junipero nanae-Cytisetum oromediterranei geographical race with *Echinopartum barnadesii* pinetosum sylvestris and altitudinal form with *Pinus nigra* subsp. *salzmannii*'. Gredos, Sistema Central, Spain
- 7, 8. Rivas-Martínez et al. (1987). 'Junipero nanae-Cytisetum oromediterranei'. Guadarrama Mountains, Sistema Central, Spain
9. Gamisans et al. (1991). 'Lonicero xylostei-Pinetum salzmannii'. Central Pyrenees, Spain
10. Fernández-González (1991). 'Senecioni carpetani-Cytisetum oromediterranei'. Guadarrama Mountains, Sistema Central, Spain
11. Fernández-González (1991). 'Luzulo forsteri-Quercetum pyrenaicae'. Guadarrama Mountains, Sistema Central, Spain
12. Ninot (1996). 'Buxo-Quercetum pubescentis'. Central Pyrenees, Spain
13. Ninot (1996). 'Hylocomio-Pinetum catalaunicae'. Central Pyrenees, Spain
14. Ninot (1996). 'Pulsatillo-Pinetum uncinatae'. Central Pyrenees, Spain
15. Valle et al. (1988). 'Junipero phoeniceae-Pinetum salzmannii'. Eastern Andalusia, Spain
16. Gamisans and Gruber (1988). 'Lonicero xylostei-Pinetum salzmannii'. Central and eastern Pyrenees, Spain
- 17, 73-76. Vigo (1979). 'Hylocomio-Pinetum catalaunicae'. Eastern Pyrenees, Spain
18. Regato and Escudero (1989). 'Vertical rupicolous community with *Pinus nigra* Arn.'. Southern Sistema Ibérico, Spain
19. Regato and Escudero (1989). 'Community on horizontal rocky plates with *Pinus nigra* Arn.'. Southern Sistema Ibérico, Spain
- 20, 51, 52. Rivas-Martínez et al. (1991). 'Rhododendro ferruginei-Pinetum uncinatae'. Western Pyrenees, Spain, France
- 21, 53. Rivas-Martínez et al. (1991). 'Rhododendro ferruginei-Abietetum albae'. Western Pyrenees, Spain, France
22. Rivas-Martínez et al. (1991). 'Salici pyrenaicae-Arctostaphyletum alpinae'. Western Pyrenees, Spain, France
- 23, 56-58. Rivas-Martínez et al. (1991). 'Arctostaphylo uvae-ursi-Pinetum uncinatae'. Western Pyrenees, Spain
- 24, 59. Rivas-Martínez et al. (1991). 'Veronico officinalis-Pinetum sylvestris'. Western Pyrenees, Navarra, Spain
25. Rivas-Martínez et al. (1991). 'Echinosparto horridi-Pinetum sylvestris'. Western Pyrenees, Spain
26. Rivas-Martínez and Géhu (1978). 'Ononido rotundifoliae-Pinetum sylvestris'. Valais, Switzerland
27. Rivas-Martínez and Géhu (1978). 'Cotino-Juniperetum sabiniae'. Valais, Switzerland
28. Rivas-Martínez and Géhu (1978). 'Rhododendro-Vaccinietum'. Valais, Switzerland
29. Losa Quintana et al. (1986). 'Daphno oleoidi-Pinetum sylvestris'. Sierra Nevada, Spain
30. López Vélez (1996). 'Daphno hispanicae-Pinetum sylvestris'. Sistema Ibérico, Albacete, Spain
31. López Vélez (1996). 'Junipero phoeniceae-Pinetum salzmannii'. Sistema Ibérico, Albacete, Spain
- 32-34. Herranz Sanz and Gómez Campo (1986). 'Daphno latifoliae-Aceretum granatensis'. Sistema Ibérico, Albacete, Spain
35. Rivas-Martínez and Cantó (1987). 'Adenocarpo hispanici-Genistetum floridae'. Guadarrama Mountains, Sistema Central, Spain
36. Rivas-Martínez and Cantó (1987). 'Erico arborea-Arctostaphyletum crassifoliae'. Guadarrama Mountains, Sistema Central, Spain
- 37-39, 48-50. Rivas-Martínez et al. (1987). 'Junipero nanae-Cytisetum oromediterranei'. Guadarrama Mountains, Sistema Central, Spain
40. Fernández-González (1991). 'Senecioni carpetani-Cytisetum oromediterranei'. Guadarrama Mountains, Sistema Central, Spain
- 41, 42. Vigo (1979). 'Buxo-Quercetum pubescentis'. Eastern Pyrenees, Spain
43. Rivas-Martínez (1963). 'Junipereto-Sarothamnetum purgantis'. Guadarrama Mountains, Sistema Central, Spain
- 44, 45. Rivas Goday and Carbonell (1961). 'Sabineto-Pinetum sylvestris'. Gudar and Javalambre mountains, Sistema Ibérico, Spain

Table 6. (continued)

46. Benabid (1984). 'Com. *Pinus pinaster* var. *maghrebiana* with *P. chusiana* var. *mauritanica*'. Rif, Morocco
47. Rivas-Martínez and Cantó (1987). 'Junipero nanae-Cytisetum oromediterranei'. Guadarrama Mountains, Sistema Central, Spain
54. Rivas-Martínez et al. (1991). 'Luzulo luteae-Loiseleurietum procumbentis'. Western Pyrenees, Spain
55. Rivas-Martínez et al. (1991). 'Carici curvulae-Empetrum hermaphroditum'. Western Pyrenees, Spain, France
- 60, 63. Regato et al. (1995). 'Lonicero xylostei-Pinetum salzmannii'. Central Pyrenees, Spain
- 61, 62. Regato et al. (1995). 'Lonicero xylostei-Pinetum salzmannii'. Sistema Ibérico, Spain
64. Regato et al. (1995). 'Thalictro tuberosi-Pinetum salzmannii'. Central Pyrenees, Spain
- 65, 66. Regato et al. (1995). 'Thalictro tuberosi-Pinetum salzmannii'. Sistema Ibérico, Spain
67. Regato et al. (1995). 'Festuco gautieri-Pinetum salzmannii'. Sistema Ibérico, Spain
68. Regato et al. (1995). 'Junipero phoeniceae-Pinetum salzmannii'. Sistema Betico, Spain
69. Regato et al. (1995). 'Hedero-Genistetum patentis'. Sistema Ibérico, Spain
70. Regato et al. (1995). 'Bupleuro-Quercetum rotundifoliae'. Eastern Pyrenees, Spain
71. Vigo (1979). 'Helleboro-Fagetum'. Eastern Pyrenees, Spain
72. Vigo (1979). 'Saxifrago-Rhododendretum'. Eastern Pyrenees, Spain
77. Braun-Blanquet (1932). 'Lithospermo-Quercetum'. Central Europa
78. Oberdorfer (1957). 'Potentillo-Quercetum'. Central Europa
79. Matuszkiewicz (1956a). 'Quercu-Potentilletum albae'. Poland
80. Passarge (1957). 'Myrtillo-Pinetum'. Germany
81. Preising (1943). 'Dicrano-Pinetum eupteridetosum'. Poland
82. Preising (1943). 'Dicrano-Pinetum typicum'. Poland
83. Matuszkiewicz (1956b). 'Pino-Quercetum berberidetosum'. Poland
84. Matuszkiewicz (1954). 'Pino-Vaccinietum myrtilli'. Poland
85. Kuoch (1954). 'Abietetum festucetosum'. Alpes, Switzerland
86. Kuoch (1954). 'Abietetum melampyretosum Carex-Variante'. Alpes, Switzerland
87. Braun-Blanquet et al. (1954). 'Piceetum montanum galietosum'. Alpes, Switzerland
88. Braun-Blanquet et al. (1954). 'Piceetum subalpinum myrtilletosum Linnaea-Variante'. Alpes, Switzerland
89. Braun-Blanquet et al. (1954). 'Piceetum subalpinum vaccinietosum, Peltigera-Hylocomium-Variante'. Alpes, Switzerland
90. Knapp (1944), in Ellenberg (1996). 'Chamaebuxo-Pinetum nigrae'. Alpes, Switzerland
91. Braun-Blanquet et al. (1954). 'Erico-Pinetum'. Alpes, Switzerland
92. Etter (1947). 'Molinio-Pinetum'. Alpes, Switzerland
93. Braun-Blanquet et al. (1954). 'Carici-Pinetum engadinensis'. Alpes, Switzerland
94. Braun-Blanquet et al. (1954). 'Erico-Mugetum'. Alpes, Switzerland
95. Braun-Blanquet et al. (1954). 'Rhododendro hirsuti-Mugetum'. Alpes, Switzerland
98. Relevés of the authors. 'Junipero nanae-Cytisetum oromediterranei geographical race with *Senecio carpetanus* pinetosum sylvestris'. Ayllón Mountains, Sistema Central, Spain

Oak groves of Quercus pyrenaica. Pinus sylvestris Derived Community (DC)

The potential area of *Q. pyrenaica* in the Sistema Central extends below 1500 m and it has been used for a long time to favour *Pinus sylvestris* crops. There are several documents from the age of Philip the Second and even earlier (Bauer 1991, Mancebo et al. 1993), which speak about *P. sylvestris* used as construction wood, for heating, and to make glass; in the Gredos Mountains some *P. nigra* subsp. *salzmannii*, stripped to obtain resin, can be found.

The oak groves of *Q. pyrenaica* of the Sistema Central (Carpetano-Iberico-Leonesa phytogeographical province), can be grouped in 3 associations:

1. Luzulo forsteri-Quercetum pyrenaicae Rivas-Martínez 1963. Guadarrama Mountains and La Alcarria, siliceous, Supramediterranean, subhumid-humid.
2. Genisto falcatae-Quercetum pyrenaicae Rivas-Martínez in Penas & T. E. Díaz 1984. Plains of Salamanca, Zamora and Orense, Gredos Mountains, siliceous, Meso-Supramediterranean, subhumid-humid.
3. Festuco heterophyllae-Quercetum pyrenaicae Br.-Bl. 1967. Sistema Ibérico in the province of Soria and the Ayllón Mountains, siliceous, Supramediterranean, humid-hyperhumid.

Luzulo-Quercetum pyrenaicae and Genisto-Quercetum pyrenaicae are enriched with *P. nigra* subsp. *salzmannii* in the more thermic localities of the eastern Gredos Mountains and western Guadarrama Mountains. Thus, relic forms of oak groves between 1100 and 1300 m can be observed (Fig. 5). This is its most frequent altitudinal distribution in the Sistema Central. *P. nigra* subsp. *salzmannii* is a western Mediterranean tree that in the Pyrenees can be included in the Quercetea ilicis, in the Sistema Ibérico in the Querco-Fagetea, in the Sistema Central in Querco-Fagetea and Cytision oromediterranei, and in the Sistema Bético in the Cytision oromediterranei (Regato et al. 1995).

In Table 3 we can observe that the natural regeneration of natural pine groves begins with *Galium rotundifolium* and *Pteridium aquilinum*, which are indicators of humidity and depth of soil. If the forest has burnt, the most humid areas are enriched with *Arctostaphylos uva-ursi*. In both cases the crops of *Pinus sylvestris* change the attributes of the soil and a distribution of the characteristic plants of Querco-Fagetea, which means that *P. sylvestris* is favoured and becomes potential as opposed to *Q. pyrenaica*. On the other hand, *A. uva-ursi* contributes to the acidity of the soil, and so *Q. pyrenaica* cannot grow, but *P. sylvestris* can. In some places of the Sistema Central altered by man, *P. sylvestris* has become potential in the dominian of Querco-Fagetea, establishing a derived community (DC), according to Kopecný et al. (1995).

Conclusions

There are two different types of *Pinus sylvestris* communities in the Sistema Central:

Natural Pine groves: with boreal and prealpine origin.

* Junipero nanae-Cytisetum oromediterranei pinetosum sylvestris, sometimes with *Pinus nigra* subsp. *salzmannii*.

Crops of Pine groves: Since the Middle Ages.

* *Pinus sylvestris* DC, in previous potential places of *Quercus pyrenaica*.

Pinus nigra subsp. *salzmannii* is more frequent in the oak grove belt (Luzulo forsteri-Quercetum pyrenaicae and Genisto falcatae-Quercetum pyrenaicae), forming relic communities.

To sum up, we give the syntaxonomic scheme of the communities studied:

- CL. VACCINIO-PICEETEA Br.-Bl. in Br.-Bl., Sissingh & Vlieger 1939
 O. Pinetalia sylvestris Oberdorfer 1956
 AL. Cytision oromediterranei R. Tx. in R. Tx. & Oberdorfer 1958 corr. Rivas-Martínez 1987 [Cytision europaei pro nom. mut., incl. Pino-Juniperetalia Rivas-Martínez 1964]. Siliceous Iberian Oromediterranean associations.
 SAL. Cytisenion oromediterranei
 AS. Junipero nanae-Cytisetum oromediterranei (Rivas Goday 1955) Rivas-Martínez 1963 <Sistema Central>
 * Geographical race with *Senecio carpetanus* <Guadarrama Mountains>
 * Geographical race with *Echinopartum barnadesii* <Gredos Mountains>
 * pinetosum sylvestris Rivas-Martínez 1963
 * Variant with *Linaria nivea*
 * Variant with *Koeleria crassipes* and *Corynephorus canescens*
 * Relic form with *Vaccinium myrtillus*
 * Altitudinal form with *Pinus nigra* subsp. *salzmannii*
- CL. QUERCO-FAGETEA Br.-Bl. & Vlieger 1937
 O. Quercetalia roboris R. Tx. 1931
 AL. Quercion robori-pyrenaicae (Br.-Bl., P. Silva & Rozeira 1956) Rivas-Martínez 1975
 SAL. Quercenion pyrenaicae Rivas-Martínez 1975. Iberian Mediterranean oak groves.
 AS. Luzulo forsteri-Quercetum pyrenaicae Rivas-Martínez 1963 <Carpetano-Iberico-Leonesa province, Alcarria>
 * Relic form with *Pinus nigra* subsp. *salzmannii* <Western Guadarrama Mountain>
 * *Pinus sylvestris* DC
- AS. Genisto falcatae-Quercetum pyrenaicae Rivas-Martínez in Penas & T. E. Díaz 1984 <Salamanca, Orense, Zamora, Gredos Mountains>
 * Relic form with *Pinus nigra* subsp. *salzmannii* <Eastern Gredos Mountains, Avila>
 * *Pinus sylvestris* DC
- AS. Festuco heterophyllae-Quercetum pyrenaicae Br.-Bl. 1967 <Sistema Ibérico, Soria, Ayllón Mountains>
 * *Pinus sylvestris* DC

Floristic appendix

The nomenclature and authorship of the taxa in the text and in the Tables follow the Catalogue des plantes de Maroc (Jahandiez and Maire 1931–1934), Flora Europaea (Tutin et al. 1964–1980), Med-Cheklist (Greuter et al. 1984–1989) and Flora iberica (Castroviejo et al. 1986–1997); for the bryophytes and lichens we have followed the Flore des bryophytes (Augier 1966) and Les lichens (Ozenda and Clauzade 1970).

There are some subspecies taxa which can be maintained with difficulty, though the greater part of the taxa of the communities with *Pinus sylvestris* in the Iberian Peninsula are well differentiated. This is the case of the subspecies of *Juniperus communis* [subsp. *alpina* (Suter) Celak, subsp. *hemisphaerica* (K. Presl) Nyman, subsp. *nana* Syme, subsp. *sibirica* Burgsd.], which have also been considered by other authors to describe new syntaxa. The same occurs with the varieties of *P. sylvestris* (var. *nevadensis* Christ, var. *olivicola* Vayr., var. *iberica* Svob., var. *pyrenaica* Svob., var. *catalaunica* Gaussen), which are not well differentiated (Amaral Franco 1986). The studies with enzymatic markers made with *Pinus nigra* must be considered. A large part of the Iberian associations is based on the Iberian-North-african distribution of *P. nigra* subsp. *salzmannii* (Blanco et al. 1996), but Aguinagalde et al. (1997) clearly establish the presence of *P. nigra* subsp. *nigra* in Navarra (Spain).

Zusammenfassung

Die Zusammensetzung der Wälder des spanischen Zentralgebirges, in denen *Pinus sylvestris* und *P. nigra* subsp. *salzmannii* vorkommen, wird unter phytosoziologischen Gesichtspunkten analysiert und dabei mit 683 weiteren europäischen Koniferengemeinschaften verglichen. Durch Anwendung der Kriterien von Kopecký & Hejný (1974), Foucault (1981), Dierschke (1993) und Kopecký et al. (1995) auf die nach Braun-Blanquet (1964) aufgenommenen Inventare werden im Innern der Iberischen Halbinsel die Klasse Vaccinio-Piceeta sowie die Ordnung Pinetalia sylvestris identifiziert. Die Vergesellschaftungen mit *Echinopartum barnadesii* und *Senecio carpetanus* werden als geographische Rassen interpretiert, und weitere Aspekte von *Junipero nanae* – *Cytisetum oromediterranei* (Subassoziationen, Varianten, Reliktformen) werden kommentiert. Vorkommen von *Pinus nigra* subsp. *salzmannii* in *Quercus pyrenaica*-Wäldern sehen die Autoren als wärmeangepaßte Reliktformen, in Gemeinschaft mit *P. sylvestris* als höheliebende Formen an.

We wish to express our gratitude to Linda Hamalainen for her linguistic help. This study has been supported by the project USP 4/97 of the San Pablo-CEU University.

References

- Aguinagalde I., Llorente F. and Benito C. 1997. Relationships among five populations of European black pine (*Pinus nigra* Arn.) using morphometric and isozyme markers. *Silvae Genetica* 46 (1): 1–5.
- Amaral Franco J. 1986. *Pinus* L. In Castroviejo et al. (ed.): *Flora iberica*. Vol. I. C. S. I. C., Madrid.
- Bauer E. 1991. *Los montes de España en la Historia*. Ministerio de Agricultura, Pesca y Alimentación, Madrid.
- Bauerochse A. and Katenhusen O. 1997. Holozäne Landschaftsentwicklung und aktuelle Vegetation im Fimbertal (Val Fenga, Tirol/Graubünden). *Phytocoenologia* 27 (3): 353–453.
- Benabid A. 1984. Étude phytoécologique des peuplements forestiers et preforestiers du Rif centro-occidental (Maroc). *Trav. Inst. Sci. Chérifien, Sér. Bot.* 34: 1–64.
- Blanco Castro E., Casado González M. A., Costa Tenorio M., Escribano Bombín R., García Antón M., Génova Fuster M., Gómez Manzaneque A. G., Gómez Manzaneque F., Moreno Saiz J. C., Morla Juaristi C., Regato Pajares P. and Sainz Ollero H. 1996. *Los bosques ibéricos. Una interpretación geobotánica*. Planeta, Barcelona.
- Bolòs O., Vigo J., Masalles R. M. and Ninot J. M. 1993. *Flora manual dels països Catalans*. Pòrtic, Barcelona.
- Braun-Blanquet J. 1932. Zur Kenntnis nordschweizerischer Waldgesellschaften. *Beih. Botan. Cbl.* 49: 7–42.
- Braun-Blanquet J., Roussine N. and Nègre R. 1952. *Les groupements végétaux de la France méditerranéenne*. CNRS, Paris.
- Braun-Blanquet J., Pallmann H. and Bach R. 1954. Pflanzensoziologische und bodenkundliche Untersuchungen im schweizerischen Nationalpark und seinen Nachbargebieten. II. Vegetation und Böden der Wald- und Zwergstrauchgesellschaften (Vaccinio-Piceetalia). *Ergebn. Wiss. Unters. Schweiz. Nationalpark N.F.* 4: 1–200.
- Braun-Blanquet J. 1964. *Pflanzensoziologie. Grundzüge der Vegetationskunde*. Springer Verlag, Wien.
- Capel Molina J. J. 1981. *Los climas de España*. Oikos-Tau, Barcelona.
- Ceballos L. 1966. *Mapa forestal de España*, 1:400.000. Ministerio de Agricultura, Madrid.
- Costa Tenorio M., Morla Juaristi C. and Sainz Ollero H. 1988. Consideraciones acerca de la evolución de paisaje vegetal de la Península Ibérica en el Cuaternario reciente. *Actes Simp. Intern. Bot. Pius Font i Quer* 2: 427–438.
- Costa Tenorio M., García Antón M., Morla Juaristi C. and Sainz Ollero H. 1990. La evolución de los bosques de la Península Ibérica: Una interpretación basada en datos paleobiogeográficos. *Ecología (Fuera de serie)* 1: 31–58.

- Cristina Peñalba M. 1994. The history of the Holocene vegetation in northern Spain from pollen analysis. *J. Ecology* 82: 815–832.
- Cristofolini G. 1997. The biodiversity of the Leguminosae-Genisteae and its genesis. *Lagascalia* 19 (1–2): 121–128.
- Dahlgren R. & Lassen P. 1972. Studies in the flora of Northern Morocco, I. Some poor fen communities and notes on a number of northern and atlantic plant species. *Bot. Notiser* 125: 439–463.
- Deil U. and Galán de Mera A. 1998. À la connaissance des groupements rupicoles calcaires du Maroc. *Bull. Inst. Bot. Rabat* (in press).
- Dierschke H. 1993. *Grundlagen und Methoden der Pflanzensoziologie*. Ulmer, Stuttgart.
- Elena Rosselló R. and Sánchez Palomares O. 1991. Los pinares españoles de *Pinus nigra* Arn.: Síntesis ecológica. *Monografías INIA* 81: 5–110.
- Ellenberg H. 1996. *Vegetation Mitteleuropas mit den Alpen*. 5. Auflage. Verlag Eugen Ulmer, Stuttgart.
- Esteve Chueca F. 1973–74. Especies y comunidades vegetales de la Sierra Nevada caliza. *Bol. Soc. Brot.* 47: 179–187.
- Etter H. 1947. Über die Waldvegetation am Südostrand des schweizerischen Mittellandes. *Ebenda* 25: 141–210.
- Fernández-González F. 1991. La vegetación del valle del Paular (Sierra de Guadarrama, Madrid), I. *Lazaroa* 12: 153–272.
- Folch i Guillèn R. 1986. *La vegetació dels països Catalans*. Ketres, Barcelona.
- Font Quer P. 1954. La vegetación. In Terán M. (ed.): *Geografía de España y Portugal*. Montaner y Simón, Barcelona.
- Foucault B. 1981. Réflexions sur l'appauvrissement des syntaxons aux limites chorologiques des unités phytosociologiques supérieures et quelques-unes de leurs conséquences. *Lazaroa* 3: 75–100.
- Galán de Mera A., Hagen de la Cerda M. A. and Vicente Orellana J. A. 1995. El índice de humedad total: Un nuevo índice climático para la delimitación de la vegetación en América Latina. *Arnaldoa* 3 (2): 87–98.
- Gamisans J. and Gruber M. 1988. Els boscos de pinassa (*Pinus nigra* subsp. *salzmannii*) als Pirineus catalans i est-aragonesos: Estudi fitosociològic. *Monogr. Inst. Piren. Ecol.* 4: 543–552.
- Gamisans J., Gruber M. and Regato P. 1991. Les forêts de *Pinus nigra* subsp. *salzmannii* des Pyrénées aragonaises. *Lazaroa* 12: 147–151.
- Gausson H., Debrach J. and Joly F. 1958. Précipitations annuelles. *Atlas du Maroc*. Comité de Géographie du Maroc, Rabat.
- Gil García M. J., Las Heras R. T. and Ruiz Zapata B. 1996. Degradación antropogénica de la vegetación en el Puerto de la Morcuera (Sierra de Guadarrama, España) durante los últimos 2000 años, en base al análisis polínico. *Bol. Real Soc. Esp. Hist. Nat., Secc. Biol.* 92 (1–4): 29–36.
- Gómez Manzaneque F. 1988. Algunos táxones interesantes del suroeste madrileño. *Stud. Bot. Univ. Salamanca* 7: 257–261.
- Herranz Sanz J. M. and Gómez Campo C. 1986. Contribución al conocimiento de la flora y vegetación de la comarca de Alcaraz (Albacete). *Caja de Ahorros de Albacete*.
- Hultén E. and Fries M. 1986. *Atlas of north european vascular plants*. Vol. I. Koeltz Scientific Books, Königstein.
- Jaccard P. 1929. Considerations sur le coefficient générique et sa signification floristique et phytosociologique. *Bull. Soc. Bot. France* 76: 47–66.
- Jan du Chene R. 1976. Étude palynologique du Miocène supérieur andalou (Espagne). *Rev. Esp. Micropal.* 9 (1): 97–114.
- Julve Ph. 1993. Synopsis phytosociologique de la France (communautés de plantes vasculaires). *Lejeunia* (n.s.) 140: 1–160.
- Kopecký K. and Hejný S. 1974. A new approach to the classification of anthropogenic plant communities. *Vegetatio* 29: 17–20.
- Kopecký K., Dostalek J. and Frantik T. 1995. The use of the deductive method of syntaxonomic classification in the system of vegetational units of the Braun-Blanquet approach. *Vegetatio* 117: 95–112.
- Kuoch R. 1954. Wälder der Schweizer Alpen mit Verbreitungsgebiet der Weißtanne. *Mitt. Schweiz. Anst. Forstl. Versuchswes.* 30: 133–260.
- Little E. L. and Crithfield W. B. 1969. Subdivisions of the genus *Pinus* (Pines). U.S.D.A. Forest. Serv. Misc. Pub. 1144.

- López Vélez G. 1996. Flora y vegetación del macizo del Calar del Mundo y sierras adyacentes del sur de Albacete. Instituto de Estudios Albaceteños, Albacete.
- Losa Quintana J. M., Molero-Mesa J. and Casares M. 1986. El paisaje vegetal de Sierra Nevada. La cuenca alta del río Genil. Universidad de Granada.
- Mancebo J. M., Molina J. R. and Camino F. 1993. *Pinus sylvestris* L. en la vertiente septentrional de la sierra de Gredos (Ávila). *Ecología* 7: 233–245.
- Matuszkiewicz A. and W. 1954. Die Verbreitung der Waldassoziationen des Nationalparks von Bia-lowieza. *Polska Akad. Nauk, Kom. Ekol.* 2: 33–60.
- Matuszkiewicz W. and A. 1956a. Pflanzensoziologische Untersuchungen im Forstrevier „Ruda“ bei Pulawy (Polen). *Acta Soc. Botan. Polon.* 25: 331–400.
- Matuszkiewicz W. and A. 1956b. Zur Systematik der Quercetalia pubescentis-Gesellschaften in Polen. *Ebenda* 25: 27–72.
- Matuszkiewicz W. and A. 1981. Das Prinzip der mehrdimensionalen Gliederung der Vegetationseinheiten, erläutert am Beispiel der Eichen-Hainbuchenwälder in Polen. In Dierschke H. (ed.): *Syntaxonomie. Ber. Int. Symp. Veget. K. Rintelen*, 123–145.
- Millar C. I. 1993. Impact of the eocene on the evolution of *Pinus* L. *Ann. Missouri Bot. Gard.* 80: 471–498.
- Montserrat P. 1975. Comunidades relícticas geomorfológicas. *Anales Inst. Bot. Cavanilles* 32 (2): 397–404.
- Müller M. J. 1982. Selected climatic data for a global set of standard stations for vegetation science. Dr. W. Junk Publi., The Hague, Boston, London.
- Ninot J. M. 1996. Estudio fitocenológico del macizo del Turbón (Prepirineo central), I: Comunidades forestales. *Doc. Phytosociol. (n.s.)* 16: 215–239.
- Oberdorfer E. 1957. Süddeutsche Pflanzengesellschaften. *Pflanzensoziol. (Jena)* 10: 1–564.
- Oberdorfer E. 1990. Pflanzensoziologische Exkursionsflora. Ulmer, Stuttgart.
- Ojeda F., Arroyo J. and Marañón T. 1995. Biodiversity components and conservation of Mediterranean heathlands in southern Spain. *Biological Conservation* 72: 61–72.
- Passarge H. 1957. Waldgesellschaften des nördlichen Havellandes. *Wiss. Abh. Deut. Akad. Landwirtschaft. Wiss. Berlin* 26: 1–139.
- Pérez Latorre A. V., Galán de Mera A., Deil U. and Cabezudo B. 1996. Fitogeografía y vegetación del sector Aljibico (Cádiz-Málaga, España). *Acta Bot. Malacitana* 21: 241–267.
- Podani J. 1994. Multivariate data analysis in ecology and systematics. A methodological guide to the SYN-TAX 5.0 package. SPB Academic Publishing bv, The Hague.
- Preisig E. 1943. Die Waldgesellschaften des Warthe- und Weichsellandes. *Arb. Zentralstelle Veget. Kart. Reiches* 1943: 1–142.
- Regato Pajares P. and Escudero A. 1989. Caracterización fitoecológica de las comunidades de *Pinus nigra* subsp. *salzmannii* en los afloramientos rocosos del Sistema Ibérico meridional. *Bot. Complutensis* 15: 149–161.
- Regato Pajares P., Génova Fúster M. M. and Gómez Manzaneque F. 1992. Las representaciones relic-tas de *Pinus nigra* Arnold en el Sistema Central español. *Bol. Real Soc. Esp. Hist. Nat., Secc. Biol.* 88 (1–4): 63–71.
- Regato P., Gamisans J. and Gruber M. 1995. A syntaxonomical study of *Pinus nigra* subsp. *salzmannii* forests in the Iberian peninsula. *Phytocoenologia* 25 (4): 561–578.
- Rehder H., Gökçeoglu M., Gebauer G. and Güleriyüz G. 1994. Die Vegetation des Uludag-Gebirges (Anatolien). *Phytocoenologia* 24: 167–192.
- Rivas Goday S. 1955. Los grados de vegetación de la península Ibérica. *Anales Inst. Bot. Cavanilles* 13: 269–331.
- Rivas Goday S. and Borja Carbonell J. 1961. Estudio de la vegetación y flórula del macizo de Gúdar y Jabalambre. *Anales Inst. Bot. Cavanilles* 19: 1–550.
- Rivas-Martínez S. 1963. Estudio de la vegetación y flora de las sierras de Guadarrama y Gredos. *Anales Inst. Bot. Cavanilles* 21 (1): 13–225.
- Rivas-Martínez S. 1964. Esquema de la vegetación potencial y su correspondencia con los suelos en la España peninsular. *Anales Inst. Bot. Cavanilles* 22: 343–404.

- Rivas-Martínez S. 1975. Mapa de vegetación de la provincia de Ávila. *Anales Inst. Bot. Cavanilles* 32 (2): 1493–1556.
- Rivas-Martínez S. 1987. Memoria del mapa de series de vegetación de España. ICONA, Madrid.
- Rivas-Martínez S., Belmonte D., Cantó P., Fernández González F., Fuente V., Moreno J. M., Sánchez Mata D. and Sancho L. G. 1987. Piornales, enebrales y pinares oromediterráneos (*Pino-Cytision oromediterranei*) en el Sistema Central. *Lazaroa* 7: 93–124.
- Rivas-Martínez S. and Cantó P. 1987. Datos sobre la vegetación de las sierras de Guadarrama y Malagón. *Lazaroa* 7: 235–257.
- Rivas-Martínez S., Fernández González F., Sánchez Mata D. and Pizarro J. M. 1990. Vegetación de la sierra de Guadarrama. *Itinera Geobot.* 4: 3–132.
- Rivas-Martínez S., Bascónes J. C., Díaz T. E., Fernández-González F. and Loidi J. 1991. Vegetación del Pirineo occidental y Navarra. *Itinera Geobot.* 5: 5–456.
- Rojo y Alboreca A. and Montero González G. 1996. El pino silvestre en la sierra de Guadarrama. Ministerio de Agricultura, Pesca y Alimentación, Madrid.
- Sánchez Mata D. 1989. Flora y vegetación del macizo oriental de la sierra de Gredos (Ávila). Diputación Provincial de Ávila.
- Schuhwerk F. 1990. Relikte und Endemiten in Pflanzengesellschaften Bayerns – eine vorläufige Übersicht. *Ber. Bayer. Bot. Ges.* 61: 303–323.
- Valle F., Mota J. F. and Gómez-Mercado F. 1988. Datos sobre la vegetación orófila de Andalucía oriental (España). *Doc. Phytosociol.* (n.s.) 11: 459–464.
- Vigo J. 1979. Les forets de coniferes des Pyrenées catalanes. Essai de revision phytocenologique. *Doc. Phytosociol.* (n.s.) 4: 929–941.
- Wendelberger G. 1963. Die Relikt-Schwarzföhrenwälder des Alpenostrandes. *Vegetatio* 11: 265–287.
- Wilmanns O. 1997. Zur Geschichte der mitteleuropäischen Trockenrasen seit dem Spätglazial – Methoden, Tatsachen, Hypothesen. *Phytocoenologia* 27 (2): 213–233.