

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 43 (1968)

Artikel: Canonical Vector Fields on Spheres.
Autor: Zvengrowski, P.
DOI: <https://doi.org/10.5169/seals-32927>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 28.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Canonical Vector Fields on Spheres

P. ZVENGROWSKI

§ 1. Introduction

We are interested in norm-preserving bilinear forms

$$M: R^r \otimes R^n \rightarrow R^n,$$

where $\otimes = \otimes_R$ and by norm preserving we mean $\|M(u \otimes v)\| = \|u\| \cdot \|v\|$. Such a form implies the existence of $r-1$ mutually orthonormal vector fields on S^{n-1} (see 1.2 below). Given n , the question of finding the largest r so that such a form exists was solved in 1923 by RADON [5], by HURWITZ [3], and again in 1942 by ECKMANN [2]. The methods of RADON and HURWITZ yield complicated iterative schemes for actually constructing the forms, which have recently been simplified by ADAMS, LAX, and PHILLIPS [1]. We now give a still simpler construction and prove certain relevant properties of the "canonical" vector fields thus obtained. In particular, they are closed under the intrinsic join operations of JAMES [4] (cf. Prop. 4.4).

Let M be a form as above and let e_0, \dots, e_{r-1} be an orthonormal basis for R^r . Then one obtains r orthogonal transformations $M_0, \dots, M_{r-1} \in O(n)$ by defining

$$M_i(v) = M(e_i \otimes v), \quad 0 \leq i \leq r-1, \quad v \in R^n.$$

Conversely, M is defined by the M_i using the formula

$$M(u \otimes v) = \sum \alpha_i M_i(v), \quad \text{where } u = \sum \alpha_i e_i \quad \text{and} \quad i = 0, \dots, r-1.$$

1.1. THEOREM: *The following are equivalent*

A: M is norm-preserving,

B: $\langle M_i(v), M_j(v) \rangle = \delta_{ij} \|v\|^2 \quad \forall 0 \leq i, j \leq r-1$ and $v \in R^n$,

C: $M_i \in O(n)$ and $M_i^t M_j + M_j^t M_i = 0, \quad i \neq j$.

This theorem has been used in one form or another by most of the above authors, and its proof is omitted.

One can assume without loss of generality that $M_0 = \text{id}$, by following M with M_0^{-1} if necessary. Then from (B) it follows that $\langle v, M_i(v) \rangle = 0, \quad 1 \leq i \leq r-1$, and hence if we restrict v to S^{n-1} , i.e. $\|v\| = 1$, we obtain

1.2. COROLLARY: $M_1(v), \dots, M_{r-1}(v)$ define a family of $r-1$ orthonormal vector fields on S^{n-1} .

Furthermore, using (C) together with $M_0 = \text{id}$ and $M_i^t M_i = 1$, we obtain

1.3. COROLLARY: $M_i + M_i^t = 0, \quad M_i^2 = -1, \quad M_i M_j + M_j M_i = 0, \quad 1 \leq i, j \leq r-1$.

1.4. DEFINITION: A norm preserving form $M: R^r \otimes R^n \rightarrow R^n$ is orthogonal to the identity if $\langle v, M(u \otimes v) \rangle = 0 \forall u \in R^r, v \in R^n$.

From the above remarks such a form is clearly equivalent to the existence of a norm preserving form $M'_0 = \text{id}$ and $M'_i = M_{i-1}$, $i \geq 1$. Furthermore, M then defines r orthonormal vector fields on S^{n-1} and M_i, M_j satisfy 1.3, $0 \leq i, j \leq r-1$.

We will use the notation $M_u = M(u \otimes -): R^n \rightarrow R^n$, $u \in R^r$. Clearly $M_u / \|u\| \in O(n)$, and if M is orthogonal to id then M_u is antisymmetric. In all cases one has the following identity:

$$\langle M(u \otimes v_1), M(u \otimes v_2) \rangle = \langle M_u v_1, M_u v_2 \rangle = \|u\|^2 \left\langle \frac{M_u}{\|u\|} v_1, \frac{M_u}{\|u\|} v_2 \right\rangle = \|u\|^2 \langle v_1, v_2 \rangle.$$

§ 2. Tensor Products of Inner Product Spaces

Let V, W be inner product spaces over a field F . Then $V \otimes_F W$ is an inner product space, where

$$\langle v_1 \otimes w_1, v_2 \otimes w_2 \rangle = \langle v_1, v_2 \rangle \langle w_1, w_2 \rangle.$$

In case $V = R^m$ and $W = R^n$, with their usual products, it is not hard to see that the resulting inner product on $R^{m \cdot n}$ is also the usual one.

The following lemma will be exceedingly useful in the proof of Theorem 3.1.

2.1. ORTHOGONALITY LEMMA: Let V, W be inner product spaces with commutative inner products and suppose $A: V \rightarrow V$ and $B: W \rightarrow W$ are endomorphisms such that

- (i) A is orthogonal to id_V , that is $\langle v, Av \rangle = 0 \forall v \in V$, or B is orthogonal to id_W
- (ii) A is symmetric and B antisymmetric, or vice-versa.

Then the two endomorphisms $\varphi = A \otimes 1$ and $\psi = 1 \otimes B$ of $V \otimes W$ are orthogonal, that is $\langle \varphi a, \psi a \rangle = 0 \forall a \in V \otimes W$.

Proof: Let $a = \sum_i v_i \otimes w_i$. Then

$$\begin{aligned} \langle \varphi a, \psi a \rangle &= \left\langle \sum_i A v_i \otimes w_i, \sum_j v_j \otimes B w_j \right\rangle \\ &= \sum_{i,j} \langle A v_i, v_j \rangle \langle w_i, B w_j \rangle. \end{aligned}$$

Now (i) clearly implies that the terms where $i=j$ vanish. Then supposing $A^t = A$, $B^t = -B$, we have

$$\begin{aligned} \langle \varphi a, \psi a \rangle &= \sum_{i < j} (\langle A v_i, v_j \rangle \langle w_i, B w_j \rangle + \langle A v_j, v_i \rangle \langle w_j, B w_i \rangle) \\ &= \sum_{i < j} (\langle v_j, A v_i \rangle \langle B w_j, w_i \rangle + \langle v_j, A v_i \rangle \langle -B w_j, w_i \rangle) \\ &= 0. \end{aligned}$$

REMARK: The representation $a = \sum_{i=1}^t v_i \otimes w_i$ is of course not unique. One can,

however, always choose it so that v_1, \dots, v_t form a given basis of V , or similarly for the w_i (but not both).

§ 3. The Basic Construction

Let $C: R^8 \otimes R^8 \rightarrow R^8$ be the Cayley multiplication. Let $i: R^7 \rightarrow R^8$ be inclusion into the last seven co-ordinates, then $C \circ (i \otimes 1): R^7 \otimes R^8 \xrightarrow{C_1} R^8$ is a norm preserving multiplication orthogonal to the identity. Now define a form $N: R^7 \otimes R^{16} \rightarrow R^{16}$ by the composition

$$R^7 \otimes R^{16} \xrightarrow{\approx} R^7 \otimes R^8 \oplus R^7 \otimes R^8 \xrightarrow{C_1 \otimes (-C_1)} R^8 \oplus R^8 \xrightarrow{\approx} R^{16}.$$


Clearly N is norm preserving, orthogonal to id, and for $0 \leq i \leq 6$ each N_i is antisymmetric. Furthermore, N_i has the form

$$N_i = \begin{pmatrix} B_i & 0 \\ 0 & -B_i \end{pmatrix}, \quad B_i \in O(8).$$

3.1. THEOREM: Let $M: R^r \otimes R^n \rightarrow R^n$, n even, be a norm-preserving form such that (a) M is orthogonal to id

(b) $M_i U = -U M_i$, $0 \leq i \leq r-1$, where $U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in O(n)$

Then the form \bar{M} defined by the composition below is norm preserving and also satisfies (a), (b), (relative to $r+8$ and $16n$):



Proof: Condition (b) follows readily from the fact that $\bar{M}_i = \begin{pmatrix} A_i & 0 \\ 0 & -A_i \end{pmatrix}$, $0 \leq i < r+7$, while $\bar{M}_{r+7} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Let $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = T$, $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = V \in O(16)$.

From (b) it follows that $M_u U = -U M_u \forall u \in R^r$. Then $U M_u$ is symmetric. Similarly, since $N_i = \begin{pmatrix} B_i & 0 \\ 0 & -B_i \end{pmatrix}$ satisfies (b), $0 \leq i \leq 6$, one sees that $V N_u$ is antisymmetric $\forall u \in R^7$ and $T N_u$ antisymmetric. Also, $V T$ is symmetric.

Now, starting with $u \otimes v \in R^{r+8} \otimes R^{16n}$, let $u = u_1 \oplus u_2 \oplus u_3 \in R^r \oplus R^7 \oplus R^1$ and $v = \sum_i v'_i \otimes v''_i \in R^n \otimes R^{16}$. Then $\bar{M}(u \otimes v) = a + b + c$, where

$$\begin{aligned} a &= \sum_j M(u_1 \otimes v'_j) \otimes T v''_j, \\ b &= \sum_k U v'_k \otimes N(u_2 \otimes v''_k), \\ c &= u_3 \sum_i v'_i \otimes V v''_i. \end{aligned}$$

To prove (a), we show $\langle v, a \rangle = \langle v, b \rangle = \langle v, c \rangle = 0$. $\langle v, a \rangle = \sum_{i,j} \langle v'_i, M_{u_1}(v'_j) \rangle \langle v''_i, T v''_j \rangle$.

Choosing $v''_i = e_i$, the standard basis for R^{16} , $\langle v''_i, T v''_j \rangle = \pm \delta_{ij}$ and $\langle v, a \rangle = \sum_i \pm \langle v'_i, M_{u_1}(v'_j) \rangle = 0$ since M is orthogonal to the identity. $\langle v, b \rangle =$

$\sum_{i,1} \langle v'_i, U v'_k \rangle \langle v''_i, N_{u_2} v''_k \rangle = 0$ by the orthogonality lemma. $\langle v, c \rangle = u_3 \sum_{i,1} \langle v'_i, v'_i \rangle$

$\langle v''_i, V v''_i \rangle = 0$ by choosing $\{v'_i\}$ orthonormal and noticing that V is orthogonal to id.

To show that \bar{M} is norm preserving, we first prove that $\langle a, b \rangle = \langle a, c \rangle = \langle b, c \rangle = 0$.

$$\begin{aligned} \langle a, b \rangle &= \sum_{j,k} \langle M_{u_1}(v'_j), U v'_k \rangle \langle T v''_j, N_{u_2}(v''_k) \rangle \\ &= \sum_{j,k} \langle U M_{u_1}(v'_j), v'_k \rangle \langle v''_j, T N_{u_2}(v''_k) \rangle. \end{aligned}$$

Choosing $v''_i = e_i$ as before, one has $\langle T v''_i, N_{u_2}(v''_i) \rangle = \pm \langle v''_i, N_{u_2}(v''_i) \rangle = 0$. Thus one need only consider the terms where $j \neq k$, which sum to zero since $U M_{u_1}$ is symmetric and $T N_{u_2}$ antisymmetric. The other two orthogonality relations are proved quite analogously, where in $\langle b, c \rangle$ one takes $\{v'_i\}$ to be the standard basis for R^n to insure that the (i, i) terms vanish. Thus

$$\|\bar{M}(u \otimes v)\|^2 = \|a\|^2 + \|b\|^2 + \|c\|^2.$$

Choosing $v''_i = e_i$, one easily sees that the individual terms in a, b, c are mutually orthogonal, being already orthogonal in the second factor. Then, since T, U , and V are all orthogonal transformations,

$$\begin{aligned} \|\bar{M}(u \otimes v)\|^2 &= \sum_i \|u_1\|^2 \|v'_i\|^2 \|v''_i\|^2 + \sum_i \|v'_i\|^2 \|u_2\|^2 \|v''_i\|^2 + u_3^2 \sum_i \|v'_i\|^2 \|v''_i\|^2 \\ &= (\|u_1\|^2 + \|u_2\|^2 + u_3^2) \sum_i \|v'_i\|^2 \|v''_i\|^2 \\ &= \|u\|^2 \|v\|^2. \end{aligned}$$

3.2. COROLLARY: If $n = s \cdot 2^{4a+b}$, s odd, $0 \leq b \leq 3$, then S^{n-1} admits $8a + 2^b - 1$ orthonormal vector fields.

Proof: If $n = s$ one has a trivial form $R^0 \otimes R^s \xrightarrow{0} R^s$. Applying the theorem “ a ” times gives a norm preserving form orthogonal to the identity

$$R^{8a} \otimes R^{s \cdot 16^a} \rightarrow R^{s \cdot 16^a}$$

(the fact that n is odd on the first iteration causes no trouble since $r = 0$ there). This is the case $b = 0$. For $b = 1, 2, 3$ one need only apply the theorem once more and observe that $\bar{M}(\mu R^{8a+2^b-1} \otimes \mu R^{s \cdot 16^a \cdot 2^b}) \subset \mu R^{s \cdot 16^a \cdot 2^b}$, where μ denotes the generic inclusion of R^m into the first m co-ordinates of R^{m+k} for any m, k . This is so because $N(\mu R^{2^b-1} \otimes \mu R^{2^b}) \subset \mu R^{2^b}$, $b = 1, 2, 3$, corresponding to the complex numbers, quaternions, and Cayley numbers respectively.

REMARK: $\varrho(n) = 8a + 2^b$ is called the Radon-Hurwitz function.

§ 4. Definition and Properties of Canonical Vector Fields

Let $R^\infty = \lim_{\rightarrow} R^m$. It is clear, using the definition of \bar{M} , that the following commutes:

$$\begin{array}{ccc} R^r \otimes R^n & \xrightarrow{M} & R^m \\ \downarrow \mu \otimes \mu & & \downarrow \mu \\ R^{r+8} \otimes R^{16n} & \xrightarrow{\bar{M}} & R^{16n} \end{array}$$

Starting with $R^0 \otimes R^1 \xrightarrow{0} R^1$, we now iterate Theorem 3.1 and pass to the limit, obtaining a norm preserving multiplication orthogonal to the identity

$$M: R^\infty \otimes R^\infty \rightarrow R^\infty.$$

Let $\mu_i: R^m \hookrightarrow R^\infty$ be the inclusion of R^m into the i 'th block of m co-ordinates, $0 \leq i$. Thus, for $i \leq n$, one has a commutative diagram

$$\begin{array}{ccc} R^m \hookrightarrow & \mu_i & \rightarrow R^\infty \\ \downarrow (\) \otimes e_i & & \uparrow \mu \\ R^m \otimes R^n & \hookrightarrow & R^{m \cdot n} \end{array}$$

Also, $\mu_0 = \mu$.

The following theorem says that M in effect gives a maximal family of orthonormal vector fields on S^{n-1} for every n .

4.1. THEOREM: If $r \leq \varrho(n) - 1$ then, for any $i \geq 0$,

$$M(\mu R^r \otimes \mu_i R^n) \subset \mu_i R^n.$$

Proof: This property will certainly hold for n if it is true for some divisor of n , the same r , and all i . Letting $n = s \cdot 2^{4a+b}$, s odd, $0 \leq b \leq 3$, it will thus suffice to prove the theorem for 2^{4a+b} , since also $\varrho(2^{4a+b}) = \varrho(n)$. In other words, we can assume without loss of generality that $n = 2^{4a+b}$. Furthermore, if the result holds for $r = \varrho(n) - 1$ it will certainly hold for smaller r , so we also take $r = \varrho(n) - 1 = 8a + 2^b - 1$.

First consider $b = 0$ and let e_0, e_1, \dots be the usual basis for R^∞ . We shall prove that if the result holds for $0 \leq i \leq 16^m - 1$ then it also holds for $0 \leq i \leq 16^{m+1} - 1$, giving an inductive proof of the theorem for the case $b = 0$ (clearly $m = 0$ furnishes a base for the induction). Write $i = t \cdot 16^m + s$, where $0 \leq s \leq 16^{m+a+1}$ and $0 \leq t \leq 15$. The inclusion $R^n = R^{16^a \mu_i} \rightarrow R^{16^{m+a+1}}$ corresponds to the composition

$$R^{16^a \mu_s} \rightarrow R^{16^{m+a}} \xrightarrow{(\cdot) \otimes e^t} R^{16^{m+a}} \otimes R^{16} \xrightarrow{\approx} R^{16^{m+a+1}}.$$

Then in the passage from M to \bar{M} , i.e., from $R^{8(m+a)-1} \otimes R^{16^{m+a}}$ to $R^{8(m+a+1)-1} \otimes R^{16^{m+a+1}}$, we have a commutative diagram

$$\begin{array}{ccc}
 R^{8(m+a+1)-1} \otimes R^{16^{m+a+1}} & \xrightarrow{\approx} & R^{8(m+a)-1} \oplus R^8 \otimes R^{16^{m+a}} \otimes R^{16} \\
 \uparrow \mu \otimes \mu_i & \nearrow (\mu, 0) \otimes \mu_s \otimes e_t & \downarrow \approx \\
 R^{8a-1} \otimes R^{16^a} & \xrightarrow{(\mu \otimes \mu_s) \otimes e_t, 0} & (R^{8(m+a)-1} \otimes R^{16^{m+a}}) \otimes R^{16} \oplus (R^{16^{m+a}} \otimes (R^8 \otimes R^{16}))
 \end{array}$$

Performing the multiplications and applying the inductive hypothesis, we find

$$M(\mu R^{8a-1} \otimes \mu_i R^{16^a}) \subset M(\mu R^{8a} \otimes \mu_s R^{16^a}) \otimes e_t \subset \mu_s R^{16^a} \otimes e_t,$$

and the latter corresponds to $\mu_i R^{16^a}$ under the isomorphism

$$R^{16^{(m+a)}} \otimes R^{16} \approx R^{16^{m+a+1}}.$$

This completes the proof for $b = 0$. A similar method works for $b = 1, 2, 3$. For example, if $b = 2$, we use the existence of quaternions to establish the cases $0 \leq i \leq 3$ (similar to the proof of Cor. 3.2) as base for the induction, then pass from $0 \leq i \leq 4 \cdot 16^m - 1$ to $0 \leq i \leq 4 \cdot 16^{m+1} - 1$

4.2. COROLLARY: *If $r \leq \varrho(n) - 1$ then the following composition defines a norm preserving multiplication orthogonal to id:*

$$R^r \otimes R^n \xrightarrow{\mu \otimes \mu_i} R^\infty \otimes R^\infty \xrightarrow{M} R^\infty \xrightarrow{\mu_i^{-1}} R^n.$$

Denoting this multiplication “ $\mathbf{M}_{i,r}^n$ ”, let us call the resultant r orthonormal vector fields on S^{n-1} “ $f_{i,r}^n$ ”. More precisely, $f_{i,r}^n: S^{n-1} \rightarrow V_{n,r+1}$ is the cross section of the fibration $V_{n,r+1} \rightarrow S^{n-1}$ such that

$$f_{i,r}^n(x) = \begin{pmatrix} x \\ x_0 \\ \vdots \\ x_{r-1} \end{pmatrix} \in V_{n,r+1}, \quad \text{where } x_j = \mathbf{M}_{i,r}^n(e_j \otimes x).$$

These are our “canonical” vector fields.

4.3. DEFINITION: Let $M: R^r \otimes R^m \rightarrow R^m$ and $N: R^r \otimes R^n \rightarrow R^n$ be norm preserving forms orthogonal to the identity. Then their intrinsic join $M*N$ is the composition

$$R^r \otimes (R^m \oplus R^n) \xrightarrow{\cong} R^r \otimes R^m \oplus R^r \otimes R^n \xrightarrow{M \oplus N} R^m \oplus R^n \xrightarrow{\cong} R^{m+n}.$$

Clearly $M*N$ is also norm preserving and orthogonal to id. If $f: S^{m-1} \rightarrow V_{m,r+1}$ and $g: S^{n-1} \rightarrow V_{n,r+1}$ are the corresponding cross sections, then their intrinsic join $f*g$ is defined as the composition

$$S^{m+n-1} \xleftarrow[\varphi]{\cong} S^{m-1} * S^{n-1} \xrightarrow{f*g} V_{m,r+1} * V_{n,r+1} \xrightarrow{\varphi} V_{m+n,r+1},$$

φ being the intrinsic join map of JAMES [4]. One easily sees that $f*g$ corresponds to $M*N$, and it is then clear that the canonical vector fields can be joined together in many ways to give other canonical fields. A typical example is the formula

$$f_{0,3}^4 * f_{1,3}^4 = f_{0,3}^8.$$

More generally, one can easily establish the following.

4.4. PROPOSITION: $f_{i,m,r}^n * f_{i,m+1,r}^n * \dots * f_{i,m+m-1,r}^n = f_{i,r}^{m \cdot n}$

REFERENCES

[1] J.F. ADAMS, P.D. LAX, and R.S. PHILLIPS, *On Matrices Whose Real Linear Combinations are Non-singular*, Proc. Am. Math. Soc. 16 (1965), 318–322.
 [2] B. ECKMANN, *Beweis des Satzes von Hurwitz-Radon*, Comment. Math. Helv. 15 (1942), 358–366.
 [3] A. HURWITZ, *Über die Komposition der quadratischen Formen*, Math. Ann. 88 (1923), 1–25.
 [4] I.M. JAMES, *The Intrinsic Join*, Proc. London Math. Soc. (3), 8 (1958), 507–535.
 [5] J. RADON, *Lineare Scharen orthogonaler Matrizen*, Abh. Math. Sem. Hamburg 1 (1923), 1–14.

Mathematics Research Institute
E.T.H., Zürich

University of Illinois