

Zeitschrift: Elemente der Mathematik
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 44 (1989)
Heft: 5

Artikel: Winch curves
Autor: Sanders, R.
DOI: <https://doi.org/10.5169/seals-41620>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 01.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Let $z = (x - y)/\|x - y\|$. Then $\gamma_z(E) = D$, $\gamma_z(C) = C$ and $\gamma_z(C \setminus D) = C \setminus D$. Hence, by (1),

$$\begin{aligned} \mu(\gamma_z(M) \cap C) &= \mu(\gamma_z(M) \cap (C \setminus D)) + \mu(\gamma_z(M) \cap D) \\ &\geq \mu(M \cap (C \setminus D)) + \mu(M \cap D) = \mu(M \cap C) + \mu(M \cap D) > \mu(M \cap C). \end{aligned}$$

Since $\gamma_z(M) \in K$, this contradicts the choice of M , so the proof is complete. \square

Let us remark that a slight variant of the proof above gives the following assertion. Let K be a non-empty closed subset of H which is also closed under the operators γ_z , i.e. which is such that $\gamma_z(A) \in K$ for all $A \in K$ and $z \in S^n$. Then K contains all caps of measure $m = \sup \{ \mu(A) : A \in K \}$.

Also, it is easily seen that the proof above implies various extensions of Theorem 3. For example, given finite sets $X, Y \subset S^n$ with $|X| = |Y|$, let us write $X \leq Y$ if for every $d > 0$, the number of pairs in X at distance at least d is not more than the number of pairs in Y at distance at least d . Furthermore, for sets $A, B \subset S^n$, let us write $A \leq B$ for the assertion that for every finite set $X \subset A$ there is a finite set $Y \subset B$ with $|Y| = |X|$ and $X \leq Y$. Then the following assertion holds. Let A be a non-empty closed subset of S^n and let C be a cap of measure $\mu(A)$. Then $C \leq A$.

Béla Bollobás

Department of Pure Mathematics and Mathematical Statistics
University of Cambridge, England

REFERENCES

- 1 Baernstein A., Taylor B. A.: Spherical rearrangements, subharmonic functions and *-functions in n -space, *Duke Math. J.* 43, 245–268 (1976).
- 2 Benyamini Y.: Two point symmetrization, the isoperimetric inequality on the sphere and some applications, *Longhorn Notes*, The University of Texas, Texas Functional Analysis Seminar, pp. 53–76, 1983–1984.
- 3 Bollobás B.: *Combinatorics*, Cambridge University Press, xii + 177, Cambridge, England, 1986.
- 4 Frankl P.: The shifting technique in extremal set theory, in «*Surveys in Combinatorics 1987*» (Whitehead C., ed.), LMS Lecture Note Series 123, Cambridge University Press, pp. 81–110, Cambridge, 1987.

Winch curves

A taut rope connects a point in the origin of a rectangular coordinate system with a point in $R(a, 0)$. If the latter starts moving along the line $x = a$, it will trail the point in the origin. For each point P of the curve that is created in this way we have $PQ = a$, where Q is the intersection of the tangent to the curve in P with the line $x = a$. This curve, known as the tractrix, is represented by an equation that can be found as follows.

In the rectangular triangle PSQ (see fig. 1) we have

$$PQ = a, \quad PS = a - x, \quad SQ = (a - x) dy/dx.$$




Figure 1. Winch curve.

Pythagoras' theorem leads to

$$a^2 = (a - x)^2 [1 + (dy/dx)^2].$$

Solving this differential equation for positive a we obtain

$$y = -\sqrt{a^2 - (a - x)^2} + a \ln C_1 \frac{a + \sqrt{a^2 - (a - x)^2}}{a - x}$$

as a solution in the first quadrant.

As the curve passes through the origin we have $y = 0$ for $x = 0$. This gives $C_1 = 1$, so the equation can be written

$$y = -\sqrt{a^2 - (a - x)^2} + a \ln \frac{a + \sqrt{a^2 - (a - x)^2}}{a - x} \tag{1}$$

In literature the tractrix is usually described as above.

In this article however we treat the trailing problem in a more general way. It will then turn out that the tractrix can be considered as a special case of a new family of curves. In order to achieve this we assume that the length of the rope PQ will be changed during trailing. In practice one could think of a vehicle Q trailing a load P and being equipped with a winch so that PQ can be shortened or lengthened during the trailing process. Usually Q will be moving at constant velocity and the length of PQ will be varied uniformly in time as well. We then have

$$RQ = y + (a - x) dy/dx = c_1 t \tag{2}$$

where t represents the time and c_1 stands for the velocity of Q along $x = a$. Furthermore we can write

$$PQ = a - c_2 t \quad (3)$$

where c_2 is the velocity with which the winch is wound up ($c_2 > 0$) or eased off ($c_2 < 0$). Elimination of t between equations (2) and (3) gives

$$PQ = a - b[y + (a - x)dy/dx]$$

with $b = c_2/c_1$.

It should be noted that the same result is obtained if the variation of the length of PQ and the velocity of Q are not constant in time on condition that the time dependencies in equations (2) and (3) are of the same form.

Writing $p = dy/dx$ and applying Pythagoras' theorem in the triangle PSQ we have

$$\{a - b[y + (a - x)p]\}^2 = (a - x)^2(1 + p^2)$$

or in the first quadrant

$$a - b[y + (a - x)p] = (a - x)\sqrt{1 + p^2}. \quad (4)$$

This equation can be written in the form $y = g(p)x + f(p)$ and is called the differential equation of d'Alembert.

Differentiating with respect to x we obtain

$$-bp - b(a - x)dp/dx + bp = -\sqrt{1 + p^2} + (a - x)\frac{p}{\sqrt{1 + p^2}}dp/dx$$

or

$$dp/dx = \frac{\sqrt{1 + p^2}}{(a - x)\left[b + \frac{p}{\sqrt{1 + p^2}}\right]}. \quad (5)$$

Integration of (5) leads to

$$-\ln(a - x) = b \ln(p + \sqrt{1 + p^2}) + \frac{1}{2} \ln(1 + p^2) + C_2.$$

Now $p = 0$ for $x = 0$, yielding $C_2 = -\ln a$, so we can write

$$\ln \frac{a}{a - x} = \ln \sqrt{1 + p^2} \{p + \sqrt{1 + p^2}\}^b$$

or

$$x = a \left[1 - \frac{1}{\sqrt{1 + p^2} \{p + \sqrt{1 + p^2}\}^b} \right] \quad (6)$$




Figure 2. Winch curves for different values of b . The numbers along the framework are given in units a .

and from equation (4) we find for $b \neq 0$

$$y = \frac{a}{b} \left[1 - \frac{pb + \sqrt{1+p^2}}{\sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b} \right]. \tag{7}$$

Equations (6) and (7) may be considered as a parameter representation of the new family of *winch curves* with parameter $p = dy/dx$.

We will now check that equations (6) and (7) yield the tractrix for $b \rightarrow 0$. For $b \rightarrow 0$ equation (6) leads to

$$x = a - \frac{a}{\sqrt{1+p^2}}. \tag{8}$$

As equation (7) is of indeterminate form for $b \rightarrow 0$ we rewrite it as

$$y = a \left[\frac{\sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b - pb - \sqrt{1+p^2}}{b \sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b} \right].$$

Now De l'Hôpital's rule gives

$$\begin{aligned} y &= \lim_{b \rightarrow 0} a \left[\frac{\sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b \ln(p + \sqrt{1+p^2}) - p}{\sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b + b \sqrt{1+p^2} \{p + \sqrt{1+p^2}\}^b \ln(p + \sqrt{1+p^2})} \right] \\ &= a \left[\ln(p + \sqrt{1+p^2}) - \frac{p}{\sqrt{1+p^2}} \right]. \end{aligned} \tag{9}$$




Figure 3. Cusped winch curves.

Eliminating p between (8) and (9) we arrive at equation (1). So the tractrix may indeed be considered as a winch curve with $b = 0$.

Figures 2 and 3 show a number of winch curves for different b .

It is assumed that the rope PQ remains tight even if it is eased off. As a consequence the curves will not or not exclusively be found in the first quadrant for $b < 0$.

In some of the curves a cusp A is shown. The coordinates of such a point can be found as follows.

Equation (6) shows that x is a continuous function of p . In A we have $dx/dp = 0$ so dp/dx goes to infinity. From equation (5) we then obtain

$$b + \frac{p_A}{\sqrt{1 + p_A^2}} = 0$$

or

$$p_A = \frac{-b}{\sqrt{1 - b^2}} \tag{10}$$

giving $-1 < b < 0$ for curves with a cusp since p_A is positive, real and finite.

Equation (10) substituted in (6) and (7) leads to the coordinates of A :

$$x_A = a[1 - \sqrt{(1 - b)^{1-b}(1 + b)^{1+b}}]$$

$$y_A = \frac{a}{b}[1 - \sqrt{(1 - b)^{2-b}(1 + b)^{2+b}}].$$

For $b \rightarrow -1$ point A approaches $(-a, -a)$ and P moves, after it has passed A , along a line that approaches the line $x = -a$.

For $b = -1$ the curve passes through $(-a, -a)$, here p is infinite.

For $b < -1$ the winch is eased off so quickly that P cannot be trailed anymore.

It will be clear that P and Q only meet each other for positive b . From (6) and (7) it can be derived that this will happen at point $(a, a/b)$.

Thanks are due to Mr H. J. de Vries for performing many calculations.

R. Sanders, Zuidwolde, The Netherlands

Kleine Mitteilung

Über die Zahlenfolge $n! + k$, $2 \leq k \leq n$

Fast jedes Buch über Zahlentheorie erwähnt die Tatsache, dass keine der Zahlen

$$n! + k \quad \text{mit} \quad 2 \leq k \leq n \tag{1}$$

eine Primzahl ist (es gibt also in der Folge der Primzahlen beliebig lange Lücken). Es scheint aber nicht allgemein bekannt zu sein, dass dieselbe Zahlenfolge auch die Unendlichkeit der Primzahl-Menge beherbergt. Dies entnimmt man dem folgenden

Satz 1. Für jedes $n > 1$ und $2 \leq k \leq n$ hat $n! + k$ entweder einen Primfaktor $> n$, oder aber k ist prim und grösser als $n/2$ und $n! + k$ ist eine Potenz von k .

Beweis. Sei $2 \leq k \leq n$. Für alle Primzahlen $p \leq n$, welche $n! + k$ teilen, ist $p|k$. Falls $p < k$, also $p \in \{2, \dots, k-1\}$ ist, gilt

$$p|n!/k \quad \text{und damit} \quad p \nmid (n!/k) + 1 = (n! + k)/k.$$

Die Zahl $(n! + k)/k$ und mit ihr $n! + k$ besitzt somit Primteiler $> n$. Hat also $n! + k$ nur Primteiler $\leq n$, so ist k prim und dies ist der einzige Primteiler von $n! + k$.

Aus

$$n! + k = k^s, \quad s \geq 2$$

folgt

$$k \nmid k^{s-1} - 1 = n!/k, \quad \text{also} \quad k > n/2.$$