

**Zeitschrift:** Elemente der Mathematik  
**Band:** 60 (2005)

**Artikel:** Ky Fan's inequality with binomial expansion  
**Autor:** Rooin, Jamal  
**DOI:** <https://doi.org/10.5169/seals-10208>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 08.11.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

---

## Ky Fan's inequality with binomial expansion

---

---

Jamal Rooin

Jamal Rooin received his M.S. from the Teacher Education University in Teheran, Iran, in 1989. In 1997 he became a Ph.D. student at that university, where he graduated in the year 2002. He is now assistant professor at the Institute for Advanced Studies in Basic Sciences in Zanjan, Iran. His mathematical research interests include the theory of inequalities, problems in analysis and functional analysis.

In this note, given  $n$  arbitrary nonnegative real numbers  $x_1, \dots, x_n$ , we denote by  $A_n$  and  $G_n$ , the unweighted arithmetic and geometric means of  $x_1, \dots, x_n$ , respectively, i.e.,

$$A_n = \frac{1}{n} \sum_{i=1}^n x_i, \quad G_n = \sqrt[n]{\prod_{i=1}^n x_i},$$

and moreover, if  $x_i \in (0, 1/2]$ , we denote by  $A'_n$  and  $G'_n$ , the unweighted arithmetic and geometric means of  $1 - x_1, \dots, 1 - x_n$ , respectively, i.e.,

$$A'_n = \frac{1}{n} \sum_{i=1}^n (1 - x_i), \quad G'_n = \sqrt[n]{\prod_{i=1}^n (1 - x_i)}.$$

In 1961 the following remarkable inequality, due to Ky Fan, was published for the first time in the well-known book *Inequalities* by Beckenbach and Bellman [2, p. 5]:

If  $x_i \in (0, 1/2]$ , then

$$\frac{A'_n}{G'_n} \leq \frac{A_n}{G_n}, \quad (1)$$

with equality holding if and only if  $x_1 = \dots = x_n$ .

Die Ungleichung von Ky Fan ist 1961 in der bekannten Enzyklopädie von Beckenbach und Bellman zum ersten Mal publiziert worden. Ausgangspunkt ist die Ungleichung zwischen dem arithmetischen und dem geometrischen Mittel, kurz AGM-Ungleichung. Die Ungleichung von Ky Fan besagt, dass sich die Diskrepanz zwischen dem arithmetischen und dem geometrischen Mittel verkleinert, wenn die gegebenen Zahlen  $x_k \in (0, \frac{1}{2}]$  durch  $x'_k := 1 - x_k$  ersetzt werden. In der vorliegenden Note wird ein weiterer, besonders einfacher Beweis dieser Ungleichung gegeben.

Inequality (1) has evoked the interest of several mathematicians and in numerous articles new proofs, extensions, refinements and various related results have been published; see the survey paper [1] and the references therein.

By replacing the apparent binomial expansion, it is easily established that the following recursive identity holds (see [3]):

$$A_n - G_n = \frac{1}{n} \sum_{k=2}^n \binom{n}{k} A_{n-1}^{\frac{n-k}{n}} \left(x_n^{1/n} - A_{n-1}^{1/n}\right)^k + x_n^{1/n} \left(A_{n-1}^{\frac{n-1}{n}} - G_{n-1}^{\frac{n-1}{n}}\right). \quad (2)$$

Using this identity, we now give a striking new proof of (1) which shows the power of binomial expansion on the one hand and the close relations between AGM and Ky Fan inequalities on the other:

Actually, dividing each side of (2) by

$$G_n = G_{n-1}^{\frac{n-1}{n}} x_n^{\frac{1}{n}} \neq 0,$$

we get the following recursive identity

$$\frac{A_n}{G_n} = \frac{1}{n} \sum_{k=2}^n \binom{n}{k} \left(\frac{A_{n-1}}{G_{n-1}}\right)^{\frac{n-k}{n}} \frac{\left(x_n^{\frac{1}{n}} - A_{n-1}^{\frac{1}{n}}\right)^k}{G_{n-1}^{\frac{k-1}{n}} x_n^{\frac{1}{n}}} + \left(\frac{A_{n-1}}{G_{n-1}}\right)^{\frac{n-1}{n}}. \quad (3)$$

Now, we proceed by induction on  $n$ . If  $n = 1$ , there is nothing to prove. Suppose that  $n \geq 2$  and (1) holds for  $n - 1$ . If  $x_1 = \dots = x_n$ , then obviously equality holds in (1). Let  $x_i$ 's not all be equal. By a rearrangement if necessary, we can suppose that  $x_n = \max_{1 \leq i \leq n} x_i$ .

Now, considering

$$\frac{A'_n}{G'_n} = \frac{1}{n} \sum_{k=2}^n \binom{n}{k} \left(\frac{A'_{n-1}}{G'_{n-1}}\right)^{\frac{n-k}{n}} \frac{\left((1-x_n)^{\frac{1}{n}} - A_{n-1}^{\frac{1}{n}}\right)^k}{G'_{n-1}^{\frac{k-1}{n}} (1-x_n)^{\frac{1}{n}}} + \left(\frac{A'_{n-1}}{G'_{n-1}}\right)^{\frac{n-1}{n}}$$

with the induction hypothesis

$$\frac{A_{n-1}}{G_{n-1}} \geq \frac{A'_{n-1}}{G'_{n-1}},$$

and taking into account that

$$\begin{aligned} G_{n-1} &< G'_{n-1}, & A_{n-1} &< A'_{n-1}, & x_n &\leq 1 - x_n, \\ A'_{n-1} &= 1 - A_{n-1}, & x_n &> A_{n-1}, \end{aligned}$$

and

$$\begin{aligned} x_n^{\frac{1}{n}} - A_{n-1}^{\frac{1}{n}} &= \frac{x_n - A_{n-1}}{\sum_{i=0}^{n-1} x_n^{\frac{i}{n}} A_{n-1}^{\frac{n-1-i}{n}}} > \frac{x_n - A_{n-1}}{\sum_{i=0}^{n-1} (1-x_n)^{\frac{i}{n}} A'_{n-1}^{\frac{n-1-i}{n}}} \\ &= |(1-x_n)^{\frac{1}{n}} - A_{n-1}^{\frac{1}{n}}|, \end{aligned}$$

we get  $\frac{A_n}{G_n} > \frac{A'_n}{G'_n}$  and the proof is completed.  $\square$

---

**References**

- [1] Alzer, H.: The inequality of Ky Fan and related results. *Acta Appl. Math.* 38 (1995), 305–354.
- [2] Beckenbach, E.F.; Bellman, R.: *Inequalities*. Springer-Verlag, Berlin 1961.
- [3] Rooin, J.: AGM inequality with binomial expansion. *Elem. Math.* 58 (2003), 115–117.

Jamal Rooin  
Department of Mathematics  
Institute for Advanced  
Studies in Basic Sciences  
Zanjan, Iran  
e-mail: rooin@iasbs.ac.ir