

**Zeitschrift:** Elemente der Mathematik  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 74 (2019)  
**Heft:** 1

**Artikel:** Logarithm and dilogarithm  
**Autor:** Kramer, Jürg / Pippich, Anna-Maria von  
**DOI:** <https://doi.org/10.5169/seals-869228>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 24.03.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

---

---

## Logarithm and dilogarithm

---

---

Jürg Kramer and Anna-Maria von Pippich

Jürg Kramer ist Professor für Mathematik an der Humboldt-Universität zu Berlin. Sein Hauptforschungsgebiet ist die arithmetische Geometrie und die Theorie der Modulformen. Darüberhinaus ist er an Fragen zur Mathematikausbildung von Studierenden, Doktoranden und Lehrkräften interessiert.

Anna-Maria von Pippich ist Juniorprofessorin für Algebra und Zahlentheorie an der Technischen Universität Darmstadt. Ihr Hauptforschungsgebiet ist die Theorie der automorphen Formen, insbesondere in Verbindung mit zahlentheoretischen Fragestellungen.

### 1 The logarithm

**1.1. A naive sequence.** Following D. Zagier, we begin with the sequence of non-zero complex numbers determined by the requirement that each element of the sequence is given as one *divided* by the product of the left-hand and right-hand neighbors of the element in question. Denoting our sequence by  $a_1, a_2, a_3, a_4, a_5, \dots$ , the defining properties of the sequence give

$$a_2 = \frac{1}{a_1 a_3},$$

Im nachfolgenden Beitrag erinnern die Autoren zunächst an die geometrische Deutung der Funktionalgleichung der klassischen Logarithmusfunktion in Termen der hyperbolischen Distanz zwischen zwei Punkten der hyperbolischen Ebene. Darauf aufbauend wird in Folge der Dilogarithmus eingeführt und dessen Werte als hyperbolische Volumina idealer hyperbolischer Tetraeder im hyperbolischen 3-Raum erkannt. Es zeigt sich dabei, dass die Funktionalgleichung des Dilogarithmus der Zerlegung gewisser hyperbolischer Körper in solche ideale hyperbolische Tetraeder entspricht. Im letzten Teil werden die bis dahin gewonnenen geometrischen Erkenntnisse zur Berechnung spezieller Werte der Dedekindschen Zetafunktion von Zahlkörpern mit Hilfe von Logarithmus und Dilogarithmus genutzt. Dies führt zu einer bis heute ungelösten Vermutung von D. Zagier zur Beschreibung der Werte Dedekindscher Zetafunktionen an den ganzzahligen Stellen.

hence the first three members of the sequence are  $a_1, a_2, (a_1 a_2)^{-1}$ . The next member of the sequence is now determined as

$$a_3 = \frac{1}{a_2 a_4} \iff a_4 = \frac{1}{a_2 a_3} = \frac{1}{a_2 (a_1 a_2)^{-1}} = a_1.$$

For the fifth member we thus compute

$$a_4 = \frac{1}{a_3 a_5} \iff a_5 = \frac{1}{a_3 a_4} = \frac{1}{(a_1 a_2)^{-1} a_1} = a_2.$$

From this we conclude that the above requirement leads to the periodic sequence

$$a_1, a_2, \frac{1}{a_1 a_2}, a_1, a_2, \frac{1}{a_1 a_2}, \dots \quad (1)$$

with period 3; here  $a_1, a_2$  are non-zero complex numbers.

**1.2. The logarithm.** Based on the sequence (1), we now ask for smooth, complex-valued functions  $f$  on  $\mathbb{C}^\times := \mathbb{C} \setminus \{0\}$ , which are characterized by the property that the sum of the values of  $f$  at any three consecutive elements of the sequence (1) is equal to zero, i.e., the function  $f$  satisfies the functional equation

$$f(a_1) + f(a_2) + f\left(\frac{1}{a_1 a_2}\right) = 0. \quad (2)$$

From complex analysis we know that the principal logarithm  $\text{Log}(z)$  ( $z \in \mathbb{C} \setminus \mathbb{R}_{\leq 0}$ ) with the power series expansion

$$\text{Log}(z) = - \sum_{n=1}^{\infty} \frac{(1-z)^n}{n} \quad (z \in \mathbb{C} : |z-1| < 1) \quad (3)$$

solves the functional equation (2) up to integral multiples of  $2\pi i$  (see, e.g., [4]). By replacing the principal logarithm by its real part, i.e., by

$$\text{Re}(\text{Log}(z)) = \log(|z|),$$

we obtain a solution of the functional equation (2), which is unique up to scaling. By the normalization condition  $f(e) = 1$ , the solution becomes unique.

**1.3. Geometric interpretation.** We now give a geometric interpretation of the results derived in the preceding two sections. The underlying geometry is given by the hyperbolic geometry of the hyperbolic plane (for which we refer to [2]) in the model of the upper half-plane

$$\mathbb{H}^2 := \{z \in \mathbb{C} \mid z = x + iy, y > 0\}$$

equipped with the metric

$$ds_{\text{hyp}} := \frac{|dz|}{y}.$$

The isometry group of  $\mathbb{H}^2$  equipped with the metric  $ds_{\text{hyp}}$  is given by the special projective linear group  $\text{PSL}_2(\mathbb{R})$  acting by fractional linear transformations on  $\mathbb{H}^2$  as follows: For  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2(\mathbb{R})$  and  $z \in \mathbb{H}^2$ , the image  $\gamma(z) \in \mathbb{H}^2$  is given by

$$\gamma(z) = \frac{az + b}{cz + d}.$$

The hyperbolic distance between two points  $z_1, z_2 \in \mathbb{H}^2$  is defined as

$$\text{dist}_{\text{hyp}}(z_1, z_2) := \inf_{\eta} \int_{\eta} ds_{\text{hyp}}, \quad (4)$$

where the infimum is taken over all continuously differentiable paths  $\eta: [0, 1] \rightarrow \mathbb{H}^2$  satisfying  $\eta(0) = z_1$  and  $\eta(1) = z_2$ . In order to give an explicit formula for the hyperbolic distance (4), one shows in a first step that the hyperbolic distance from the point  $i\lambda_1$  to  $i\lambda_2$  with  $\lambda_1, \lambda_2 \in \mathbb{R}$  and  $\lambda_2 \geq \lambda_1$  is given by  $\log(\lambda_2/\lambda_1)$ . In fact, this is easily seen: Given a path  $\eta(t) = x(t) + iy(t)$  ( $t \in [0, 1]$ ,  $\eta(0) = i\lambda_1$ ,  $\eta(1) = i\lambda_2$ ), we have

$$\begin{aligned} \text{dist}_{\text{hyp}}(i\lambda_1, i\lambda_2) &= \inf_{\eta} \int_0^1 \frac{\sqrt{x'(t)^2 + y'(t)^2}}{y(t)} dt = \int_0^1 \frac{|y'(t)|}{y(t)} dt \\ &= \left[ \log(y(t)) \right]_0^1 = \log\left(\frac{\lambda_2}{\lambda_1}\right), \end{aligned}$$

since the infimum, in fact the minimum, is achieved for the path satisfying  $x'(t) = 0$  for all  $t \in [0, 1]$ , i.e., for the path  $\eta(t) = iy(t)$ . In a second step one then proves the existence of an element  $\gamma \in \text{PSL}_2(\mathbb{R})$  with the property  $\gamma(z_1) = i\lambda_1$  and  $\gamma(z_2) = i\lambda_2$ , where  $\lambda_1, \lambda_2 \in \mathbb{R}$  with  $\lambda_2 \geq \lambda_1$  satisfy

$$\frac{\lambda_2}{\lambda_1} = \frac{1 + \left| \frac{z_1 - z_2}{z_1 - \bar{z}_2} \right|}{1 - \left| \frac{z_1 - z_2}{z_1 - \bar{z}_2} \right|}. \quad (5)$$

Using the invariance of the hyperbolic distance by the action of  $\text{PSL}_2(\mathbb{R})$ , we find that the hyperbolic distance between  $z_1$  and  $z_2$  equals the *logarithm* of the ratio  $\lambda_2/\lambda_1$  given by formula (5), i.e., we have the relation

$$\text{dist}_{\text{hyp}}(z_1, z_2) = \log(\lambda_2/\lambda_1).$$

After having “rediscovered” the logarithm as the hyperbolic distance between two points in the upper half-plane  $\mathbb{H}^2$ , we next provide a hyperbolic geometric interpretation characterizing the naive sequence (1), which underlies the functional equation of the logarithm (see Figure 1).

To this end, we recall that the boundary of the upper half-plane  $\mathbb{H}^2$  is given by the real line  $\mathbb{R}$ , the  $x$ -axis, together with the point  $i\infty$  at infinity, i.e., the real projective line  $\mathbb{P}^1(\mathbb{R})$ . Furthermore, the geodesics in the upper half-plane are given by Euclidean semicircles in




Figure 1

$\mathbb{H}^2$ , which are orthogonal to the  $x$ -axis, or straight lines in  $\mathbb{H}^2$ , which are parallel to the  $y$ -axis. Therefore, the unique geodesic passing through  $z_1$  and  $z_2$  is given by the semicircle  $\mathcal{L}$  from  $z'_1$  to  $z'_2$  as depicted in the preceding figure. Using an appropriate  $\gamma \in \text{PSL}_2(\mathbb{R})$ , the geodesic  $\mathcal{L}$  can be mapped onto the geodesic from  $0$  to  $i\infty$  such that  $\gamma(z_1) = i\lambda_1$  and  $\gamma(z_2) = i\lambda_2$ .

Let us now connect our geometric insight to the naive sequence (1). For this we give ourselves three complex numbers  $z_1, z_2, z_3 \in \mathbb{H}^2$  lying on a hyperbolic line  $\mathcal{L}$  as depicted in Figure 2.




Figure 2

Denoting by  $[z_j, z_k]$  ( $j, k = 1, 2, 3$  with  $j \neq k$ ) the closed segment of  $\mathcal{L}$  consisting of all the points of  $\mathcal{L}$  lying between  $z_j$  and  $z_k$ , we may assume without loss of generality that  $z_2 \in [z_1, z_3]$  as shown in the above figure. By means of formula (5) we are now able to

determine real numbers

$$\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R} \quad \text{satisfying} \quad \lambda_1 \leq \lambda_2 \leq \lambda_3$$

such that  $\gamma(z_j) = i\lambda_j$  ( $j = 1, 2, 3$ ) for a suitable  $\gamma \in \text{PSL}_2(\mathbb{R})$ . Now, we introduce the positive real numbers  $a_j := \lambda_{j+1}/\lambda_j$  ( $j = 1, 2, 3$ ), where the indices are to be read modulo 3, that is,

$$a_1 = \frac{\lambda_2}{\lambda_1}, \quad a_2 = \frac{\lambda_3}{\lambda_2}, \quad a_3 = \frac{\lambda_1}{\lambda_3}.$$

Since we obviously have  $a_3 = (a_1 a_2)^{-1}$ , the sequence  $a_1, a_2, a_3, \dots$  gives rise to the naive sequence (1). The functional equation (2), i.e.,

$$\log(a_1) + \log(a_2) + \log(a_3) = 0 \quad \iff \quad \log\left(\frac{\lambda_2}{\lambda_1}\right) + \log\left(\frac{\lambda_3}{\lambda_2}\right) = \log\left(\frac{\lambda_3}{\lambda_1}\right)$$

now becomes equivalent to the obvious additive relation among the hyperbolic distances between the points  $z_1, z_2$ , and  $z_3$  lying on the geodesic  $\mathcal{L}$ , namely

$$\text{dist}_{\text{hyp}}(z_1, z_2) + \text{dist}_{\text{hyp}}(z_2, z_3) = \text{dist}_{\text{hyp}}(z_1, z_3).$$

## 2 The dilogarithm

**2.1. A more sophisticated sequence.** Next, we slightly vary the construction of the sequence studied in Subsection 1.1 by switching from the multiplicative to the additive viewpoint. More specifically, again following D. Zagier, we now ask for a sequence of non-zero complex numbers characterized by the property that each element of the sequence is given as one *minus* the product of the left-hand and right-hand neighbors of the element in question. Denoting our sequence by  $a_1, a_2, a_3, a_4, a_5, a_6, a_7, \dots$ , the defining properties of the sequence give

$$a_2 = 1 - a_1 a_3 \quad \iff \quad a_3 = \frac{1 - a_2}{a_1},$$

hence the first three members of the sequence are  $a_1, a_2, (1 - a_2)/a_1$ . The next member of the sequence is now determined as

$$a_3 = 1 - a_2 a_4 \quad \iff \quad a_4 = \frac{1 - a_3}{a_2} = \frac{a_1 + a_2 - 1}{a_1 a_2}.$$

For the fifth member we thus compute

$$a_4 = 1 - a_3 a_5 \quad \iff \quad a_5 = \frac{1 - a_4}{a_3} = \frac{1 - \frac{a_1 + a_2 - 1}{a_1 a_2}}{\frac{1 - a_2}{a_1}} = \frac{1 - a_1}{a_2}.$$

Turning to the sixth member of the sequence, we compute

$$a_5 = 1 - a_4 a_6 \iff a_6 = \frac{1 - a_5}{a_4} = \frac{1 - \frac{1-a_1}{a_2}}{\frac{a_1+a_2-1}{a_1 a_2}} = a_1.$$

It is now not a surprise that the seventh member in sequence is given by  $a_2$ , namely we have

$$a_6 = 1 - a_5 a_7 \iff a_7 = \frac{1 - a_6}{a_5} = \frac{1 - a_1}{\frac{1-a_1}{a_2}} = a_2.$$

From this we conclude that the above requirement leads to the periodic sequence

$$a_1, a_2, \frac{1 - a_2}{a_1}, \frac{a_1 + a_2 - 1}{a_1 a_2}, \frac{1 - a_1}{a_2}, a_1, a_2, \frac{1 - a_2}{a_1}, \frac{a_1 + a_2 - 1}{a_1 a_2}, \frac{1 - a_1}{a_2}, \dots \quad (6)$$

with period 5; here  $a_1, a_2$  are non-zero complex numbers.

**2.2. The dilogarithm.** In analogy to the situation of the first section relating the sequence (1) to the logarithm, we now ask for smooth, complex-valued functions  $f$  on  $\mathbb{C}^\times$ , which are characterized by the property that the sum of the values of  $f$  at any five consecutive elements of the sequence (6) is equal to zero, i.e., the function  $f$  satisfies the functional equation

$$f(a_1) + f(a_2) + f\left(\frac{1 - a_2}{a_1}\right) + f\left(\frac{a_1 + a_2 - 1}{a_1 a_2}\right) + f\left(\frac{1 - a_1}{a_2}\right) = 0. \quad (7)$$

Inspired by the solution in the case of the previous section, namely the power series expansion (3) with  $z$  replaced by  $1 - z$ , i.e.,

$$-\text{Log}(1 - z) = \sum_{n=1}^{\infty} \frac{z^n}{n} \quad (z \in \mathbb{C} : |z| < 1),$$

we now consider the dilogarithm  $\text{Li}_2(z)$  defined by the power series expansion

$$\text{Li}_2(z) := \sum_{n=1}^{\infty} \frac{z^n}{n^2} \quad (z \in \mathbb{C} : |z| < 1); \quad (8)$$

the definition of  $\text{Li}_2(z)$  on the whole complex plane then follows by analytic continuation. As in the case of the logarithm, this function is multi-valued and it turns out not to satisfy the functional equation (7). However, a suitable combination of the dilogarithm  $\text{Li}_2(z)$  and the solution of the functional equation (2), i.e.,  $\log(|z|)$ , leads to a positive answer of our question. Consider the so-called Bloch–Wigner dilogarithm  $D(z)$ , defined as

$$D(z) := \text{Im}(\text{Li}_2(z)) + \arg(1 - z) \log(|z|)$$

for  $z \in \mathbb{C} \setminus \mathbb{R}$ ; for  $z \in \mathbb{R}$ , we put  $D(z) = 0$ . It can then be shown that the Bloch–Wigner dilogarithm  $D(z)$  provides a solution of the functional equation

$$D(a_1) + D(a_2) + D(a_3) + D(a_4) + D(a_5) = 0 \quad (9)$$

for the generating members  $a_1, a_2, a_3, a_4, a_5$  of the 5-periodic sequence (6), which is unique up to scaling. For later purposes, we note that the Bloch–Wigner dilogarithm  $D(z)$  satisfies several other functional equations, such as

$$D\left(\frac{1}{z}\right) = -D(z), \quad D\left(1 - \frac{1}{z}\right) = D(z), \quad (10)$$

which are formal consequences of (9). For more details, we refer the reader to [9], pp. 8–11.

**2.3. Geometric interpretation.** In analogy to Subsection 1.3, we now give a geometric interpretation of the results derived in the preceding two sections. The underlying geometry is now given by the hyperbolic geometry of the hyperbolic 3-space (for which we refer to [3]) in the model of the upper half-space

$$\mathbb{H}^3 := \{(z, w) \in \mathbb{C} \times \mathbb{R} \mid z = x + iy, w > 0\}$$

equipped with the metric

$$ds_{\text{hyp}} := \frac{\sqrt{dx^2 + dy^2 + dw^2}}{w}. \quad (11)$$

The line element (11) gives rise to the volume element

$$d\mu_{\text{hyp}} := \frac{dx \wedge dy \wedge dw}{w^3},$$

which allows to compute volumes with respect to the hyperbolic metric.

The isometry group of  $\mathbb{H}^3$  equipped with the metric  $ds_{\text{hyp}}$  is given by the special projective linear group  $\text{PSL}_2(\mathbb{C})$  acting by fractional linear transformations on  $\mathbb{H}^3$  as follows: For  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2(\mathbb{C})$  and  $(z, w) \in \mathbb{H}^3$ , the image  $\gamma(z, w) \in \mathbb{H}^3$  is given by

$$\gamma(z, w) = \left( \frac{(az + b)(\bar{c}\bar{z} + \bar{d}) + a\bar{c}w^2}{|cz + d|^2 + |c|^2w^2}, \frac{w}{|cz + d|^2 + |c|^2w^2} \right).$$

In analogy to the 1-dimensional case, we will now see that the Bloch–Wigner dilogarithm  $D(z)$  computes the hyperbolic volumes of certain bodies in  $\mathbb{H}^3$ .

To this end, we recall that the boundary of the upper half-space  $\mathbb{H}^3$  consists of the complex plane  $\mathbb{C}$  given by  $\{(z, w) \in \mathbb{H}^3 \mid w = 0\}$  and the point  $\infty$  at infinity, i.e., it is given by the complex projective line  $\mathbb{P}^1(\mathbb{C})$ ; to simplify our notation, we will write  $z$  instead of  $(z, 0)$  for points on the boundary  $\mathbb{C} \subset \mathbb{P}^1(\mathbb{C})$ . Furthermore, the geodesics in the upper half-space are given by Euclidean (possibly degenerate) semicircles in  $\mathbb{H}^3$ , which are orthogonal to the boundary, as depicted in Figure 3.






Figure 3

An ideal hyperbolic tetrahedron is a geodesic 3-simplex all whose vertices  $z_1, z_2, z_3, z_4$  lie on the boundary  $\mathbb{P}^1(\mathbb{C})$  of  $\mathbb{H}^3$ . We denote it by  $T(z_1, z_2, z_3, z_4)$ , which keeps track of the ordering of the vertices and thus of the underlying orientation of the ideal hyperbolic tetrahedron. We note that for a given  $\gamma \in \text{PSL}_2(\mathbb{C})$ , the 3-simplex

$$T(\gamma(z_1), \gamma(z_2), \gamma(z_3), \gamma(z_4))$$

is again an ideal hyperbolic tetrahedron of the same hyperbolic volume as  $T(z_1, z_2, z_3, z_4)$ . Introducing the cross-ratio

$$[z_1 : z_2 : z_3 : z_4] := \frac{(z_1 - z_3)(z_2 - z_4)}{(z_1 - z_4)(z_2 - z_3)} \in \mathbb{P}^1(\mathbb{C}),$$

and choosing  $\gamma \in \text{PSL}_2(\mathbb{C})$  such that

$$\gamma(z_1) = \infty, \gamma(z_2) = 0, \gamma(z_3) = 1,$$

a short calculation yields

$$T(\gamma(z_1), \gamma(z_2), \gamma(z_3), \gamma(z_4)) = T(\infty, 0, 1, [z_1 : z_2 : z_3 : z_4]).$$

This shows that in order to compute the hyperbolic volume of an ideal hyperbolic tetrahedron  $T(z_1, z_2, z_3, z_4)$ , one is reduced to calculate the hyperbolic volume of

$$T(\infty, 0, 1, [z_1 : z_2 : z_3 : z_4]),$$

which looks as shown in Figure 4.

As predicted by the preceding consideration, the hyperbolic volume of  $T(z_1, z_2, z_3, z_4)$  depends only on the cross-ratio  $[z_1 : z_2 : z_3 : z_4]$ . A theorem going back to Lobachevsky now states that this volume equals the *Bloch–Wigner dilogarithm* of the corresponding cross-ratio, i.e., we have the relation

$$\text{vol}_{\text{hyp}}(T(z_1, z_2, z_3, z_4)) = D([z_1 : z_2 : z_3 : z_4]),$$




Figure 4

taking into account the orientation, namely, the volume is positive if the orientation is and negative otherwise. From this and the definition of the dilogarithm we see that the ideal hyperbolic tetrahedron  $T(z_1, z_2, z_3, z_4)$  is flat if and only if the cross-ratio is real. Furthermore, we see that the orientation of  $T(z_1, z_2, z_3, z_4)$  given by the ordering of its vertices agrees with the orientation induced from  $\mathbb{H}^3$  if and only if the cross-ratio has positive imaginary part.

After having “rediscovered” the Bloch–Wigner dilogarithm as the hyperbolic volume of an ideal hyperbolic tetrahedron in the upper half-space  $\mathbb{H}^3$ , we next provide a hyperbolic geometric interpretation characterizing the naive sequence (6), which underlies the functional equation of the Bloch–Wigner dilogarithm.

For this we give ourselves five complex numbers  $z_1, z_2, z_3, z_4, z_5$  lying on the boundary  $\mathbb{P}^1(\mathbb{C})$  of  $\mathbb{H}^3$ . We then introduce the complex numbers  $b_j := [z_1 : \dots : \hat{z}_j : \dots : z_5]$  ( $j = 1, \dots, 5$ ), where the entry marked with the hat has to be omitted, i.e., we consider the cross-ratios

$$\begin{aligned} b_1 &= [z_2 : z_3 : z_4 : z_5], & b_2 &= [z_1 : z_3 : z_4 : z_5], & b_3 &= [z_1 : z_2 : z_4 : z_5], \\ b_4 &= [z_1 : z_2 : z_3 : z_5], & b_5 &= [z_1 : z_2 : z_3 : z_4]. \end{aligned}$$

Now, letting

$$a_1 := \frac{1}{b_1}, \quad a_2 := 1 - \frac{1}{b_2}, \quad a_3 := \frac{1}{b_3}, \quad a_4 := 1 - \frac{1}{b_4}, \quad a_5 := \frac{1}{b_5}, \quad (12)$$

and reading indices modulo 5, it can easily be shown that the recursion formula

$$a_j = 1 - a_{j-1}a_{j+1} \quad (j = 1, \dots, 5)$$

holds. For example, if  $j = 2$ , one has to show that  $a_2 = 1 - a_1a_3$ , which, using (12), amounts to show that  $b_2 = b_1b_3$ . The latter equality can be immediately verified, since we

have

$$\begin{aligned}
b_1 \cdot b_3 &= [z_2 : z_3 : z_4 : z_5] \cdot [z_1 : z_2 : z_4 : z_5] \\
&= \frac{(z_2 - z_4)(z_3 - z_5)}{(z_2 - z_5)(z_3 - z_4)} \cdot \frac{(z_1 - z_4)(z_2 - z_5)}{(z_1 - z_5)(z_2 - z_4)} \\
&= \frac{(z_1 - z_4)(z_3 - z_5)}{(z_1 - z_5)(z_3 - z_4)} = [z_1 : z_3 : z_4 : z_5] = b_2.
\end{aligned}$$

Consequently, the sequence  $a_1, a_2, a_3, a_4, a_5$  above gives rise to the 5-periodic sequence (6).

After these preliminary computations, we now find using the functional equations (10) that

$$\begin{aligned}
&D(b_1) - D(b_2) + D(b_3) - D(b_4) + D(b_5) \\
&= -D\left(\frac{1}{b_1}\right) - D\left(1 - \frac{1}{b_2}\right) - D\left(\frac{1}{b_3}\right) - D\left(1 - \frac{1}{b_4}\right) - D\left(\frac{1}{b_5}\right) \\
&= -D(a_1) - D(a_2) - D(a_3) - D(a_4) - D(a_5) = 0,
\end{aligned}$$

taking into account the functional equation (9). In other words, we have proven that the functional equation (9) is equivalent to the following additive relation among the hyperbolic volumes of the ideal hyperbolic tetrahedra  $T(z_1, \dots, \hat{z}_j, \dots, z_5)$  ( $j = 1, \dots, 5$ ), namely

$$\sum_{j=1}^5 (-1)^j \text{vol}_{\text{hyp}}(T(z_1, \dots, \hat{z}_j, \dots, z_5)) = 0. \quad (13)$$

The relation (13) can also be checked directly geometrically by taking five complex numbers  $z_1, z_2, z_3, z_4, z_5$  lying on the boundary  $\mathbb{P}^1(\mathbb{C})$  of  $\mathbb{H}^3$  and then verifying that the alternating sum of the hyperbolic volumes of the five ideal hyperbolic tetrahedra

$$T(z_1, \dots, \hat{z}_j, \dots, z_5)$$

vanishes. This fact can be *seen* quite easily, if we choose

$$z_1 = 0, z_2 = z', z_3 = 1, z_4 = \infty, z_5 = z,$$

where we can pick  $z'$  without loss of generality in such a way that it lies inside of the Euclidean triangle determined by  $z_1, z_3, z_5$ . We then find that the union of the two ideal hyperbolic tetrahedra

$$T(z_1, \hat{z}_2, z_3, z_4, z_5), T(z_1, z_2, z_3, \hat{z}_4, z_5)$$

equals the union of the three ideal hyperbolic tetrahedra

$$T(\hat{z}_1, z_2, z_3, z_4, z_5), T(z_1, z_2, \hat{z}_3, z_4, z_5), T(z_1, z_2, z_3, z_4, \hat{z}_5),$$

from which the claimed relation immediately follows.

### 3 Special values of zeta functions

In this section we illustrate how the logarithm and dilogarithm together with their geometric interpretations presented in the previous sections as well as their generalizations, called polylogarithms, can be used to compute special values of zeta functions attached to number fields. However, in general, one is still far from having complete results and thus these special values are to be viewed as mysterious quantities.

For this section, we assume that the reader has some basic knowledge of algebraic number theory as one can find it for example in the book [6] of S. Lang.

**3.1. Zeta functions of number fields.** The Riemann zeta function  $\zeta(s)$  is defined by the series

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s},$$

where  $s \in \mathbb{C}$  with  $\operatorname{Re}(s) > 1$ . This function is known to have a meromorphic continuation to the whole complex  $s$ -plane with a simple pole at  $s = 1$  with residue 1, which can be restated as

$$\operatorname{res}_{s=1} \zeta(s) = \lim_{s \rightarrow 1} (s - 1)\zeta(s) = 1.$$

The Riemann zeta function encodes basic arithmetic properties of the integers such as the fundamental theorem of arithmetic by means of its Euler product expansion and the infinitude of the number of primes through its pole at  $s = 1$ .

The special values of  $\zeta(s)$  at positive even integers have been computed already by L. Euler as

$$\zeta(2m) = \frac{(-1)^{m-1} (2\pi)^{2m} B_{2m}}{2(2m)!} \quad (m \in \mathbb{N}_{>0}),$$

where the  $B_{2m}$ 's are the so-called Bernoulli numbers, which are rational numbers. In other words, the special values  $\zeta(2m)$  are rational multiples of even powers of  $\pi$ , i.e.,  $\zeta(2m) \in \mathbb{Q}[\pi^2]$ . In contrast to the special values of  $\zeta(s)$  at positive even integers, their values at positive odd integers are still quite mysterious. For example, it was only in the 80s of the last century that R. Apéry was able to prove that the special value  $\zeta(3)$  is irrational (see [1]). Although alternative proofs of this result have meanwhile been found, none of them seems to generalize to other odd zeta values.

More generally, let now  $F/\mathbb{Q}$  be a number field. The Dedekind zeta function  $\zeta_F(s)$  attached to  $F$  is defined as

$$\zeta_F(s) := \sum_{\mathfrak{a} \subseteq \mathcal{O}_F} \frac{1}{N(\mathfrak{a})^s},$$

where again  $s \in \mathbb{C}$  with  $\operatorname{Re}(s) > 1$  and the sum is taken over all non-trivial integral ideals  $\mathfrak{a}$  contained in the ring of integers  $\mathcal{O}_F$  of  $F$ ; here,  $N(\mathfrak{a})$  denotes the norm of  $\mathfrak{a}$ , which is given by the quantity  $|\mathcal{O}_F/\mathfrak{a}|$ . One easily checks that  $\zeta(s) = \zeta_{\mathbb{Q}}(s)$ . As in the case of

the Riemann zeta function it can be shown that the Dedekind zeta function  $\zeta_F(s)$  has a meromorphic continuation to the whole complex  $s$ -plane with a simple pole at  $s = 1$ . Also in this more general case, the Dedekind zeta function encodes important arithmetic information about the ring of integers  $O_F$ , such as the unique factorization of integral ideals into powers of prime ideals and the infinitude of the number of prime ideals. As will be explained below, it is expected that the logarithm, dilogarithm and, more generally, the polylogarithms will allow to “compute” special values of Dedekind zeta functions of number fields at positive integers.

**3.2. The logarithm and the residue of  $\zeta_F(s)$  at  $s = 1$ .** In this paragraph we will show how the logarithm and its geometric characterization by means of hyperbolic lengths of geodesics in the upper half-plane  $\mathbb{H}^2$  presented in the first section is encoded in the residue of the Dedekind zeta function  $\zeta_F(s)$  at  $s = 1$ . To illustrate this phenomenon, it is best to start with the case of a real quadratic extension  $F = \mathbb{Q}(\sqrt{D})$ , where  $D \in \mathbb{N}$  is a positive discriminant, i.e.,  $D \equiv 0$  or  $1 \pmod{4}$ . The group of units  $O_F^\times$  of the ring of integers  $O_F$  of  $F$  is generated, modulo sign, by the so-called fundamental unit  $\varepsilon_D > 1$ . By the basic theory of integral binary quadratic forms it can then be shown that there exists  $\gamma \in \mathrm{PSL}_2(\mathbb{R})$  such that

$$\delta := \gamma \begin{pmatrix} \varepsilon_D & 0 \\ 0 & 1/\varepsilon_D \end{pmatrix} \gamma^{-1} \in \mathrm{PSL}_2(\mathbb{Z}),$$

and we see that the hyperbolic distance between the point  $z := \gamma(i) \in \mathbb{H}^2$  and  $\delta(z) \in \mathbb{H}^2$  is given as

$$\mathrm{dist}_{\mathrm{hyp}}(z, \delta(z)) = \mathrm{dist}_{\mathrm{hyp}}(i, \varepsilon_D^2 i) = 2 \log(\varepsilon_D).$$

Projecting the geodesic line from  $z$  to  $\delta(z)$  to the quotient space  $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}^2$ , it thus turns out that we obtain a primitive closed geodesic of hyperbolic length  $2 \log(\varepsilon_D)$ . The detailed analysis in [7] in fact shows that *all* the primitive closed geodesics on  $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}^2$  are obtained in this way, where a fixed real quadratic extension  $F = \mathbb{Q}(\sqrt{D})$  together with its fundamental unit  $\varepsilon_D$  gives rise to exactly  $h_F$  different primitive closed geodesics, with  $h_F$  denoting the class number of  $F$ . The residue of  $\zeta_F(s)$  at  $s = 1$  now counts the lengths of all the  $h_F$  primitive closed geodesics weighted by the square root of the discriminant, i.e., we have the formula

$$\mathrm{res}_{s=1} \zeta_F(s) = \lim_{s \rightarrow 1} (s-1) \zeta_F(s) = \frac{2 h_F \log(\varepsilon_D)}{\sqrt{D}}. \quad (14)$$

Turning now to the general case, we let  $F/\mathbb{Q}$  be any number field with ring of integers  $O_F$  and discriminant  $D_F$ . It is known that  $F$  comes with  $r_1$  real embeddings and  $r_2$  pairs of complex embeddings, where  $[F : \mathbb{Q}] = r_1 + 2r_2$ . Dirichlet’s unit theorem now states that the group  $O_F^\times$  of units modulo the roots of unity in  $O_F^\times$  is a free abelian group generated by  $r := r_1 + r_2 - 1$  units  $\varepsilon_1, \dots, \varepsilon_r$ . Denoting the real and half of the complex embeddings by  $\sigma_1, \dots, \sigma_{r_1+r_2}$ , the residue of  $\zeta_F(s)$  at  $s = 1$  is finally given by the so-called class number formula, which reads

$$\mathrm{res}_{s=1} \zeta_F(s) = \lim_{s \rightarrow 1} (s-1) \zeta_F(s) = \frac{2^{r_1} (2\pi)^{r_2} h_F \det((\log |\sigma_j(\varepsilon_k)|^{\ell_j})_{j,k=1,\dots,r})}{\omega_F \sqrt{|D_F|}}, \quad (15)$$

where  $\omega_F$  is the number of roots of unity in  $O_F^\times$  and  $e_j = 1$ , if  $\sigma_j$  is a real embedding, and  $e_j = 2$  otherwise. One easily checks that formula (14) is a special case of the general class number formula (15), and we also recognize that the special value of  $\zeta_F(s)$ , or rather its residue, at  $s = 1$  can be expressed in terms of logarithms evaluated at distinguished elements of  $F$ .

**3.3. The dilogarithm and the value of  $\zeta_F(s)$  at  $s = 2$ .** In this paragraph we will show how the dilogarithm and its geometric characterization by means of hyperbolic volumes of ideal hyperbolic tetrahedra in the upper half-space  $\mathbb{H}^3$  presented in the second section is encoded in the value of the Dedekind zeta function  $\zeta_F(s)$  at  $s = 2$ . To illustrate this fact, it is best to start with the case of an imaginary quadratic extension  $F = \mathbb{Q}(\sqrt{-D})$ , where  $-D \in \mathbb{Z}$  is a negative discriminant, i.e.,  $D \equiv 0$  or  $3 \pmod{4}$ . It is then known that the Bianchi group  $\mathrm{PSL}_2(O_F)$ , with  $O_F$  denoting again the ring of integers of  $F$ , acts strongly discontinuously on  $\mathbb{H}^3$  and the resulting quotient space  $\mathrm{PSL}_2(O_F) \backslash \mathbb{H}^3$  is an oriented hyperbolic 3-orbifold (i.e., a hyperbolic 3-manifold with the exception of finitely many singular points due to the torsion elements of  $\mathrm{PSL}_2(O_F)$ ) of finite volume. This hyperbolic volume has been computed by M.G. Humbert, who found the formula

$$\mathrm{vol}_{\mathrm{hyp}}(\mathrm{PSL}_2(O_F) \backslash \mathbb{H}^3) = \frac{D\sqrt{D}}{4\pi^2} \zeta_F(2).$$

Since, on the other hand, every hyperbolic 3-manifold or 3-orbifold can be triangulated into ideal hyperbolic tetrahedra, the volume  $\mathrm{vol}_{\mathrm{hyp}}(\mathrm{PSL}_2(O_F) \backslash \mathbb{H}^3)$  can also be expressed in terms of special values of dilogarithms. For example, for  $D = 3$ , one is thus led to the formula

$$\zeta_{\mathbb{Q}(\sqrt{-3})}(2) = \frac{2}{27} \pi^2 \sqrt{3} D \left( \frac{1 + \sqrt{-3}}{2} \right) \underset{\mathbb{Q}^\times}{\sim} \pi^2 \sqrt{3} D \left( \frac{1 + \sqrt{-3}}{2} \right),$$

where the latter notation means that equality holds up to a non-zero rational number. Turning now to the general case, where  $F/\mathbb{Q}$  is any number field with ring of integers  $O_F$  and discriminant  $D_F$ , and letting  $\sigma_1, \dots, \sigma_{r_2}$  denote the  $r_2$  complex embeddings, the value of  $\zeta_F(s)$  at  $s = 2$  is given as

$$\zeta_F(2) \underset{\mathbb{Q}^\times}{\sim} \pi^{2(r_1+r_2)} \sqrt{|D_F|} \det(D(\sigma_j(\xi_k)))_{j,k=1,\dots,r_2}, \quad (16)$$

where  $\xi_1, \dots, \xi_{r_2}$  are suitable elements of  $F$ . We thus recognize that the special value of  $\zeta_F(s)$  at  $s = 2$  can indeed be expressed in terms of dilogarithms evaluated at distinguished elements of  $F$ .

**3.4. Zagier's conjecture.** In order to state this conjecture about the values of the Dedekind zeta function  $\zeta_F(s)$  of number fields  $F$  at positive integers  $s = m$ , we introduce the  $m$ th polylogarithm by generalizing the series expansion (8) of the dilogarithm to

$$\mathrm{Li}_m(z) := \sum_{n=1}^{\infty} \frac{z^n}{n^m} \quad (z \in \mathbb{C} : |z| < 1),$$

and – in analogy to the Bloch–Wigner dilogarithm – the corrected function

$$\mathcal{L}_m(z) := \operatorname{Re}_m \left( \sum_{j=0}^{m-1} \frac{2^j B_j}{j!} \log(|z|)^j \operatorname{Li}_{m-j}(z) \right),$$

where  $\operatorname{Re}_m$  equals the real or the imaginary part, when  $m$  is odd or even, respectively, and  $B_j$  refer again to the Bernoulli numbers. One immediately verifies that

$$\mathcal{L}_1(z) = -\log(|1-z|) \quad \text{and} \quad \mathcal{L}_2(z) = D(z).$$

Letting  $n_+ := r_1 + r_2$  and  $n_- := r_2$ , as well as  $\pm = (-1)^{m-1}$  (and thus  $\mp = (-1)^m$ ), Zagier's conjecture (see [8]) then states that

$$\zeta_F(m) \underset{\mathbb{Q}^\times}{\sim} \pi^{mn_\mp} \sqrt{|D_F|} \det \left( \mathcal{L}_m(\sigma_j(\zeta_k))_{j,k=1,\dots,n_\pm} \right); \quad (17)$$

here  $D_F$  is the discriminant of  $F$ ,  $\sigma_j$  denote the  $r_1$  real and the  $r_2$  complex embeddings of  $F$ , and  $\zeta_1, \dots, \zeta_{n_\pm}$  are suitable elements of  $F$ . It is straightforward to check that Zagier's conjectured formula (17) coincides with formulas (15) and (16), if  $m = 1$  and  $m = 2$ , respectively. This conjecture has been proven for  $m = 3$  by A.B. Goncharov (see [5]). The general case remains open to this date and is the subject of current research.

## References

- [1] R. Apéry: *Irrationalité de  $\zeta(2)$  et  $\zeta(3)$* . Luminy Conference on Arithmetic. Astérisque **61** (1979), 11–13.
- [2] A.F. Beardon: *The geometry of discrete groups*. Graduate Texts in Mathematics **91**. Corrected second printing. Springer-Verlag, New York, 1995.
- [3] J. Elstrodt, F. Grunewald, J. Mennicke: *Groups acting on hyperbolic space*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [4] E. Freitag, R. Busam: *Complex analysis*. Universitext. Second edition. Springer-Verlag, Berlin, 2009.
- [5] A.B. Goncharov: *Geometry of configurations, polylogarithms, and motivic cohomology*. Adv. Math. **114** (1995), 197–318.
- [6] S. Lang: *Algebraic number theory*. Graduate Texts in Mathematics **110**. Second edition. Springer-Verlag, New-York, 1994.
- [7] P. Sarnak: *Class numbers of indefinite binary quadratic forms*. J. Number Th. **15** (1982), 229–247.
- [8] D. Zagier: *Polylogarithms, Dedekind zeta functions and the algebraic K-theory of fields*. In: Arithmetic Algebraic Geometry, Texel, 1989. Progress in Mathematics **89**, Birkhäuser Boston, 1991, pages 391–430.
- [9] D. Zagier: *The dilogarithm function*. In: Frontiers in Number Theory, Physics, and Geometry II. Springer-Verlag, Berlin, 2007, pages 3–65.

Jürg Kramer  
 Institut für Mathematik  
 Humboldt-Universität zu Berlin  
 Unter den Linden 6  
 D-10099 Berlin, Germany  
 e-mail:

kramer@math.hu-berlin.de

Anna-Maria von Pippich  
 Fachbereich Mathematik  
 Technische Universität Darmstadt  
 Schloßgartenstraße 7  
 D-64289 Darmstadt, Germany  
 e-mail:

pippich@mathematik.tu-darmstadt.de