

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 86 (1993)
Heft: 3

Artikel: Late Aptian-Early Albian radiolaria of the Windalia radiolarite (type section), Carnarvon Basin, Western Australia
Autor: Ellis, Glynn
Kapitel: 2: Material and methods
DOI: <https://doi.org/10.5169/seals-167268>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 26.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The age of the Windalia Radiolarite has been based previously on associated foraminifera and megafossils. Whitehouse (1926, 1927) suggested the Windalia to be equivalent in age to the Tambo Series (Albian) in Queensland based of the presence of similar belemnites. Raggatt (1936) considered the Winning Group to be Upper Albian and equivalent to the radiolarian rocks described by Hinde (1893) from the Northern Territory. Condon (1954) attributed a Late Cenomanian age for the Windalia Radiolarite based on foraminifera identified by Edgell (1952) and on the presence of large discoidal ammonites belonging to the Family Acanthoceratidae. Subsequent foraminiferal studies by Glaessner (1955) and Belford (1959; for the succeeding Gearle Siltstone), and reinterpreted ammonite identifications, which include the Family Aconoceratidae, by Brunnschweiler (1959) place the Windalia Radiolarite within the Late Aptian to Early Albian.

2. Material and methods

Assemblages examined for this study come from the type section of the Windalia Radiolarite at Windalia Hill (Lat. 23°16'S, Long. 114°48'E) on Winning Station (Fig. 1). Only the lower part of the formation is exposed, conformably overlying the Muderong Shale (2 samples of which are used in this study). A detailed lithostratigraphy of the type section is given below and in Fig. 3. A total of 25 samples were collected, of which 14 were analyzed for their radiolarian (and other fossil) content. Selection of samples for processing was based on observed freshness and friability. Sample preparation generally followed standard foraminiferal processing techniques. About 5 cm³ of sample was broken down to mm-sized pieces and then boiled with sodium pyrophosphate (Calgon™) and a small amount of 10% hydrogen peroxide. Successive drying and retreatments were necessary to assist disaggregation of samples with high clay contents. The sediment was then washed through 150 µm and 63 µm sieves to remove the clay. Final residues were dried and stored in plastic vials.

The 63–150 µm and 150 µm–2 mm size fractions were examined and the fossil residue (radiolaria, foraminifera, sponge spicules) placed on a counting tray and collected into grided cardboard slides. The characteristics of the radiolarian thanatoconoesis for each sample were first noted on a minimum representative count of 300 specimens. The entire residue from each sample was then searched several times for rare taxa and well-preserved forms for illustration. This ensured a better perception of the true distribution of specific radiolaria that may have been unevenly dispersed on the counting tray due to size and/or shape. Qualitative estimates of species abundance are tabulated in Table 1.

Preservation of the radiolaria ranges from very poor to good. Compaction of the sediment during lithification has resulted in many broken specimens, and preservation of the finer structures of some radiolaria is rare. With many specimens superficial clay still adhered to surfaces after washing the sediment, and it was necessary to use vigorous additional cleaning (e.g. ultrasound, reboiling and sieving) to aid determinations. Unfortunately, these techniques resulted in broken specimens. The radiolaria are illustrated by scanning electron micrographs taken with a PHILLIPS SEM 505 at the University of Western Australia and with a CAMSCAN Series 4 SEM at the Université de Lausanne. Specimens were mounted onto SEM plugs using double sided tape and then coated with carbon and gold in a vacuum evaporator. The amount of coating required to prevent “charging” varied with shell type and preservation. Generally, forms with spongy tests (e.g. Spongodiscacea) required a heavier coating than forms having latticed meshwork. Transmitted light identification of some radiolaria was carried out using immersion oil or by preparing strewn slides with molten “Lakeside Cement™”. Although these are quick techniques allowing viewing of internal structures, the slides are generally of poor quality and the resulting photographs are not suitable for publication.

Tab. 1. Distribution of radiolaria from the type section of the Windalia Radiolarite at Windalia Hill.

TAXA	WINDALIA RADIOLARITE															
	WIND 24	WIND 23	WIND 21	WIND 19	WIND 17	WIND 16	WIND 15	WIND 13	WIND 10	WIND 9	WIND 8	WIND 7	WIND 5	WIND 4	Muderong Shale	
<i>Acaeniotyle diaphorogona</i>	+	+														
<i>Acaeniotyle longispina</i>																
<i>Acaeniotyle</i> sp. cf. <i>A. diaphorogona</i>																
<i>Acaeniotyle</i> (?) sp. A																
<i>Acaeniotyle</i> (?) sp. B																
<i>Actinommid</i> gen. & sp. indet																
<i>Actinomma</i> (?) <i>pleiadesensis</i>																
<i>Alievium</i> (?) sp. A																
<i>Alievium</i> (?) sp. B																
<i>Amphipyndax stocki</i>																
<i>Angulobracchia crassa</i>																
<i>Arachnosphaera exilis</i>																
<i>Archaeocenosphaera euganea</i>																
<i>Archaeodictyomitra siliteri</i>																
<i>Archaeodictyomitra vulgaris</i>																
<i>Archaeospongoprunum carrierensis</i>																
<i>Archaeospongoprunum diversispina</i>																
<i>Archaeospongoprunum klingi</i>																
<i>Archaeospongoprunum</i> sp. cf. <i>A. tehaemaensis</i>																
<i>Archaeospongoprunum</i> sp. cf. <i>A. praelongum</i>																
<i>Archaeospongoprunum</i> sp.																
<i>Artocapsa ultima</i>																
<i>Crucella messinae</i>																
<i>Crucella</i> sp.																
<i>Cyrtocalpis operosa</i>																
<i>Dicanthocapsa</i> sp. cf. <i>D. ancus</i>																
<i>Gonglyothorax cephalocrypta</i>																
<i>Haliomma</i> sp.																
<i>Hemicyptocapsa</i> sp. cf. <i>H. simplex</i>																
<i>Histastrum aster</i>																
<i>Holocryptocanium barbui barbui</i>																
<i>Mesosaturminus hueyi</i> group																
<i>Mita</i> sp.																
<i>Napora dumitricai</i>																

Species abundance is defined as follows:

- A = abundant – more than 30 specimens per 300
- C = common – 15–29 specimens
- F = few – 3–14 specimens
- R = rare – 1–2 specimens
- + = very rare – fewer than 2 specimens per 1000
- ? = doubtful identification

