

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 95 (2002)
Heft: 3

Artikel: Brachiopods from the Plattenwald Bed (Albian, Cretaceous) of the Helvetic Alps of Vorarlberg (Austria)
Autor: Sulser, Heinz / Friebe, J. Georg
DOI: <https://doi.org/10.5169/seals-168968>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Brachiopods from the Plattenwald Bed (Albian, Cretaceous) of the Helvetic Alps of Vorarlberg (Austria)

HEINZ SULSER¹ & J. GEORG FRIEBE²

Key words: Brachiopods, new species, Plattenwald Bed, conditions of accumulation, Helvetic nappes, Albian, Cretaceous, Vorarlberg, Austria

ABSTRACT

The brachiopods of the Plattenwald Bed (Albian, Cretaceous) of diverse localities in the Helvetic nappes of Vorarlberg (Austria) are described. The most abundant faunal element are terebratulids (*Moutonithyris dutempleana*, showing a distinct dimorphism: A “obtusa type”; B “dutempleana type”), followed by rare rhynchonellids (*Orbirhynchia parkinsoni*, *Burrirhynchia tripartita*) and a new terebratulid, *Gemmarcula ? plattenwaldensis* sp. nov. Further brachiopods are very rare and have not yet been studied. The brachiopods are discussed with regard to the lithology and the depositional environment of the Plattenwald Bed and with regard to their possible provenance.

ZUSAMMENFASSUNG

Die Brachiopoden der Plattenwald-Schicht (Albian, Kreide) von verschiedenen Fundorten der Helvetischen Decken von Vorarlberg (Österreich) werden beschrieben. Die Fauna umfasst als häufigstes Element Terebratuliden (*Moutonithyris dutempleana* mit deutlichem Dimorphismus: A “obtusa-Typ”; B “dutempleana-Typ”), die wesentlich selteneren Rhynchonelliden *Orbirhynchia parkinsoni* und *Burrirhynchia tripartita* und ein neuer Terebratulide, *Gemmarcula ? plattenwaldensis* sp. nov. Weitere Brachiopoden sind sehr selten und noch nicht näher untersucht. Die Brachiopoden werden im Zusammenhang mit der Lithologie und den Ablagerungsbedingungen der Plattenwald-Schicht und ihrer möglichen Herkunft diskutiert.

1. Introduction

The knowledge of Early Cretaceous brachiopods in the Helvetic nappes mainly rests on ancient publications such as those of Ooster (1863), Bachmann (1864), Kaufmann (1877) and on generalized fossil lists from geological publications. In the Swiss and French Jura Mountains the knowledge of the corresponding brachiopods can be based on the pioneering work of Loriol (1864, 1868) and Pictet (1872), dating back to the 19th century, and revisions by Burri (1953, 1956), Gaspard (1989, 1997, 1999) and Middlemiss (1981, 1983, 1984, 1989).

Towards the end of the Early Cretaceous, during the final drowning phase of the latest Aptian, the widespread carbonate platforms („Schrattenkalk“ Formation) ceased to develop. Due to a fundamental change in the conditions of sedimentation an episode of very slow accumulation took place, resulting in condensed phosphatic horizons which are part of the

Garschella Formation (Gebhard 1985, Föllmi 1989a, Föllmi et al. 1994). In these rocks of limited extent (e.g. Durschlägi and Sellamatt Beds in Switzerland) brachiopods occur only sporadically. One of these phosphatic layers, however, the so-called Plattenwald Bed in the eastern Helvetic of Vorarlberg, has yielded not only a rich ammonite fauna, allowing biostratigraphical dating (Föllmi 1989b), but also sponges, endobenthic and epibenthic bivalves, gastropods, echinoderms, and a small, diverse brachiopod fauna. These brachiopods, listed in Heim & Seitz (1934), have hitherto never been figured or described. As they constitute an important episode in the history of the Alpine Cretaceous, we decided to study them in more detail in the present paper. An improved taxonomy may help to compare them with other coeval brachiopod communities and may contribute to paleogeographic questions.

¹ Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Str. 4, CH-8006 Zurich, Switzerland

² Vorarlberger Naturschau, Marktstrasse 33, A-6850 Dornbirn, Austria




Fig. 1. Location map of the Plattenwald Bed (Cretaceous, Albian) in Vorarlberg (Austria). P: Plattenwald; O: Orsanka; S: Sattelberg; Go: Gopfberg; R: Rotenbchalpe; Gu: Gütle; L: Lädtobel; A: Achbrücke.

2. Geological setting

Mainly specimens housed in the Vorarlberger Naturschau, Dornbirn (Austria) were used for this study. Most of them had been collected by the founder of the museum, Siegfried Fussenegger, and are listed in Heim & Seitz (1934). Some specimens originate from (formerly) private collections. The brachiopods were found at the following localities (Fig. 1):

Plattenwald (GKR coordinates -51405 / 242355), about 0.75 km NE of Klaus. A small outcrop described by Heim & Seitz (1934), which was already sampled by Fussenegger, was designed as stratotype of the „Plattenwald-Schicht“ by Föllmi & Ouwehand (1987). This phosphatic condensed bed lies above a dark, glauconitic sandstone („Brisi-Sandstein“) succeeded by 0.4 m glauconitic sandstone with abundant extraclasts, but rare macrofossils („Klauser Schichten“). The rich ammonite fauna of the Plattenwald Bed (Föllmi 1989b) contains predominantly *Leymeriella* spp. and *Douvilleiceras* ex gr. *mammillatum* (Schlotheim) (Lower Albian: tardefurcata- and mammillatum-zones). Younger ammonites are rare (up to inflatum-zone).

The rock is extremely weathered leaving phosphatic internal casts in a sandy matrix. An additional outcrop (GKR coordinates -51550 / 242300 / ± 40 m) can be found approximately 150 m west of the stratotype. A gentle slope is extremely burrowed by fossil collectors. Most ammonites (except small fragments) have been removed, but brachiopods (as well as bivalves) are still abundant.

Orsanka, about 1.25 km ENE of Klaus. East of the stratotype the Plattenwald Bed is exposed in several small outcrops (GKR coordinates of main outcrop: -50820 / 242435). Ammonites indicate the mammillatum-zone as well as the tardefurcata-zone by a recent find of *Leymeriella intermedia* Spath. One outcrop provided large specimens of *Moutonithyrus dutempleana* (d' Orbigny).

Sattelberg (GKR coordinates -52730 / 241880). The steeply dipping Plattenwald Bed is exposed on a slope about 1 km W of Klaus. It is overlain by pelagic limestone of the Seewen Formation („Seewerkalk“). As in the stratotype ammonites represent the tardefurcata- and mammillatum-zones, but also very rarely the dentatus-zone.

Gopfberg (GKR coordinates -32400 / 247270). This outcrop in the Brengenzlerwald area approximately 1.2 km SE of Reuthe was described by Heim & Seitz (1934) as a phosphatic bed overlying glauconitic sandstone („Brisi-Sandstein“). Besides some undetermined specimens of *Phylloceras*, *Leymeriella pseudoregularis* Seitz and *Douvilleiceras* ex gr. *mammillatum* (Schlotheim) are the only ammonites found in this outcrop.

Rotenbchalpe (GKR coordinates -37200 / 250150). No lithological description is given by Heim & Seitz (1934) of this location, approximately 2.9 km SSW of Schwarzenberg. Ammonites (determined by Seitz) indicate a Lower to early Middle Albian age.

Gütle – Kraftwerk Ebensand (GKR coordinates -41980 / 249000). The extremely fossiliferous bed was found near the entrance of the romantic Alploeh Gorge 4.75 km SE of Dornbirn. The strongly weathered outcrop, which was sampled by Siegfried Fussenegger, is not accessible any more. The Middle Albian genus *Hoplites* spp. is common (dentatus-zone). Occasional specimens of




Fig. 2. Piece of rock (75 mm high) of the Plattenwald Bed with *Moutonithyris dutempleana* (d'Orbigny) and fossil debris. Inventory number P. 16577.

Hysterocheras, *Dipoloceras* and *Mortoniceras* represent the early Upper Albian inflatum-zone.

Lädtofel near Ebnit (GKR coordinates -44625 / 246060). An outcrop of the Plattenwald Bed was found approximately 0.2 km N of the church of Ebnit during construction work for a forest road. The phosphatic bed overlies bioturbated glauconitic sandstone and is succeeded by pelagic limestone of the Seewen Formation. Again strongly weathered portions provided the best fossils. *Hoplites* spp. is common (dentatus-zone). *Anahoplites intermedius* Spath and *A. aff. praecox* (sensu Föllmi 1989b) are characteristic of the lower lorica-zone. One fragment of *Mortoniceras* sp. represents the inflatum-zone. However, *Douvilleceras* ex gr. *mammillatum* (Schlotheim) is still present (one fragment). Today the outcrop is overgrown by grass and thus no longer accessible.

Achbrücke near Schwarzenberg (GKR coordinates -36475 / 251470). This outcrop on the old road from Schwarzenberg to the former railway station (approximately 1.4 km S of the church of Schwarzenberg) was sampled by Siegfried Fussenegger during construction works following a rockfall event. The 0.7 m thick, phosphatic, fossiliferous layer (representing the Plattenwald Bed) directly overlies limestone of the Barremian to Lower Aptian Schratenkalk Formation. It is succeeded by massive, glauconitic, sandy limestone of the Aubrig Bed (Garschella Formation). This outcrop differs from all other locations by the occurrence of large specimens of *Mortoniceras* spp. and *Prohysterocheras* (*Goodhallites*) aff. *goodhalli* (J. Sowerby) (upper inflatum-zone). Other ammonites include *Puzosia* spp. and *Beudanticeras* spp.

Additional material was recovered from recently collected material by the authors at the following localities: Plattenwald, Sattelberg, and Orsanka.

3. Material and methods

As it was practically impossible to extract undestroyed brachiopods mechanically out of the hard rock, they had to be isolated by sieving from loose, weathered material, preserved as internal casts, sometimes with adhering fragments of the test (Fig. 2). The material is of poor quality, and the complete study of successive serial sections was not possible.

In favourable cases, parts of the cardinalia were preserved silicified and could be prepared by repeated treatment with 12% hydrochloric acid, followed by rinsing with water. By this procedure the hard matrix was scrubbed to sandy grains. In this way fragments of internal structures were isolated.

All figured brachiopods are deposited at the Vorarlberger Naturschau, Dornbirn (Austria).

4. Paleontology

Terebratulids predominate with about 90% of all brachiopods. Rhynchonellids and terebratellids count for about 10% of the total. Since all brachiopods are preserved as internal moulds the following descriptions are based thereupon.

Order	Rhynchonellida Kuhn 1949
Superfamily	Pugnacoidea Williams et al. 1996
Family	Pugnacidae Rzhonsnitskaya 1956
Subfamily	Lacunosellinae Smirnova 1963
Genus	<i>Orbirhynchia</i> Pettitt 1954 [type-species: <i>Orbirhynchia orbignyi</i> Pettitt 1954]

Orbirhynchia parkinsoni Owen 1960

(Fig. 3, 4)

pars *Terebratula sulcata* Parkinson 1811: auctt.

pars 1852–1854 *Rhynchonella sulcata* (Park.); Davidson: 85–87; pl. 10, fig. 18–20 (only)

1913 *Rhynchonella sulcata* (Park.); Jacob & Fallot: 66; pl. 9, fig. 14–17

1918 *Stolmorhynchia sulcata* (Park.); Buckman: 46

1925 *Rhynchonella deluci* Pict.; Schaad: 15

1925 *Rhynchonella sulcata* Park.; Schaad: 15

1934 *Rhynchonella* cf. *sulcata* (Park.); Heim & Seitz: 198, 229, 230

1960 *Orbirhynchia parkinsoni* Owen: 250–252; pl. 5, fig. 2a–c, textfig. 1

1965 *Rhynchonella sulcata* (Park. 1811) Davidson; Almérás: 384

1974 *Orbirhynchia parkinsoni* Owen; Dieni et al.: 210–211; pl. 38, fig. 10, 11

1977 *Orbirhynchia parkinsoni* Owen; Popiel-Barczyk: 32–35; pl. 2, fig. 1–6; textfig. 5–8

1980 *Orbirhynchia parkinsoni* f. *rencurelensis* (Jac. & Fall.); Middlemiss & Owen: 204; textfig. 8

1999 *Orbirhynchia parkinsoni* Owen; Sulser: 70 (with fig.)

For the completion of synonymy compare Owen (1960) and Popiel-Barczyk (1977).

Holotype in Owen (1960), pl. 5, fig. 2a–c from the Upper Albian (Cambridge Greensand) of Cambridge, England.

Occurrence and stratigraphy

Orbirhynchia parkinsoni is reported from the Albian of England (Cambridge, Isle of Wight), France (Dépt. Isère, Savoie), central Sardinia, western Switzerland (?), Spain, and from the Lower Cenomanian of Poland (Holy Cross Mountains, region of Cracow, Tatra).



Fig. 3. *Orbirhynchia parkinsoni* Owen. Specimen from Plattenwald. Dorsal, ventral, lateral, and anterior view. 2x natural size. Inventory number P. 6271.




Fig. 4. *Orbirhynchia parkinsoni* (Owen). Transverse serial sections of a specimen (length = 12.4, width = 12.1, thickness = 8.5 mm) from Klaus. Numbers give distance in mm from the ventral apex. 5 × natural size.

Description

External characters

Dimensions (mm) *	Length (L)	Width (W)	Thickness (T)	W : L	T : L
Majority (80%) of specimens	10.5–15.5	12.0–17.0	6.0–9.0	1.08–1.20	0.60–0.70
Extreme values	8.3/17.3	9.4/18.3	5.4/10.5	0.97/1.32	0.53/0.74

* Number of specimens : 21 (Plattenwald: 17; Sattelberg: 3; Achbrücke: 1)

Shell oval to subtrigonal in outline. Position of greatest width at half-length or somewhat more anteriorly. Shell biconvex, curvature of dorsal valve more pronounced. Dorsal valve with scarcely perceptible median fold, somewhat more marked anteriorly. Ventral valve with recognizable sinus near the anterior commissure, widening to a regularly formed, broad unipli-cation, 2–4 mm high. Shell covered with 20–28 (24 on average) moderately angular, non-bifurcating costae, 4–6 in the unipli-cation. Beak tapering, suberect, beak-ridges ill-defined, rounded. Foramen circular, hypothyrid. Interarea short, flat. Apical angle varying between 100° and 120°, on average 110°.

Internal characters (Fig. 4)

Ventral valve: dental lamellae subparallel or slightly convergent. Hinge-teeth massive, inserted in broad dental sockets,

inner and outer socket-ridges present. – Dorsal valve: cardinal process not observed. Hinge-plates wide, flat. Falcifer crura, given off dorsally, developed almost simultaneously with hinge-plates. Dorsal septum not present.

Discussion

Orbirhynchia parkinsoni was founded on „*Terebratulula sulcata*“ Parkinson, 1811 and has a long history which was recapitulated by Owen (1960). The allocation to the genus *Orbirhynchia* by Owen (loc. cit.) was based on the recognition of hinge plates bearing falcifer crura and the absence of a dorsal septum which are the main diagnostic features of the subfamily Lacunosellinae, represented in the Cretaceous by the single genus *Orbirhynchia*. This genus abounds in Upper Cretaceous with small and globular species, whereas *Orbirhynchia parkinsoni* has retained the trilobate shape and the relatively coarse ribs of its Jurassic ancestors, such as species of *Lacunosella* [*L. arolica* (Oppel 1865), *L. sparsicosta* (Quenstedt 1858) and others]. “Missing links” between these late Jurassic forms and *O. parkinsoni*, demonstrating some relations in external morphology might occur in the Early Cretaceous, with *Lacunosella hoheneggeri* (Suess 1859) and *Lacunosella contracta* (d’Hom-bres-Firmas 1842) as possible candidates (M. Sandy, personal communication).

Some other rhynchonellids occurring in the terminal Aptian and the Albian show a striking external homeomorphy with *O. parkinsoni*. *Rhynchonella deluci* Pictet 1872 was shown to have



Fig. 5. *Burrirhynchia tripartita* (Pictet). Specimen from Lädtobel. Dorsal, ventral, lateral, anterior, and posterior view. Anterior and posterior views with dorsal valve up. $3 \times$ natural size. Inventory number P. 8935.

radulifer crura and a dorsal septum and was assigned to the genus *Cyclothyris* (Calzada 1976 and unpublished results, personal communication). The general shapes of both *Cyclothyris deluci* and *O. parkinsoni* are very similar. A clear separation may become difficult when based on external morphology only. The main discriminating characters are the denser ribs (33–46 costae) of *C. deluci*, the more accentuated convexity of the dorsal valve, the often somewhat irregular and asymmetric front-margin, the narrower apical angle (90° – 100°), and the truncated anterior end.

Further cyclothyrids bearing a strong resemblance to *O. parkinsoni* are apparently restricted to the Aptian of England. They were described by Owen (loc. cit.) as species of the genus *Lamellaerhynchia* Burri, 1953. They are not further discussed here.

The specimens from the Plattenwald Bed are assigned to *Orbirhynchia parkinsoni* on the grounds of the shell characters and the presence of falcifer crura. To our knowledge this species has not been identified from the Helvetic Alps yet. They are of small size when compared to the holotype and the topotype specimens of the Cambridge area which are almost twice as large. However, they are comparable with conspecific finds in south-eastern France and east-central Sardinia. No attempt has been made to assign these forms to one of the four variations (*paludensis*, *paucicostata*, *rencurelensis*, *salazacensis*) which Jacob & Fallot (1913) proposed and which, in part, were taken over by Dieni et al. (1974) and Middlemiss & Owen (1980).

Rhynchonellids from diverse outcrops of the Alpine Albian were reported under different names. The poor definition of these “species” was the main reason for numerous misidentifications. Schaad (1925) and Heim & Seitz (1934) expressed their doubts about correct naming when they listed from the Plattenwald Bed: *Rh. deluci*, *Rh. cf. sulcata* and *Rh. sp.*, respectively. Jacob & Tobler (1906) and Fichter (1934) mentioned *Rh. deluci* from the Helvetic Alps in central Switzerland, and Delamette et al. (1997) listed it from the western Helvetic. The real nature of these rhynchonellids remains to be solved.

Order Rhynchonellida Kuhn 1949
 Superfamily Rhynchonelloidea Gray 1848
 Family Rhynchonellidae Gray 1848
 Subfamily Cyclothyridinae Makridin 1955
 Genus *Burrirhynchia* Owen 1962
 [type species: *Rhynchonella leightonensis* Lamplugh & Walker 1903]

***Burrirhynchia tripartita* (Pictet 1872)**

(Fig. 5)

- 1872 *Rhynchonella tripartita* Pictet: 44–45, 54; pl. 200, fig. 4,5
 1934 *Rhynchonella tripartita* Pictet; Heim & Seitz: 209
 1962 “*Rhynchonella*” *tripartita* Pictet (assigned to *Burrirhynchia*); Owen: 60
 1965 *Rhynchonella tripartita* Pictet 1872; Alméras: 384
 1986 *Burrirhynchia tripartita* (Pictet); Lobacheva: 129–133
 1999 *Burrirhynchia tripartita* (Pictet); Sulser: 87

Lectotype (designated here) in Pictet (1872, pl. 200, fig. 4a–4d), from the „Gault inférieur” of Perte-du-Rhône (Dépt. Ain, France). Repository: Museum of Natural History (Geneva, Switzerland); specimen number: A II-39-6706.

Occurrence and stratigraphy

Burrirhynchia tripartita is a rare species and, besides the type-locality, known only from the Albian of Clar (Dépt. Var, France) and Georgia in southern Russia (Kvakhadze 1973; cit. by Lobacheva 1986).

Description

External characters

Dimensions (mm) *	Length (L)	Width (W)	Thickness (T)	W : L	T : L
	9.3–17.7	8.8–22.0	7.6–14.5	0.80–1.24	0.68–0.82

*Number of specimens: 5 (Plattenwald: 1; Rotenbachalpe: 2; Gütle: 1; Tädtobel: 1)

Shell rounded-subpentagonal in outline, globose, almost equal in length and width, or spade-like and a little longer than broad. Position of greatest width variable, greatest thickness

in posterior third of length. Dorsal valve much inflated, laterally compressed. Ventral valve less convex than dorsal, with a sinus and a marked anterior geniculation, forming a right angle and giving rise to a lingual extension towards the front-margin. Wide trapezoidal or rectangular uniplication. Each valve covered with 35–50 fine costellae which, immediately before reaching the anterior edge, disappear and fuse to a pointed zig-zag-line, suggesting an antichotomic pattern of ribs. The dorsal apex is short and tumid and pressed down to the ventral valve so that the hinge line draws a conspicuous half-circular inflexion (Fig. 5, dorsal and posterior view). After a straight, oblique section the commissure follows the geniculation of the sinus and turns in a sharp angle dorsalward (Fig. 5, lateral view). Foramen small, cyclothyrid, beak ridges rounded, apical angle 80°–90°.

Internal characters

Shortage of material did not allow the taking of serial sections. A few internal characters could be observed on internal molds: a dorsal septum, extending to about one third of valve length and persistent, subparallel dental lamellae (fig. 5, ventral and posterior view).

Discussion

Burrirhynchia tripartita may be synonymous with *Burrirhynchia leightonensis* (Lamplugh & Walker 1903) from the Aptian/Albian boundary of England, chosen as type-species of *Burrirhynchia* (Owen 1962, Owen et al. 1968). Both nominal species are much alike. They have in common some noticeable characters, such as the described type of uniplication, the course of the lateral commissure, and the unusual pattern of costellation at the anterior commissure, well illustrated by Pictet (1872, pl. 200, fig. 4c and 4e). As long as the internal structures remain unknown, they have to be treated as separate species.

Owen (1988) suggested that two other rhynchonellids may be adaptations of *Burrirhynchia leightonensis* due to changing environments. These are the Cenomanian *Burrirhynchia sigma* (Schloenbach 1867) with a sigmoidal uniplication and *Burrirhynchia devoniana* Owen 1988, at present known only from a single locality in England. The latter has a similar anterior commissure as *B. tripartita*, but is covered with more marked costae.

Burrirhynchia tripartita is the rarest brachiopod of the Plattenwald Bed. It has been reported sporadically from only few and distant localities. The few specimens we disposed of display a wide and discontinuous variation of size. This brachiopod seems to grow in variable directions during ontogenetical development. A specimen of 22.0 mm width is almost three-fold as large as the smallest one (8.8 mm wide). In spite of the great span of shell size the typical characters are kept constant. It is a question, whether this behaviour is due to environmental factors, e.g. changing supply of nutrients or whether it reflects an abnormal pathological development.

Order	Terebratulida Waagen 1883
Superfamily	Terebratuloidea Gray 1840
Family	Terebratulidae Gray 1840
Subfamily	Rectithyrididae Muir-Wood 1965
Genus	<i>Moutonithyris</i> Middlemiss 1976 [type species: <i>Terebratula moutoniana</i> d'Orbigny 1849]

Moutonithyris dutempleana (d'Orbigny 1851*)

(Fig. 6–10)

- 1812 *Terebratula buplicata* J. Sowerby: 201: pl. 90, fig. 1 (only)
 1847 *Terebratula Dutempliana* (sic) d'Orbigny: 93
 1851 *Terebratula Dutempleana* d'Orbigny: pl. 511, fig. 1–3, 6, 7 only
 1871 *Terebratula buplicata* Sow.: Quenstedt: 381, pl. 48, fig. 13, 61–67
 ? 1872 *Terebratula dutempleana* d'Orbigny: Pictet: 82: pl. CCV, fig. 1–5
 1872 *Terebratula buplicata* Sow.: Stoliczka: 19: pl. 4, fig. 2–13
 1872 *Terebratula buplicata* var. *dutempleana*: Stoliczka: 20: pl. 4, fig. 14–17
 1874 *Terebratula buplicata* Sow.: Davidson: 33: pl. 5, fig. 1, 2
 ? 1896 *Terebratula dutempleana* d'Orbigny: Loriol: 142: pl. 5, fig. 10, 11
 1903 *Terebratula buplicata* var. *dutempleana*: Lamplugh & Walker: 251: pl. 17, fig. 1a–b
 1925 *Terebratula dutempleana* d'Orb.: Schaad: 15
 1925 *Terebratula buplicata* Sow.: Schaad: 15
 1934 *Terebratula dutempleana* d'Orbigny: Heim & Seitz: 198, 217, 221, 229, 230
 1965 *Terebratula dutempleana* d'Orbigny 1847: Alméras: 385
 ? 1972 *Praelongithyris dutempleana* (Orb.): Smirnova: 66: pl. 5, fig. 3
 1974 “*Terebratula*” *dutempleana* d'Orbigny: Dieni, Middlemiss & Owen: 185–190: pl. 33, fig. 2; text-fig. 9
 1976 “*Terebratula*” *dutempleana* d'Orbigny: Gaspard: pl. 1–5
 1976 *Moutonithyris dutempleana* (d'Orbigny): Middlemiss: 65
 1983 *Biplicatoria ferruginea* Cooper: 174–175: pl. 21, fig. 1–6; pl. 67, fig. 12, 13, 23, 24
 1983 *Biplicatoria hunstantonensis* Cooper: 176: pl. 20, fig. 21–27
 1988 *Moutonithyris dutempleana* (d'Orb.): Gaspard: 197–201: pl. 24, fig. 1–8; pl. 25, fig. 1–6; textfig. 78, 79
 1988 *Moutonithyris dutempleana* (d'Orbigny): Owen: 122–125: pl. 10; pl. 11, fig. 4–6; pl. 18, fig. 23, 24; text-fig. 21
 1999 *Moutonithyris dutempleana* (d'Orbigny): Sulser: 147 (with fig.)

For the completion of synonymy compare Dieni et al. (1974), Gaspard (1988) and Owen (1988)

* We prefer the year 1851 as the correct publication date (instead of 1847–1851), because d'Orbigny published the plates in 1851

Lectotype, designated by Dieni et al. (1974), in J. Sowerby (1815), pl. 90, fig. 1 from the Cambridge Greensand (Albian?) of the Castle Hill, Cambridgeshire (England).

Occurrence and stratigraphy

Moutonithyris dutempleana is a widespread species in the Albian of England (Cambridge and Norfolk area) and of many localities in France. It is widely distributed in the terminal levels of the Early Cretaceous in the Alpine regions of Europe. It further occurs in the Carpathians of Poland, in northern Spain, Ibiza, and doubtfully in the Cenomanian of England and northern Germany. Possible occurrences are reported from the Albian of Madagascar, the Caucasus, the Crimea and India. Reports from the french and swiss Jura region require confirmation.




Fig. 6. *Moutonithyris dutempleana* (d'Orbigny). Morph A. 2 × natural size. A: Specimen with oval outline and weakly saddled uniplication from Plattenwald. Dorsal and anterior view. Inventory number P. 17123. B: Specimen with pear-like, smoothly pentagonal outline from Sattelberg. Dorsal view. Inventory number P. 17124. C: Relatively broad specimen with slightly curved lateral commissure and almost rectimarginate front, from Plattenwald. Dorsal, lateral, anterior, and posterior view. Inventory number P. 17125.

Description

External characters

Dimensions (mm) *	Length (L)	Width (W)	Thickness (T)	W : L	T : L
	15.1–25.4	12.0–21.2	8.3–16.2	0.64–0.97	0.49–0.79

*Number of specimens: over 100 (Plattenwald: 43; Sattelberg: 65; Orsanka: 23)

Shell elongated-oval or pear-like in outline, with greatest width at or just anterior of mid-length, biconvex, with maximum convexity near the apex, either on dorsal or ventral valve. Anterior commissure variable: uniplicate to more or less biplicate, with either none or very low median depression, or distinctly sulcinate to episulcate. Lateral commissure arched in ventral direction. Beak sub-erect to erect. Foramen large, labiate. Deltidium not visible. Apical angle 60–80°.

Two variants, designated as morph A and B, can be distinguished:

Morph A (Fig. 6) is almost rectimarginate or displays a very flat uniplicate broad anterior elevation of not more than 1–2 mm. In rare specimens a tendency towards a weak biplication can be observed. The lateral commissure is nearly straight.

Morph B (Fig. 7) has a sulcinate or episulcate anterior commissure with distinct plicae and a pronounced median

sinus of 2–5 mm depth. The lateral commissure is distinctly curved. Impressions of the vascular system are occasionally visible (Fig. 7A, dorsal view), and, also rarely, an accumulation of growth-lines.

Internal characters (Fig. 8)

The internal morphology could be studied only partially by transverse sections. Some insight was obtained by isolated parts of the brachidium (Fig. 8, left). A distinct cardinal process is flanked by deep dental sockets (Fig. 8, right). Hinge plates horizontal, becoming convex later on, keeled at the inner edge. Crural processes not observed. Loop situated at about one third of shells length, partially exposed, probably narrow. No difference of internal characters between morph A and morph B was observed.

Discussion

The gradual change of Sowerby's species name „*biplicata*” to the now valid „*dutempleana*” is illustrated in the long list of synonyms. In the past this was a controversial species and the subject of many misinterpretations. After the internal structures had been studied in specimens of the type region, first by Middlemiss (in Dieni et al. 1974) and later by Gaspard (1988)




Fig. 7. *Moutonithyris dutempleana* (d'Orbigny). Morph B. $2 \times$ natural size. A: Specimen with distinct subpentagonal outline, strongly curved lateral commissure and episulcate front, from Plattenwald. Dorsal, lateral, and anterior view. On dorsal valve traces of the vascular system can be recognized. Inventory number P. 17126. B: Relatively broad specimen with much inflated dorsal valve, from Sattelberg. Dorsal and lateral view. Inventory number P. 17127. C: Average specimen from Plattenwald. Dorsal and lateral view. Inventory number P. 17128.

and Owen (1988), *Terebratula dutempleana* was assigned to the genus *Moutonithyris*. Related species could then be discriminated and partly assigned to pre-existing genera such as *Praelongithyris* and *Cyrtothyris*. The important diagnostic character of *Moutonithyris*, the hammer-shaped hinge plates, was well recognized in the Plattenwald specimens.

In many localities of the Alpine Albian forms ascribed to *Moutonithyris dutempleana* occur in phosphatic and siliciclastic beds. Internal structures are mostly destroyed, and identification has to be based on external morphology alone. Its presence elsewhere in the Helvetic Alps was reported by Renevier (1854), Moesch (1878, 1881, 1894), Jacob & Tobler (1906),

Oberholzer (1933), Fichter (1934), and other authors. However, paleontological confirmation remains to be done for these localities. In cases of undated stratigraphy the risk of confusion with *Moutonithyris moutoniana* (d'Orbigny) which does not reach the Albian stage, is possible. Normally, *M. moutoniana* displays a more circular outline and is less biplicate, but in unusual large individuals which show a tendency to become elongate and to develop a distinct biplication, both species look much the same. *M. moutoniana* colonized the marginal areas of the Tethys during the whole Early Cretaceous. It is a rather common element in the Aptian of the Helvetic, and it seems likely that *M. dutempleana* became its successor by replacement (Middlemiss 1973, 1979).




Fig. 8. *Moutonithyris dutempleana* (d'Orbigny). Specimens from Plattenwald. Left: Morph A. Almost complete brachidium with parts of the transverse band, exposed from dorsal side by acid treatment. Convex crura with distinct inner edges, a character of *Moutonithyris*. $4 \times$ natural size. Right: Morph B. Ventral view of deep and long dental sockets enclosing a porous cardinal process. The distal parts of the crura and the transverse band are not preserved. $6 \times$ natural size.




Fig. 9. Population of *Moutonithyris dutempleana* (d'Orbigny) from Plattenwald. Shell length and thickness of 21 specimens of morph A (thickness:length ratio = 0.49–0.62) and 16 specimens of morph B (thickness:length ratio = 0.60–0.79).

External and internal morphology leave no doubt that the populations of the Plattenwald Bed can be assigned to *M. dutempleana*. The multitude of specimens shows a shell length of 15–22 mm. Larger individuals were found at Orsanka. In all sampled localities *M. dutempleana* occurs in two different forms, named morph A und morph B. Their internal structures are identical and their distribution is approximately 50:50. We think that both forms should be considered morphotypes of one species, since practically no transient forms can be observed and so a clear-cut (sexual?) dimorphism results. The different shell shapes of morph A and B can be discriminated by biometrical means. Specimens of morph B display a thickness:length ratio which is always greater than 0.6 whereas those of morph A are below this (Fig. 9). Otherwise the width:length ratio shows about a similar distribution for both morphs, and it is not possible to tell them apart (Fig. 10).

A brachiopod, very closely related to *M. dutempleana*, was named by J. de C. Sowerby (1825) *Terebratula obtusa*, associated with *dutempleana* in the Albian Greensand of the Cambridge area. This companionship was registered at many other places where *M. dutempleana* occurs. Davidson (1852–54) namely distinguished both *dutempleana* and *obtusa*, but united them in his synonymy

of *Terebratula buplicata* (now *Moutonithyris dutempleana*). The internal structures of both „species“ are identical. Gaspard (1988, 1997) treated both nominal taxa as different species and maintained that *Terebratula (Moutonithyris) obtusa* is confined to the Cambridge Greensand. We do not opt for either interpretation. However, in an informal manner morph A might be called the „*obtusa* type“ and morph B the „*dutempleana* type“, because these affinities are quite obvious.

Superfamily Terebratelloidea King 1850
 Family Dallinidae Beecher 1893
 Subfamily Gemmarculinae Elliott 1947
 Genus *Gemmarcula* Elliott 1947 [type species:
Terebratula truncata J. de C. Sowerby 1826]

***Gemmarcula ? plattenwaldensis* sp. nov.**
 (Fig. 11, 12)

Terebratula cf. *moreana* sensu Seitz: *in schedulis*.

Derivation of name: *plattenwaldensis* – according to the stratotype locality of the „Plattenwald-Schicht“.




Fig. 10. Same population of *Moutonithyris dutempleana* (d'Orbigny) as in fig. 9. Shell length and width of 21 specimens of morph A (width:length ratio = 0.64–0.89) and 16 specimens of morph B (width:length ratio = 0.69–0.97).

Holotype in Fig. 11A, from the Plattenwald Bed (Albian). Specimen number: P. 10318. Type locality: Plattenwald near Klaus (Vorarlberg, Austria). Paratype in Fig. 11B, same locality. Specimen number: P. 11923.

Description

External characters

Dimensions (mm) *	Length (L)	Width (W)	Thickness (T)	W : L	T : L
	9.9–13.3	9.5–12.7	8.0–10.0	0.94–1.01	0.62–0.84

*Number of specimens: 9 (Plattenwald: 6; Sattelberg: 1; Gopfberg: 2)

Shell biconvex, stout, rounded-pentagonal to subcircular in outline, oval to subcircular in profile, length and width equal, greatest width and greatest thickness in the middle of the shell. Dorsal valve with three broad rounded folds and deeply furrowed spaces in between, often with distinct muscle scars. Ventral valve with two corresponding strong folds on each side of a deeply incised sinus. Occasionally with a thickened area

near the lateral margins. Test (when preserved) covered with about 24 radiating, intercalate costellae. Anterior commissure antiplicate. Beak stout, suberect, truncated by a large foramen. Beak-ridges rounded, hinge-line narrow, interarea triangular. Apical angle 80–90°.

Internal characters (Fig. 12)

A short sequence of sections revealed a broad transverse band (Fig. 12, section 5.2 mm). Some further details of the internal structures were detected on weathered internal moulds: a long dorsal septum, extending to at least 75% of total shell length, and muscle scars (diductors) on the posterior half of the valve (compare Fig. 11A, dorsal valve).

Discussion

Only two specimens were available for sectioning. The study of the internal morphology is therefore incomplete, but further details were detected on the surface of the internal moulds: a long loop and a high and long dorsal septum, extending to



Fig. 11. *Gemmarcula? plattenwaldensis* sp. nov. $2 \times$ natural size. A: Holotype from Plattenwald. Dorsal, ventral, lateral, and anterior view. Anterior view with dorsal valve up. On the dorsal valve the muscle scars are visible. Inventory number P. 10318. B: Specimen from Plattenwald. Dorsal valve with preserved fragments of ribbed test. Inventory number P. 11923. C: Specimen from Gopfberg. Posterior view. Inventory number P. 11451. D: Specimen from Gopfberg. Posterior view, showing long septum and dental lamellae. Inventory number P. 11450.



Fig. 12. *Gemmarcula? plattenwaldensis* sp. nov. Transverse serial sections of a specimen (length = 10.5 mm, width = 10.1 mm, thickness = 9.0 mm) from Plattenwald. Numbers give distance in mm from the posterior end. $3 \times$ nat. size.

about 75% of total shell length. There is scarcely any doubt that these brachiopods are terebratellids, but a satisfactory identification on the grounds of relevant known species was not possible. The species of the Plattenwald Bed under study was named *plattenwaldensis* sp. nov.

O. Seitz compared these terebratellids with *Terebratula* cf. *moreana* (collection labels) without giving an authorship for this taxon. *Terebratula moreana* d'Orbigny (1851, pl. 506, fig. 13–16) shows affinities to *T. reticulata* Schlotheim and *T. puscheana* Roemer both of which Owen (1977) synonymized with his *Oblongarcula alemannica*, a Hauterivian species of the Hilsconglomerat in northern Germany and in France (Dépt. Haute-Marne). A common character of these forms are three broad folds on the dorsal valve which are seen also on the Plattenwald specimens, but their general shell morphology is distinctly different.

A Cenomanian species of northern Germany, *Megerleia* (?) *hercynica* Schloenbach 1867, assigned to *Gemmarcula* Elliott 1947 by Owen (loc. cit.), and the supposedly synonymous *Terebratella kurskensis* Hofmann 1869 differ from *plattenwaldensis* in having seven or rarely five dorsal folds and a much shorter septum. *Arenaciarcula acuticosta* Owen 1977 of the Lower Albian of England (Bedfordshire), originally described by Lamplugh & Walker (1903) as Schloenbach's species mentioned above, and „*Eudesia*“ *tekedikensis* Smirnova 1972 from the Upper Albian of Turkmenistan differ from

the Plattenwald individuals by their pattern of costae and the fan-shaped outline.

A strong evidence to assign the species *plattenwaldensis* either to the genus *Gemmarcula* Elliott 1947 or *Arenaciarcula* Elliott 1959 is lacking. The broad transverse band may be indicative for both genera. Therefore an open nomenclature is used.

Other brachiopods (Fig. 13)

The Plattenwald Bed furnished some further brachiopods in very small number. A reliable description is not possible, but they are mentioned and figured in order to complete the faunal spectrum of the brachiopod assemblage. The one, a thick globular brachiopod with a massive, much incurved umbo and a well visible mark of the dorsal septum on the surface of the mould has some affinity to representatives of *Kingena* (Fig. 13A). Other forms like the preceding one are less thick and fan-shaped, with a maximum width shifted to the anterior line (Fig. 13B).

5. Conclusions

The brachiopods of the Plattenwald Bed are a minor and accessory element in a fauna which is clearly dominated by ammonites. These indicate a time span which reaches from the




Fig. 13. Unidentified brachiopods from Plattenwald. 2 × natural size. A: Dorsal valve with very long septum. Inventory number P. 5885. B: Dorsal and lateral view of a more globular specimen. Collection H. S.

uppermost Aptian to the top of the Albian (Föllmi 1989b). This is in agreement with the age of the brachiopods, as far as their life span is known. *Burrirhynchia tripartita* is known hitherto only from the tardefurcata-zone and thus represents the Lower Albian. *Moutonithyris dutempleana* and *Orbirhynchia parkinsoni* are characteristic for the Upper Albian and basal Cenomanian. The Early and terminal Albian are well documented by the brachiopods.

When we look at the brachiopod assemblage in the Tethyan (Alpine) and Subtethyan (Jura) domains, a parallelism in the evolution of Early Cretaceous stages is obvious (Gaspard 1999). One of the reasons is the development of similar facies in the „Schrattenkalk“ and in the Urgonian, culminating and widespread in the Barremian and Aptian. In the Albian the brachiopod communities change. They developed in highly condensed, geographically limited, spot-like areas and with a very slow sedimentation rate. As a whole, the Plattenwald brachiopods can be characterized as the remains of a small, adaptive group of generalists with affinities to a fauna of the boreal type.

In the Plattenwald Bed a normal oxygen supply with a tendency to eutrophism is assumed (Föllmi, personal communication). The brachiopods are generally of small size, but in one locality (Orsanka) larger individuals of *Moutonithyris dutempleana* coexist with small ones. In view of the long time during which the Plattenwald Bed accumulated, one can argue that brachiopods of different size did not live simultaneously, but reflect periods of more or less favorable conditions of life.

Gemmarcula ? plattenwaldensis sp. nov. shows a remarkable morphological steadiness with regard to size, shell ornament and shape. This low variability may reflect a life in protected niches. The species is rare yet, perhaps an indication that favorable conditions did probably not last for long.

Acknowledgements

Museums work would be impossible without donations from private collectors. Some of the brachiopods studied were donated to the museum by Joe Vincenz (Ludesch) and the late R. Gabriel (Weiler). Prof. Dr. K. Föllmi (Neuchâtel, Switzerland), Dr. Danièle Gaspard (Paris) and Dr. Michael Sandy (Dayton OH, U.S.) were so kind to read thoroughly the manuscript. They gave valuable and substantial suggestions for improvement.

REFERENCES

- ALMÉRAS, Y. 1965: Bibliographie des brachiopodes du Crétacé Inférieur (France et pays limitrophes). Etat de nos connaissances actuelles sur la répartition stratigraphique. Mém. Bur. Rech. Géol. Min. No. 34, 373–391.
- BACHMANN, I. 1864: Ueber die cretacischen Brachiopoden des Pilatus, Vitznauerstocks und der Hochflue. Mitt. naturf. Ges. Bern 190–193.
- BUCKMAN, S.S. 1918: The Brachiopoda of the Namyau beds, northern Shan States, Burma. Mem. geol. Surv. India, Palaeont. indica [n.s.] 3/2, 1–299, pls. 1–21 (1917).
- BURRI, F. 1953: Beiträge zur Systematik der Brachiopoden aus der untersten Kreide im westschweizerischen Jura-gebirge. Eclogae geol. Helv. 46/2, 269–285, 1 pl.
- 1956: Die Rhynchonelliden der Unteren Kreide (Valanginien-Barrémien) im westschweizerischen Jura-gebirge. Eclogae geol. Helv. 49/2, 599–701, 15 pls.
- CALZADA, B.S. 1976: Braquiopodos infracretácicos del Levante español. Trab. Mus. geol. Semin. Barcelona 14 (nr. 149), 1–86, pls. 1–14.
- COOPER, G.A. 1983: The Terebratulacea (Brachiopoda). Triassic to Recent: A study of the Brachidia (Loops). Smithson. Contrib. Paleobiology nr. 50, 445 p., 77 pls.
- DAVIDSON, TH. 1852–1854: A Monograph of the British Fossil Brachiopoda. II. The Cretaceous Brachiopoda. Mon. Palaeont. Soc. London 1/2, 1–117, pls. 1–12.
- DELAMETTE, M., CHAROLLAIS, J., DECROUEZ, D. & CARON, M. 1997: Les grès verts helvétiques (Aptien moyen-Albien supérieur) de Haute-Savoie, Valais et Vaud (Alpes occidentales franco-suissees). Publ. Dép. géol. paléont. Univ. Genève 23, 400 pp., 40 pls.
- DIENI I., MIDDLEMISS F.A. & OWEN E.F. 1974: The Lower Cretaceous Brachiopods of east-central Sardinia. Boll. Soc. Paleontologica Italiana 12/2, 166–216, pls. 32–38, text-figs. 1–18 (1973).
- ELLIOTT, G.F. 1947: The development of a British Aptian brachiopod. Geologist's Assoc. Proc. 58, 144–159.
- 1959: Six new genera of Mesozoic Brachiopoda. Geol. Mag. 96, 146–148.
- FICHTER, H.J. 1934: Geologie der Bauen-Brisen-Kette am Vierwaldstättersee und die zyklische Gliederung der Kreide und des Malms der helvetischen Decken. Beitr. geol. Karte Schweiz [N.F.] 69, 128 pp., 3 pls.
- FÖLLMI, K.B. (1989a): Evolution of the Mid Cretaceous triad: platform carbonates, phosphatic sediments, and pelagic carbonates along the northern Tethys margin. Lecture Notes in Earth Sciences 23, 153 pp.
- (1989b): Beschreibung neugefundener Ammonoidea aus der Vorarlberger Garschella-Formation (Aptian-Albian). Jahrb. Geol. B.-A. 132/1, 105–189, Wien.
- FÖLLMI, K.B. & OUWEHAND, P.J. (1987): Garschella – Formation und Götzis – Schichten (Aptian-Coniacian): Neue stratigraphische Daten aus dem Helvetikum der Ostschweiz und des Vorarlbergs. Eclogae geol. Helv. 80/1, 141–191.
- FÖLLMI, K.B., WEISSERT, H., BISPING, M. & FUNK, H. 1994: Phosphogenesis, carbon-isotope stratigraphy, and carbonate-platform evolution along the Lower Cretaceous northern Tethyan margin. Geol. Soc. America Bull. 106, 729–746.
- GASPARD, D. 1988: Sellithyridinae, Terebratulidae du crétacé d'Europe occidentale. Dynamique des populations, systématique et évolution. Cahiers de paléontologie 242 p., 28 pls.
- 1989: Explosion de brachiopodes au crétacé inférieur dans la mer épicontinentale jurassienne. Rev. Paléobiol. vo. spéc.: 81–102.
- 1997: Distribution and recognition of phases in Aptian-Turonian (Cretaceous) brachiopod development in NW Europe. Geologica carpathica 48/3, 145–161. Bratislava.
- 1999: Distribution of Lower Cretaceous brachiopods in Europe (Berriasian to Aptian). Bull. Soc. géol. France 170/3, 311–326.
- GEBHARD, G. 1985: Kondensiertes Apt und Alb im Helvetikum (Allgäu und Vorarlberg) – Biostratigraphie und Fauneninhalt. In: H.A. Kollmann (Ed.): Beiträge zur Stratigraphie und Paläogeographie der mittleren Kreide Zentral-Europas. Österr. Akad. Wiss. Schriftenreihe Erdwiss. Komm. 7, 271–279.

- HEIM ARN. & SEITZ O. 1934: Die mittlere Kreide in den helvetischen Alpen von Rheintal und Vorarlberg und das Problem der Kondensation. Denkschr. Schweiz. Naturf. Ges. 69/2, 185–310, 3 pls.
- HOFMANN, E. 1869: Materials for the Geology of Russia 1, 1–99, pls. 1–19. St. Peterbourg.
- JACOB, CH. & FALLOT, P. 1913: Etude sur les Rhynchonelles portlandiennes, néocènes et méso-crétacées du Sud-Est de la France. Abh. Schweiz. Paläont. Ges. 39/2, 82 pp., 11 pls.
- JACOB, C. & TOBLER, A. 1906: Etude stratigraphique et paléontologique du Gault de la Vallée de la Engelberger Aa (Alpes calcaires suisses, environs du Lac des quatre cantons). Abh. Schweiz. Paläont. Ges. 33/5, 26 pp., 2 pls.
- KAUFMANN, F.J. 1877: Kalkstein- und Schiefergebiete der Kantone Schwyz und Zug und des Bürgenstocks bei Stans. Beitr. geol. Karte Schweiz 14/2, 180 pp., 5 pls.
- LAMPLUGH, G.W. & WALKER, J.F. 1903: On a fossiliferous band at the top of the Lower Greensand near Leighton Buzzard (Bedfordshire). Q. J. geol. Soc. London 59, 234–265, pls. 16–18.
- LORIOU, P. DE 1864: Description de quelques brachiopodes crétacés. Mém. Soc. phys. Hist. nat. Genève 17/2, 437–447, 1 pl.
- 1868: Monographie des couches de l'étage Valangin des carrières d'Arzier (Vaud). Matér. Paléont. Suisse 4/2, 110 pp., 9 pls.
- LOBACHEVA, S.V. 1986: Early Cretaceous brachiopods of the South USSR and some new data on Capillithyrinae. In: P.R. Racheboef & Ch. Emig: Les brachiopodes fossiles et actuels. 129–134. 1^{er} Congrès Internat. Brachiopodes, Brest, France. 1985. Collection „Biostratigraphie du Paléozoïque“. Univ. Bretagne Occidentale.
- MIDDLEMISS, F.A. 1973: The geographical distribution of Lower Cretaceous Terebratulacea in western Europe. In: R. Casey & P.F. Rawson (ed.): The boreal Lower Cretaceous. Geol. J. spec. Issue 5, 111–120.
- 1976: Lower Cretaceous Terebratulidina of northern England and Germany and their geological background. Geol. Jb. A30, 21–104, 11 pls., 30 text-figs.
- 1979: Boreal and Tethyan brachiopods in the European Early and Middle Cretaceous. In: J. Wiedmann (Ed.): Aspekte der Kreide Europas. Internat. Union geol. Sci. [A] 6, 351–361. Stuttgart.
- 1981: Lower Cretaceous Terebratulidae of the Jura region. 1. Revision of some species described by Pictet and De Loriol (1872). Eclogae geol. Helv. 74/3, 701–733.
- 1983: Lower Cretaceous Terebratulidae of the Jura region. 2. The Aptian fauna of La Presta and Noirvaux Dessus and the Upper Albian fauna of La Vraconne. Eclogae geol. Helv. 76/3, 669–687.
- 1984: Lower Cretaceous Terebratulidae of the Jura region. 3. *Terebratula salevensis* De Loriol and the genera *Praelongithyris* and *Tropeothyris*. Eclogae geol. Helv. 77/3, 583–617.
- 1989: The distribution of Terebratulidae in the Hauterivian of the Jura region. Mém. Soc. neuchâtel. Sci. nat. 1, 183–186.
- MIDDLEMISS, F.A. & OWEN, E.F. 1980: Condensed Cretaceous limestone in a section near Xiquena, Betic of Malaga, SE Spain. II. Albian (-Cenomanian) brachiopods. Proc. konink. nederl. Akad. wetensch. [B] 83/2, 201–208.
- MOESCH, C. 1878: Zur Paläontologie des Sentisgebirges. Über neue und weniger bekannte Petrefakten aus der Kreide des Sentisgebirges. In: A. Escher v.d. Linth: Geologische Beschreibung der Sentis-Gruppe. Beitr. geol. Karte Schweiz 13, 16 pp., 3 pls.
- 1881: Geologische Beschreibung der Kalkstein- und Schiefergebilde der Kantone Appenzell, St. Gallen und Schwyz. Beitr. geol. Karte Schweiz 14/3, 336 pp., 4 pls.
- 1894: Geologische Beschreibung des Centralgebietes der Schweiz. III. Abteilung: Geologische Beschreibung der Kalk- und Schiefergebirge zwischen dem Reuss- und Kienthal. Beitr. geol. Karte Schweiz 24, 307 pp., 35 pls.
- OBERHOLZER, J. 1933: Geologie der Glarneralpen. Beitr. geol. Karte Schweiz [N.F.] 28, 626 pp., 29 pls.
- OOSTER, W.A. 1863: Pétrifications remarquables des Alpes Suisses. Synopsis des brachiopodes fossiles des Alpes Suisses: 71 pp., 20 pls. Genf & Basel.
- ORBIGNY, A. d' 1847–1851: Paléontologie française. Description des Mollusques et rayonnés fossiles. Terrains crétacés. 4. Brachiopodes. 390 pp. (1847), pls. 490–599 (1851). Paris.
- OWEN E.F. 1960: A note on „*Rhynchonella*“ *sulcata* (Parkinson) from the Lower Cretaceous of Great Britain. Ann. & Mag. Nat. Hist. [13] 2, 248–256.
- 1962: The brachiopod genus *Cyclothyris*. Bull. Brit. Mus. (nat. Hist.) Geol. 11/2, 49–72, 3 pls.
- 1977: Evolutionary trends in some Mesozoic Terebratulacea. Bull. Brit. Mus. (Nat. Hist.) Geol. 28, 205–253, 3 pls.
- 1988: Cenomanian brachiopods from the Lower Chalk of Britain and northern Europe. Bull. Brit. Mus. nat. Hist. (Geol.) 44/2, 65–175.
- OWEN, E.F., RAWSON, P.F. & WHITHAM, F. 1968: The Carstone (Lower Cretaceous) of Melton, East Yorkshire, and its brachiopod fauna. Proc. Yorkshire geol. Soc. 36/4, 513–524, pls. 22–23.
- PARKINSON J. 1811: Observations of some of the strata in the neighbourhood of London and on the fossil remains contained in them. Trans. Geol. Soc. London 1, 324–354, 26 pls.
- PETTIT, N.E. 1954: A monograph on the Rhynchonellidae of the British Chalk. Palaeontogr. Soc. London. 1–26, pls. 1–2 (1950), 27–52, pl. 3 (1954).
- PICTET, F.J. 1872: In Pictet, F.J. & Campiche, G.: Description des fossiles du terrain crétacé des environs de Sainte-Croix. 5. partie: Brachiopodes (terminé par P. De Loriol). Matér. Paléont. Suisse. 158 pp., 14 pls.
- POPIEL-BARCZYK E. 1977: A further study of Albian-Cenomanian brachiopods from the environs of Annopol on the Vistula with some remarks on related species from the Cracow region, Poland. Prace Muz. Ziemi 26, 25–54, pls. 1–7.
- RENEVIER, E. 1854: Seconde note sur la géologie des Alpes vaudoises. Bull. Soc. vaud. Sci. nat. 4, 204–218.
- SCHAAD, H. 1925: Geologische Untersuchungen in der südlichen Vorarlberger Kreide-Flysch-Zone zwischen Feldkirch und Hochfreschen. Diss. Univ. Zürich.
- SCHLOENBACH, U. 1867: Über die Brachiopoden der norddeutschen Cenoman-Bildungen. Geogn.-paläont. Beiträge (E.W. Benecke) 1/3, 399–506, 3 pls.
- SMIRNOVA, T.N. 1972: Early Cretaceous brachiopods of the Crimea and the northern Caucasus. Akad. Nauk SSSR. 139 pp., 13 pls. (in russian). Moscow.
- SOWERBY, J. 1812: The Mineral Conchology of Great Britain: or coloured figures ... B. Meredith (London). Vol. 1, 234 p., 102 pls.
- STOLICZKA, F. 1872: The Cretaceous fauna of southern India. IV. The Brachiopoda. Mem. geol. Surv. India. Palaeont. indica 4/1, 1–32, pls. 1–7.
- SULSER, H. 1999: Die fossilen Brachiopoden der Schweiz und der angrenzenden Gebiete. Juragebirge und Alpen: 315 pp. Paleont. Inst. Mus. Univ. Zürich.
- WILLIAMS, A., CARLSON, S. J., BRUNTON, C.H.C., HOLMER, I.E. & POPOV, L.E. 1996: A supra-ordinal classification of the Brachiopoda. Phil. Transact. Roy. Soc. London (series B) 351, 1171–1193, 6 fig.

Manuscript received April 22, 2002

Revision accepted August 22, 2002

