

Zeitschrift: Eclogae Geologicae Helvetiae
Herausgeber: Schweizerische Geologische Gesellschaft
Band: 96 (2003)
Heft: 2

Artikel: Geology of the NW Indian Himalaya
Autor: Steck, Albrecht
Kapitel: Acknowledgments
DOI: <https://doi.org/10.5169/seals-169014>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 29.03.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

the Himalayan frontal thrusts with the probably synchronous Kishtwar and Larji-Kullu-Rampur domes in the middle of the High Himalaya. The models of Beaumont et al. (1994, 1996) and Escher & Beaumont (1997) suggest that the initiation of up-warping at the backstop may be controlled by reduction in the convergent material that can be accommodated by the subduction channel. During the late orogenic and active Himalayan phase of dextral transpression between India and Asia, the dome structures continue to be uplifted and are often limited by younger normal faults and flexures such as the N-striking Tso Moriri fault and the Tso Kar and Yurdi flexures. The latter is together with the WNW-striking Sanku flexure responsible for the active uplift of the Suru syntaxis (Suru dome).

Conclusion

In conclusion, the Himalayan range was built up during the convergence of the Indian and Asian plates by a typical succession of orogenic phases (Masclé 1985, Le Fort 1996, Hodges 2000), where the preceding phase influences the next younger one. The main phases of the NW Himalaya are enumerated in the following list:

- 1) The Late Cretaceous and Paleocene **Transhimalayan batholith phase** (prothimalayan phase, Hodges 2000), characterised by the 103-50 Ma Andean type Ladakh magmatism, the accretion of the Dras-Nindam arc, the accretion and later obduction of the Spongtang immature island arc forming the southern active border of Asia and forearc sediment deposition. The Transhimalayan batholith, together with the Asian mantle wedge, form the Asian backstop for the Himalayan range.
- 2) The **Shikar Beh phase**: the intra continental NE-verging Shikar Beh range of an unknown, probably late Paleocene age,
- 3) The Eocene **North Himalayan phase** (eohimalayan phase, Hodges 2000) creating the SW-directed North Himalayan accretionary wedge.
- 4) The late Eocene-Miocene **High Himalayan phase** (neohimalayan phase, Hodges 2000): The zone of dry intra-crustal melting below the North Himalayan range and the Shikar Beh nappe stack determined the future position of the Main Central thrust at the base of the High Himalayan or "Crystalline" nappe.
- 5) The late Miocene to present **Lesser Himalayan phase**, with the formation of the deep-seated intracrustal Main boundary thrust.
- 6) The active **Subhimalayan phase** with the Subhimalayan thrust in front and the Active Himalayan thrust at the base of the present Himalayan accretionary wedge.

Acknowledgments

Financial support by the Foundation of the 450th Anniversary of the University of Lausanne and the Dr. Joachim de Giacomo Foundation of the Swiss Academy of Natural Sciences are gratefully acknowledged. The Lausanne earth science Himalaya research project was sponsored by the Swiss National Science Foundation and the Herbette Foundation of the Faculty of Sciences of the University of Lausanne for more than twenty years. The reviewer Clark Burchfiel and Jean-Pierre Burg are thanked for their critical reading of the manuscript.

REFERENCES

- AGRARWAL, R.P. 1993: Geology and Biostratigraphy of the Upper Siwalik of Samba area, Jammu Foothills. *J. Himalayan Geol.* 4, 227-236.
- ALSDORF, D., BROWN, L., NELSON, K.D., MAKOVSKY, Y., KLEMPERER, S. & ZHAO, W. 1998: Crustal deformation of the Lhasa terrane, Tibet plateau from Project INDEPTH deep seismic reflection profiles. *Tectonics* 17, 501-519.
- APPEL, E., MÜLLER, R. & WIDDER, R.W. 1991: Paleomagnetic results from the Tibetan sedimentary series of the Manang area (north central Nepal). *Geophys. J. Int.* 104, 255-266.
- APPEL, E., PATZELT, A. & CHOUKER, C. 1995: Secondary palaeoremanence of Tethyan sediments from the Zaskar Range (NW Himalaya). *Geophys. J. Int.* 122: 227-242.
- ARGAND, E. 1924: La tectonique de l'Asie. *Congr. géol. int. Bruxelles 1922, XIIIe session.* 171-372.
- AUDEN, J.B. 1934: The Geology of the Krol belt. *Rec. Geol. Surv. India* 67/4 357-454.
- AVOUAC, J.P., BOLLINGER, L., LAVÉ, J., CATTIN, R. & FLOUZAT, M. 2001: Le cycle sismique en Himalaya. *C.R. Acad. Sci. (Paris)* 333, 513-529.
- AZMI, R.J. & PANCHOLI, V.P. 1983: Early Cambrian (Tommotian) conodonts and other shelly microfauna from the Upper Krol of Mussoorie Syncline, Garhwal Lesser Himalaya, with remarks on the Pricambrian-Cambrian boundary. *Himalayan Geol.* 11, 360-372.
- BAGATI, T.N. 1990: Lithostratigraphy and facies variation in the Spiti basin (Tethys), Himachal Pradesh, India. *J. Himalayan Geol.* 1, 35-47.
- BAGATI, T.N. 1991: Evolution of the Tethyan sedimentary basin in the western Himalaya. In: S.K. TANDON, C.C. PANT & S.M. CASSHYAP (eds.): *Sedimentary basins of India, tectonic context.* Gyanodaya Prakashan, Nainital, 218-235.
- BASSOULET, J.P., BELLIER, J.P., COLCHEN, M., MARCOUX, J. & MASCLÉ, G. 1978a: Découverte de Crétacé supérieur calcaire pélagique dans le Zaskar (Himalaya du Ladakh). *Bull. Soc. géol. France*, 7/20, 961-964.
- BASSOULET, J.P., COLCHEN, M., MARCOUX, J. & MASCLÉ, G. 1978b: Une transversale de la zone de l'Indus de Khalsi à Phothaksar, Himalaya du Ladakh. *C.R. Acad. Sci. (Paris)* 286, série D, 563-566.
- BASSOULET, J.P., COLCHEN, M., JUTEAU, TH., MARCOUX, J. ET MASCLÉ, G. 1980: L'édifice des nappes du Zaskar (Ladakh-Himalaya). *C.R. Acad. Sci. (Paris)* 290D, 389-392.
- COLCHEN, M., JUTEAU, T., MARCOUX, J., MASCLÉ, G. & REIBEL, G. 1983: Geological studies in the Indus Suture Zone of Ladakh (Himalaya). In: GUPTA, V.J. (Ed.): *Stratigraphy and Structure of Kashmir and Ladakh, Himalaya.* (p. 96-124). – Hindustan. Publ. Corp., Delhi.
- BASSOULET, J.P., COLCHEN, M., GILBERT, E., MARCOUX, J., MASCLÉ, G., SUTRE, E. & VAN HAVER, T. 1984: L'orogène himalayen au Crétacé. *Mém. Soc. géol. France*, N.S. 147, 9-20.
- BAUD, A., ARN, R., BUGNON, P., CRISINEL, A., DOLIVO, E., ESCHER, A., HAMMERSCHLAG, J.G., MARTHALER, M., MASSON, H., STECK, A. & TIÈCHE, J.C. 1982: Le contact Gondwana-péri-Gondwana dans le Zaskar oriental (Ladakh, Himalaya). *Bull. Soc. géol. France*, 24, 341-361.
- BAUD, A., ATUDOREI, V. & SHARP, Z. 1996: Late Permian and Early Triassic Evolution of the Northern Indian margin: Carbon Isotope and Sequence Stratigraphy. *Geodynamica Acta* 9, 57-77.
- GAETANI, M., GARZANTI, E., FOIS, E., NICORA, A. & TINTORI, A. 1984: Geological observations in southeastern Zaskar and adjacent Lahul area (northwestern Himalaya). *Eclogae geol. Helv.* 77, 171-197.