

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	8 (1962)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON THE CONSTRUCTION OF RELATED EQUATIONS FOR THE ASYMPTOTIC THEORY OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS ABOUT A TURNING POINT
Autor:	Langer, Rudolph E.
Kapitel:	3. The resultant.
DOI:	https://doi.org/10.5169/seals-37963

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 02.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The relations (2.5), (2.6) and (2.7) imply that $p+q = n$. We shall suppose the notation to be so assigned as to yield $p \leq q$. It follows then also that

$$\sum_{i=0}^k b_i(z) c_{k-i}(z) = p_{k,0}(z), \quad k = 0, 1, 2, \dots, n, \quad (2.8)$$

it being understood that the value 0 is to be assigned to any symbol $b_i(z)$ or $c_{k-i}(z)$ which, by virtue of its subscript, is not present in the formulas (2.7).

A change of dependent variable can be applied to the equation (1.1) to give it a form for which the coefficient $b_1(z)$ is identically zero. Considerable simplifications of the formulas result therefrom. We shall not resort to that normalization, however, refraining from it in order to keep the roles of the factors $B(\chi)$ and $\Gamma(\chi)$ interchangeable.

3. THE RESULTANT.

It is a hypothesis that the two polynomials (2.7) are relatively prime over the z -region. If we denote their resultant by $\Delta(z)$, we have accordingly

$$\Delta(z) \neq 0, \quad (3.1)$$

with

$$\Delta(z) \equiv \begin{vmatrix} 1 & b_1 & b_2 & \cdots & b_p & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & 1 & b_1 & b_2 & \cdots & b_p & 0 & \cdots & \cdots & 0 \\ \vdots & \vdots \\ 0 & \cdots & 0 \\ 1 & c_1 & c_2 & \cdots & \cdots & \cdots & c_q & 0 & \cdots & 0 \\ 0 & 1 & c_1 & \cdots & \cdots & \cdots & c_q & 0 & \cdots & 0 \\ \vdots & \vdots \\ 0 & \cdots & \cdots & 0 & 1 & c_1 & c_2 & \cdots & \cdots & c_q \end{vmatrix} \quad (3.2)$$

We shall find use for the relation (3.2). However, for future use we find it convenient to formulate the relative primary of

the factors $B(\chi)$ and $\Gamma(\chi)$ also in an alternative form. This is done as follows:

Let K designate the familiar matrix

$$K(z) = \begin{bmatrix} 0 & 0 & - & - & - & - & - & -b_p \\ 1 & 0 & - & - & - & - & - & -b_{p-1} \\ 0 & 1 & 0 & - & - & - & - & -b_{p-2} \\ - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - \\ 0 & - & - & - & - & 1 & -b_1 & \end{bmatrix} \quad (3.3)$$

The eigen-values of this are the roots x_i , $i = 1, 2, \dots, p$ of the equation $B(x) = 0$. If we designate by ξ_i an eigen-vector corresponding to x_i we have

$$K^h \xi_i = x_i^h \xi_i, \quad h = 1, 2, \dots, q,$$

and accordingly

$$\Gamma(K) \xi_i = \Gamma(x_i) \xi_i.$$

Thus $\Gamma(x_i)$ is an eigenvalue of the matrix $\Gamma(K)$, and, since the product of the eigenvalues is the determinant of the matrix, we have

$$|\Gamma(K)| = \prod_{i=1}^p \Gamma(x_i).$$

Observing that no factor on the right is zero, and giving to $\Gamma(K)$ its explicit form, we conclude with the result

$$|\sum_{j=0}^q c_j(z) K^{q-j}(z)| \neq 0 \quad (3.4)$$

4. TWO DIFFERENTIAL OPERATORS OF THE ORDERS p AND q .

Let the functions $\beta_j(z, \lambda)$ and $\gamma_i(z, \lambda)$ be taken to be polynomials of the degree $(r - 1)$ in $1/\lambda$, thus

$$\begin{aligned} \beta_j(z, \lambda) &= \sum_{v=0}^{r-1} \frac{\beta_{j,v}(z)}{\lambda^v}, \quad \beta_{j,0}(z) \equiv b_j(z); \quad j = 1, 2, \dots, p, \\ \gamma_i(z, \lambda) &= \sum_{v=0}^{r-1} \frac{\gamma_{i,v}(z)}{\lambda^v}, \quad \gamma_{i,0}(z) \equiv c_i(z); \quad i = 1, 2, \dots, q. \end{aligned} \quad (4.1)$$