

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 27 (1981)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: APPLICATION OF TOPOLOGY TO PROBLEMS ON SUMS OF SQUARES
Autor: Dai, Z. D. / Milgram, R. J.
DOI: <https://doi.org/10.5169/seals-51752>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 01.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

APPLICATION OF TOPOLOGY TO PROBLEMS ON SUMS OF SQUARES

by Z. D. DAI, T. Y. LAM ¹⁾ and R. J. MILGRAM ¹⁾

If $n = 1, 2, 4$ or 8 , the classical n -square identities imply that the product of two sums of n squares in any commutative ring A is also a sum of n squares in A . On the other hand, by a classical theorem of Hurwitz [L, p. 137], one knows that the same statement cannot hold for other natural numbers n .

One can study the same problem over *fields* instead of over commutative rings. Here, the solution of the problem is also known, albeit somewhat different. According to a remarkable theorem of Pfister [P], if $n = 2^m$ is any 2-power, and if u, v are sums of n squares in a field F , then their product uv is also a sum of n squares in F . (This implies that the set of nonzero elements in F which are a sum of $n = 2^m$ squares in F is a group under multiplication.) On the other hand, Pfister has also shown that the above statement cannot hold for all fields if n is not of the form 2^m .

Back to sums of squares in commutative rings again, the above two paragraphs suggest that, in considering the multiplication problem, it is perhaps more reasonable to confine one's attention to *units* of a ring A which are sums of 2^m squares in A . Writing $n = 2^m$ and $U(A)$ for the group of units in A , one can ask:

- (*) If $u, v \in U(A)$ are sums of n squares in A ,
is $uv \in U(A)$ also a sum of n squares in A ?

This is equivalent to asking if the set of units in A which are a sum of $n = 2^m$ squares in A is a *group* under multiplication. This problem, first raised by R. Baeza, appeared as "Question 12" in Knebusch's collection [K₂] of open problems in the Proceedings of the Quadratic Form Conference in Kingston, Ontario in 1976. Generalizing the work of Pfister, Knebusch [K₁] has shown that (*) has an affirmative answer in case A is a (commutative) *semilocal* ring.

¹⁾ Supported in part by NSF.

In this note, we shall furnish the following solution to Baeza's problem:

THEOREM 1. *The answer to (*) is affirmative (for all commutative rings A) iff $n = 1, 2, 4$ or 8 .*

In view of the classical square identities mentioned before, we need only show the "only if" part of the theorem. The idea of the proof is to apply (*) in a "generic" setting, and then use suitable topological machinery to derive the conclusion $n = 1, 2, 4$ or 8 . The topological result needed here is Adams' famous theorem $[A_1]$ on the nonexistence of Hopf invariant one. Surprisingly, this algebraic application of Adams' Theorem, though reasonably straightforward, seems to have escaped the notice of both algebraists and topologists.

Let n be a natural number for which (*) holds for any commutative ring A . We shall prove that $n = 1, 2, 4$ or 8 . (In the following, we do not need to assume n to be a power of 2 to begin with, though this would follow from Pfister's theorem.)

Let A be the ring obtained by localizing the polynomial ring

$$\mathbf{R} [x_1, \dots, x_n, y_1, \dots, y_n]$$

at the multiplicative set generated by

$$u = x_1^2 + \dots + x_n^2 \quad \text{and} \quad v = y_1^2 + \dots + y_n^2.$$

Then, by (*), the unit $uv \in U(A)$ is a sum of n squares in A , say

$$uv = \left(\frac{f_1}{u^r v^s} \right)^2 + \dots + \left(\frac{f_n}{u^r v^s} \right)^2, \quad f_i \in \mathbf{R} [x, y].$$

Clearing denominators, we get a polynomial equation:

$$(1) \quad \begin{aligned} & x_1^2 + \dots + x_n^2)^{2r+1} (y_1^2 + \dots + y_n^2)^{2s+1} \\ & = f_1(x, y)^2 + \dots + f_n(x, y)^2. \end{aligned}$$

Now we make the following key observation:

LEMMA 1. *Each $f_i(x, y)$ above is a "biform" in (x, y) , of bidegree $(2r+1, 2s+1)$ (i.e. viewing the y 's as constants, f_i is a form of degree $2r+1$ in x , and, viewing the x 's as constants, f_i is a form of degree $2s+1$ in y).*

Proof. View each f_i as a polynomial in x , and let f_{ij} denote its homogeneous component of degree j in x . We may write

$$f_i = f_{i,p} + f_{i,p+1} + \dots + f_{i,q} \quad (1 \leq i \leq n)$$

where p, q are independent of i . If $q > 2r + 1$, a comparison of terms of x -degree $2q$ on the two sides of (1) shows that

$$\sum_{i=1}^n f_{i,q}^2 = 0,$$

and hence $f_{i,q} = 0$ for all i . Similarly, if $p < 2r + 1$, we must have $f_{i,p} = 0$ for all i . Hence, f_i is a form in x of degree $2r + 1$. By symmetry, we infer that f_i is also a form in y of degree $2s + 1$. Q.E.D.

Now let x, y be points on the unit sphere S^{n-1} . The equation (1) above implies that the n -tuple

$$(f_1(x, y), \dots, f_n(x, y))$$

is also a point on S^{n-1} . Thus,

$$(x, y) \mapsto (f_1(x, y), \dots, f_n(x, y))$$

induces a polynomial (and hence continuous) mapping:

$$\mu: S^{n-1} \times S^{n-1} \rightarrow S^{n-1}.$$

Fix a base point $b \in S^{n-1}$. Then the compositions

$$S^{n-1} \rightarrow S^{n-1} \times \{b\} \xrightarrow{\mu} S^{n-1}$$

$$S^{n-1} \rightarrow \{b\} \times S^{n-1} \xrightarrow{\mu} S^{n-1}$$

are *odd* mappings, since each f_i has bidegree $(2r + 1, 2s + 1)$. By the theorem of Borsuk [B], these odd mappings from S^{n-1} to itself must have odd (topological) degrees, say, $2r' + 1$ and $2s' + 1$. Thus, the mapping μ has "type" $(2r' + 1, 2s' + 1)$ in the sense of Hopf [H₁].

Now by the Hopf Construction, the map μ induces a continuous map $\sigma: S^{2n-1} \rightarrow S^n$. Let

$$H: \pi_{2n-1}(S^n) \rightarrow \mathbf{Z}$$

be the Hopf invariant on the homotopy group $\pi_{2n-1}(S^n)$. According to Hopf [H₁, § 6], the homotopy class $[\sigma] \in \pi_{2n-1}(S^n)$ has Hopf invariant

$$H[\sigma] = \pm (2r' + 1)(2s' + 1),$$

which is an odd number. By Adams' theorem [A₁] on the nonexistence of Hopf invariant one (or odd Hopf invariant), one knows that this is possible only if $n = 1, 2, 4$ or 8 . This completes the proof of Theorem 1.

Adams' original solution of the Hopf invariant one problem took 85 pages, but there exists a proof using the powerful machinery of topological K -theory (cf. [A₆], [A₇, p. 137]) which, according to M. Atiyah, "can be written on a postcard". Thus, our Theorem 1 does admit a "short" proof. In fact, using K -theory, it is possible to obtain a more general version of Theorem 1. This will be deduced from the following topological statement:

THEOREM 2. *Let $\mu: S^{k-1} \times S^{n-1} \rightarrow S^{n-1}$ be a continuous mapping such that*

$$\mu(-x, y) = \mu(x, -y) = -\mu(x, y)$$

for all $x \in S^{k-1}$ and $y \in S^{n-1}$ (cf. [H₂]). Then $k \leq \rho(n)$, where ρ is the Hurwitz-Radon function.

(Recall that, if $n = 2^{4a+b} n_0$ where n_0 is odd and $b = 0, 1, 2$ or 3 , then, by definition, $\rho(n) = 8a + 2^b$.)

Before proving this theorem, let us first record several of its remarkable consequences in algebra. The first one is a result on real common zeros of biforms.

COROLLARY 1. *Let*

$$x = (x_1, \dots, x_k), y = (y_1, \dots, y_n).$$

Let $f_i(x, y)$ ($1 \leq i \leq n$) be biforms in (x, y) of odd bidegrees $(2r_i + 1, 2s_i + 1)$. If $k > \rho(n)$, then the real loci of $f_i = 0$ in the multiprojective space $\mathbf{RP}^{k-1} \times \mathbf{RP}^{n-1}$ have a common point.

Proof. If otherwise, we would have a mapping μ as in Theorem 2 defined by

$$\mu(x, y) = (f_1(x, y)/g(x, y), \dots, f_n(x, y)/g(x, y))$$

where $g(x, y) = (\sum f_i(x, y)^2)^{1/2}$.

COROLLARY 2. *Let*

$$F(x, y) = F(x_1, \dots, x_k; y_1, \dots, y_n)$$

be a biform of bidegree (d, e) where d, e are not multiples of 4, and $k > \rho(n)$. Suppose that

$$F(x_o, y_o) = 0 \ (x_o \in \mathbf{R}^k, y_o \in \mathbf{R}^n) \Rightarrow x_o = 0 \text{ or } y_o = 0.$$

Then F cannot be a sum of n squares in $\mathbf{R}[x, y]$.

Proof. This is clear from the above Corollary and the argument given in Lemma 1.

The next corollary may be viewed as a nonlinear generalization of the classical Hurwitz-Radon Theorem [L, p. 137]:

COROLLARY 3. For

$$x = (x_1, \dots, x_k), y = (y_1, \dots, y_n)$$

and fixed integers $r, s \geq 0$, the following statements are equivalent:

- (1) $(x_1^2 + \dots + x_k^2)^{2r+1} (y_1^2 + \dots + y_n^2)^{2s+1}$ is a sum of n squares in $\mathbf{R}[x, y]$;
- (2) $(x_1^2 + \dots + x_k^2)^{2r+1} (y_1^2 + \dots + y_n^2)^{2s+1}$ is a sum of n squares in $\mathbf{Z}[x, y]$;
- (3) $k \leq \rho(n)$.

Proof. (2) \Rightarrow (1) is obvious.

(1) \Rightarrow (3) follows from Corollary 2.

(3) \Rightarrow (2): It is enough to prove (2) for $r = s = 0$. This follows from $[G_1]$ or $[G_2]$.

For a commutative ring A , let $S_m(A)$ denote the set of sums of m squares in A , and let $US_m(A) = U(A) \cap S_m(A)$.

COROLLARY 4. For fixed integers k and n , the following statements are equivalent:

- (1) For any commutative \mathbf{R} -algebra A , $US_k(A) \cdot US_n(A) \subseteq US_n(A)$;
- (2) For any commutative ring A , $S_k(A) \cdot S_n(A) \subseteq S_n(A)$;
- (3) $k \leq \rho(n)$.

Proof. This is clear from Corollary 3 and the localization argument we have given before. (Note that Theorem 1 is a special case of this Corollary since it is well-known that $n \leq \rho(n)$ iff $n = 1, 2, 4$ or 8 .)

We shall now begin the proof of Theorem 2, using tools from K -theory, especially Adams' work on the J -homomorphism. For any finite CW -complex X , let $KO(X)$ denote the K -group of virtual real vector bundles over X , and

$\widetilde{KO}(X)$ the reduced K -group (modulo trivial bundles). Let $J(X)$ denote the group of stable fiber homotopy equivalence classes of virtual sphere bundles over X , and $\tilde{J}(X)$ the reduced J -group. The canonical J -homomorphism $J: KO(X) \rightarrow J(X)$ induces a homomorphism $\tilde{J}: \widetilde{KO}(X) \rightarrow \tilde{J}(X)$. We shall use Adams' results in the following form (see $[A_2, (7.4)]$, $[A_4, (3.5)]$ and $[A_5]$):

ADAMS' THEOREM. For $X = \mathbf{RP}^m$, \tilde{J} is an isomorphism $\widetilde{KO}(X) \cong \tilde{J}(X)$. The group $\widetilde{KO}(X)$ is cyclic of order $2^{\phi(m)}$ where $\phi(m)$ is the number of positive integers $\leq m$ which are congruent to $0, 1, 2$ or $4 \pmod{8}$. A generator for $\widetilde{KO}(X)$ is given by the canonical line bundle ξ_m over \mathbf{RP}^m .

On the product $S^{k-1} \times S^{n-1}$, we have an involution defined by

$$T(x, y) = (-x, -y);$$

let E be the quotient space $S^{k-1} \times S^{n-1}/T$. We have an $(n-1)$ -sphere bundle $\eta: E \rightarrow \mathbf{RP}^{k-1}$: this is the associated sphere bundle of the Whitney sum $n \cdot \xi_{k-1}$. Note that E has an involution $\tau(x, y) = (x, -y)$ which on each fiber is the antipodal map.

Assume that we have a continuous map

$$\mu: S^{k-1} \times S^{n-1} \rightarrow S^{n-1},$$

as in Theorem 2. Then μ induces a map $\bar{\mu}: E \rightarrow S^{n-1}$ which is equivariant with respect to the involution τ on E and the antipodal map on S^{n-1} . We have a fiber map

$$(\eta, \bar{\mu}): E \rightarrow \mathbf{RP}^{k-1} \times S^{n-1} \quad (\text{trivial bundle over } \mathbf{RP}^{k-1})$$

which (by the theorem of Borsuk again) has odd degree d on each fiber. By the "mod d -Dold Theorem" $[A_3, (1.1)]$, there exists an integer $e \geq 0$ such that $d^e \cdot \eta$ is fiber homotopy equivalent to a trivial bundle. Since $\tilde{J}(\mathbf{RP}^{k-1})$ is 2-primary, this implies that $\eta = 0$ in $\tilde{J}(\mathbf{RP}^{k-1})$. Pulling back to $\widetilde{KO}(\mathbf{RP}^{k-1})$, we have $n \cdot \xi_{k-1} = 0$ in $\widetilde{KO}(\mathbf{RP}^{k-1})$, so by Adams' Theorem, n is divisible by $2^{\phi(k-1)}$. Let $n = 2^{4a+b} n_0$ where n_0 is odd and $b = 0, 1, 2$ or 3 . If $k > \rho(n) = 8a + 2^b$, then

$$\phi(k-1) \geq \phi(8a+2^b) = 4a + b + 1,$$

contradicting $2^{\phi(k-1)} \mid n$. Therefore, we have $k \leq \rho(n)$ as desired.

In a recent communication to us, I. M. James has suggested a similar proof of Theorem 2. He points out that a more general discussion of similar structures from which Theorem 2 follows may be found in $[W]$ and in $[J, \text{Sec. 7}]$.

REFERENCES

- [A₁] ADAMS, J. On the nonexistence of elements of Hopf invariant one. *Annals of Math.* 72 (1960), 20-104.
- [A₂] ——— Vector fields on spheres. *Annals of Math.* 75 (1962), 603-632.
- [A₃] ——— On the groups $J(X)$, I. *Topology* 2 (1963), 181-195.
- [A₄] ——— On the groups $J(X)$, II. *Topology* 3 (1965), 137-171.
- [A₅] ——— On the groups $J(X)$, III. *Topology* 3 (1965), 193-222.
- [A₆] ADAMS, J. and M. ATIYAH. K -theory and the Hopf invariant. *Quart. J. Math.* 17 (1966), 31-38.
- [A₇] ATIYAH, M. F. *K-Theory*. W. A. Benjamin, Inc., Reading, Massachusetts, 1967.
- [B] BORSUK, K. Drei Sätze über die n -dimensionale Euklidische Sphäre. *Fund. Math.* 20 (1933), 177-190.
- [G₁] GABEL, M. R. Generic orthogonal stably free projectives. *J. Algebra* 29 (1974), 477-488.
- [G₂] GERAMITA, A. and N. PULLMAN. A theorem of Hurwitz and Radon and orthogonal projective modules. *Proc. Amer. Math. Soc.* 42 (1974), 51-56.
- [H₁] HOPF, H. Über die Abbildungen von Sphären auf Sphären niedrigerer Dimension. *Fund. Math.* 25 (1935), 427-440.
- [H₂] ——— Ein topologischer Beitrag zur reellen Algebra. *Comment. Math. Helv.* 13 (1940), 219-239.
- [J] JAMES, I. M. *The Topology of Stiefel Manifolds*. London Math. Soc. Lecture Notes Series, Vol. 24, Cambridge University Press, 1976.
- [K₁] KNEBUSCH, M. Runde Formen über semilokalen Ringen. *Math. Ann.* 193 (1971), 21-34.
- [K₂] ——— Some open problems. In: *Proc. Quadratic Form Conference* (ed. G. Orzech), Queen's Papers in Pure and Applied Math., Vol. 55, pp. 361-370, Queen's University, Kingston, Ontario, 1977.
- [L] LAM, T. Y. *The Algebraic Theory of Quadratic Forms*. W. A. Benjamin, Inc., 1973.
- [P] PFISTER, A. Multiplikative quadratische Formen. *Arch. Math.* 16 (1965), 363-370.
- [W] WOODWARD, L. M. Vector fields on spheres and a generalization. *Quart. J. Math.* 24 (1973), 357-366.

(Reçu le 30 novembre 1980)

Z. D. Dai

Institute of Mathematics
Peking, China

T. Y. Lam

University of California
Berkeley, California 94720

R. J. Milgram

Stanford University
Stanford, California 94305

Vide-leer-empty