

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	27 (1981)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	 IDENTITIES FOR PRODUCTS OF GAUSS SUMS OVER FINITE FIELDS
Autor:	Evans, Ronald J.
Kapitel:	8. Character sum analogues of (1), (1a) and (1b)
DOI:	https://doi.org/10.5169/seals-51748

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 01.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Now replace u by tu to get

$$(28) \quad E = \sum_{u, t} \chi_1 \chi_2 \chi_3(t) \bar{\chi}_1 \bar{\chi}_2(1-t) \bar{\chi}_1(u) \chi_1 \chi_2(1-u) \{1 + \phi(ut)\} \\ = J(\chi_1 \chi_2 \chi_3, \bar{\chi}_1 \bar{\chi}_2) J(\bar{\chi}_1, \chi_1 \chi_2) + J(\chi_1 \chi_2 \chi_3 \phi, \bar{\chi}_1 \bar{\chi}_2) J(\bar{\chi}_1 \phi, \chi_1 \chi_2),$$

where the Jacobi sums J are defined above (18). Since $\chi_1^2, \chi_2^2, \chi_3^2$, and $\chi_1 \chi_2$ are nontrivial, (5) now follows from (18).

Remark. If $\chi_1 \chi_2, \chi_1 \chi_3$, or $\chi_2 \chi_3$ is trivial, we can easily evaluate E directly from its definition. Otherwise, E can be evaluated simply from (28).

8. CHARACTER SUM ANALOGUES OF (1), (1a) AND (1b)

Let $\chi_1, \chi_2, \chi_3, \phi$ be characters on $GF(q)$, where ϕ has order 2, $p > 2$. Set $t_0 = 1$. The discriminant of the polynomial

$$F(y) = \sum_{i=0}^n t_i y^{n-i}$$

is a polynomial in t_1, \dots, t_n which shall be denoted by D_n . Write

$$E_n = \sum_{i=0}^n (-1)^i t_i.$$

We conjecture that the following analogues of (1), (1a), (1b) hold for each $n \geq 1$:

$$(29) \quad \sum_{t_1, \dots, t_n \in GF(q)} \chi_1(t_n) \chi_2(E_n) \chi_3 \phi(D_n) = \prod_{j=0}^{n-1} \frac{-G(\chi_3^{j+1}) G(\chi_1 \chi_3^j) G(\chi_2 \chi_3^j)}{G(\chi_3) G(\chi_1 \chi_2 \chi_3^{n+j-1})},$$

provided that the n characters $\chi_1 \chi_2 \chi_3^{n+j-1}$ ($0 \leq j \leq n-1$) are all nontrivial;

$$(29a) \quad \sum_{t_1, \dots, t_n \in GF(q)} \chi_1(t_n) \chi_3 \phi(D_n) \zeta^{T(t_1)} = \prod_{j=0}^{n-1} \frac{-G(\chi_3^{j+1}) G(\chi_1 \chi_3^j)}{G(\chi_3)}$$

for all χ_1, χ_3 ; and

$$(29b) \quad \sum_{t_1, \dots, t_n \in GF(q)} \chi_3 \phi(D_n) \zeta^{\frac{p+1}{2} T(t_1^2 - 2t_2)} = \prod_{j=0}^{n-1} \frac{-\phi(2) G(\phi) G(\chi_3^{j+1})}{G(\chi_3)}$$

for all χ_3 .

Formulas (29), (29a), and (29b) have been verified by computer for some small primes q with $n = 3, 4$. Of course the formulas are well known for $n = 1$. For

$n = 2$, (29a) and (29b) are not hard to prove, and (29) follows from the proof of (4). (For example, if χ_3 in (29) is trivial, one makes use of (19) and (21)-(23).)

For $n = 3$, we can prove (29b), but not (29) or (29a). The *ad hoc* proof given below appears to shed little light on the general case.

THEOREM. For $n = 3$, (29b) holds.

Proof. All rational fractions below are to be interpreted as integers $(\bmod p)$; for example, $\frac{1}{2}$ equals $(p+1)/2$. We must show that

$$(30) \quad A = \sum_{t, u, v} \chi \phi(D_3) \zeta^T \left(\frac{v^2}{2} - u \right) = - \frac{\phi(2) G^3(\phi) G(\chi^3) G(\chi^2)}{G^2(\chi)}$$

for any character χ on $GF(q)$, where

$$D_3 = u^2v^2 - 4u^3 - 4tv^3 - 27t^2 + 18tuv.$$

First suppose that $p = 3$. Then

$$\begin{aligned} A &= \sum_{t, u, v} \chi \phi(u^2v^2 - u^3 - tv^3) \zeta^T \left(\frac{v^2}{2} - u \right) \\ &= \sum_{v \neq 0} \sum_u \zeta^T \left(\frac{v^2}{2} - u \right) \sum_t \chi \phi(u^2v^2 - u^3 - t) + \sum_{t, u} \phi \chi^3(-u) \zeta^T(-u) \\ &= 0 - q G(\phi \chi^3) = -q G(\chi \phi), \end{aligned}$$

since $G(\psi^p) = G(\psi)$ for any character ψ . Now (30) follows with the aid of (26).

Next, suppose that $p > 3$. Completing the square in t , one has

$$D_3/27 = c - \left(t + \frac{2v^3}{27} - \frac{uv}{3} \right)^2,$$

where $c = \frac{4}{27} \left(\frac{v^2}{3} - u \right)^3$. Thus

$$\begin{aligned} \bar{\chi} \phi(27) A &= \sum_{t, u, v} \chi \phi(c - t^2) \zeta^T \left(\frac{v^2}{2} - u \right) \\ &= \sum_{u, v} \zeta^T \left(\frac{v^2}{2} - u \right) \sum_t \chi \phi(c - t) \{1 + \phi(t)\} \\ &= \sum_{u, v} \zeta^T \left(\frac{v^2}{2} - u \right) \sum_t \chi \phi(c - t) \phi(t) \\ &= K \sum_{\substack{u, v \\ c=0}} \zeta^T \left(\frac{v^2}{2} - u \right) + J \sum_{u, v} \chi(c) \zeta^T \left(\frac{v^2}{2} - u \right), \end{aligned}$$

where $K = \chi\phi(-1) \sum_t \chi(t)$ and $J = \sum_t \chi\phi(1-t)\phi(t)$.

Thus

$$\bar{\chi}\phi(27)A = K \sum_v \zeta^T\left(\frac{v^2}{6}\right) + \chi\left(\frac{4}{27}\right) J \sum_{u,v} \chi^3\left(\frac{v^2}{3} - u\right) \zeta^T\left(\frac{v^2}{2} - u\right).$$

Replace u by $u + \frac{v^2}{3}$ to obtain

$$\begin{aligned} \bar{\chi}\phi(27)A &= -K\phi(6)G(\phi) + \chi\left(\frac{4}{27}\right) J \sum_{u,v} \chi^3(-u) \zeta^T\left(\frac{v^2}{6} - u\right) \\ &= \phi(6)G(\phi) \left\{ -K + \chi\left(\frac{4}{27}\right) J G(\chi^3) \right\}. \end{aligned}$$

If χ is trivial, then $K = \phi(-1)(q-1)$, $J = -\phi(-1)$, and $G(\chi^3) = 1$, and the desired result (30) follows. If χ is nontrivial, then $K = 0$ and

$$J = -G(\chi\phi)G(\phi)/G(\chi)$$

by (18), and (30) follows with the aid of (26).

REFERENCES

- [1] ASKEY, R. Some basic hypergeometric extensions of integrals of Selberg and Andrews. *SIAM J. Math. Anal.* 11 (1980), 938-951,
- [2] BOYARSKY, M. p -adic gamma functions and Dwork cohomology. *Trans. Amer. Math. Soc.* 257 (1980), 359-369.
- [3] DAVENPORT, H. und H. HASSE. Die Nullstellen der Kongruenzzetafunktionen in gewissen zyklischen Fällen. *J. Reine Angew. Math.* 172 (1934), 151-182.
- [4] EVANS, R., J. PULHAM and J. SHEEHAN. On the number of complete subgraphs contained in certain graphs. *J. Combinatorial Theory* (to appear).
- [5] GRAS, G. Sommes de Gauss sur les corps finis. *Publ. Math. Besançon* 1 (1977-1978), 1-71.
- [6] GROSS, B. and N. KOBLITZ. Gauss sums and the p -adic Γ -function. *Annals of Math.* 109 (1979), 569-581.
- [7] SELBERG, A. Private correspondence, Summer, 1980.
- [8] STICKELBERGER, L. Über eine Verallgemeinerung der Kreistheilung. *Math. Ann.* 37 (1890), 321-367.
- [9] THOMASON, A. Ph.D. Thesis, Cambridge University, 1979.

(Reçu le 18 septembre 1980)

Ronald J. Evans

Department of Mathematics
University of California, San Diego
La Jolla, CA 92093